

**Question 1 (12 Marks) Begin a SEPARATE sheet of paper**


	<b>Marks</b>
(a) Evaluate $\frac{e+1}{\pi}$ , correct to three decimal places.	<b>2</b>
(b) Find $\theta$ , to the nearest degree, if $\sin \theta = \frac{4 \sin 57^\circ}{6.7}$	<b>2</b>
(c) What is the centre and radius of a circle with equation $(x+2)^2 + (y-3)^2 = 2.25$ .	<b>2</b>
(d) The mean of 3, 5, 7, $x$ is 6.75. What is the value of $x$ ?	<b>1</b>
(e) If $x = 2.35$ , evaluate the expression $ -3-4x $	<b>1</b>
(f) Factorise $3x^2 - 5x - 2$ .	<b>1</b>
(g) Express 2.32 radians as an angle in degrees, correct to the nearest minute.	<b>1</b>
(h) Write down the domain and range for $y = \sqrt{x^2 - 9}$ .	<b>2</b>

**Question 2 (12 Marks) Begin a SEPARATE sheet of paper**

	<b>Marks</b>
(a) Differentiate the following:	
(i) $e^{0.5}$ .	<b>1</b>
(ii) $\log_e (x^2 - 3x)$ .	<b>1</b>
(iii) $\frac{2x-1}{\cos x}$	<b>1</b>
(b) Integrate: $\int \sin \frac{x}{2} dx$	<b>2</b>
(c) Evaluate: $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{5}{6} e^{3x} dx$ (Leave your answer in exact form).	<b>2</b>
(d) Given $\log_m p = 1.75$ and $\log_m q = 2.25$ . Find the following:	
(i) $\log_m pq$	<b>1</b>
(ii) $\log_m \frac{q}{p}$	<b>1</b>
(iii) $\sqrt[3]{pq}$ in terms of $m$ .	<b>3</b>

**Question 3 (12 Marks) Begin a SEPARATE sheet of paper****Marks**

The number plane shows the line  $n$  with equation  $x + 2y = 5$ .  
The point  $P$  is  $(2, 4)$  and  $O$  is the origin.




- |  |          |
|--|----------|
| (a) Find the coordinates of the midpoint $M$ , of the interval $OP$ .  | <b>1</b> |
| (b) Show that the point $M$ lies on the line $n$ .   | <b>2</b> |
| (c) Find the gradient of the line $OP$ .   | <b>1</b> |
| (d) Show that the line $n$ is the perpendicular bisector of the interval $OP$ .                                | <b>3</b> |
| (e) Line $n$ meets the $x$ -axis at $Q$ . Find the co-ordinates of $Q$ .                                       | <b>1</b> |
| (f) A line is drawn through $O$ parallel to $PQ$ and it meets line $n$ in $R$ . Find the co-ordinates of $R$ . | <b>2</b> |
| (g) What sort of quadrilateral is $PQOR$ ? Give reasons for your answer.                                       | <b>2</b> |

**Question 4 (12 Marks) Begin a SEPARATE sheet of paper**

	<b>Marks</b>
(a) Let $m$ and $n$ be positive whole numbers where $m > n$ .	
(i) Show that a triangle with sides $m^2 + n^2$ , $m^2 - n^2$ , $2mn$ obeys Pythagoras' Theorem.	<b>2</b>
(ii) Which Pythagorean Triad is generated when $m = 3$ and $n = 2$ ?	<b>1</b>
(b) Consider the function $f(x) = x - 2 \log_e x$ , for $x > 0$ .	
(i) Find the first <b>and</b> second derivative of $f(x)$ .	<b>2</b>
(ii) Find the co-ordinates of the turning point(s) and determine their nature.	<b>2</b>
(iii) Show that there are no points of inflexion.	<b>2</b>
(iv) What is the maximum value of $x - 2 \log_e x$ in the domain $1 \leq x \leq 5$ ?	<b>1</b>
(v) Sketch the curve $y = x - 2 \log_e x$ , for $0 < x \leq 5$ .	<b>2</b>

**Question 5 (12 Marks) Begin a SEPARATE sheet of paper****Marks**

(a)



The diagram above shows the graph of  $y = \sin x$  for  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .

- (i) Copy the diagram onto your exam paper. 2  
 On the same set of axes, graph  $y = \cos 2x$  for  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .
- (ii) Show that the two graphs intersect where  $x = \frac{\pi}{6}$ . 1
- (iii) Calculate the exact area enclosed between the two curves. 3
- (b) Show that  $\frac{\sec \theta - \sec \theta \cos^4 \theta}{1 + \cos^2 \theta} = \sin \theta \tan \theta$ . 3
- (c) (i) Find the value(s) of  $k$  for which  $x^2 + (2 - k)x + 2 \cdot 25 = 0$  has equal roots. 2  
 (ii) Find the value(s) of  $k$  for which  $y = kx + 1$  is a tangent to  $y = x^2 + 2x + 3 \cdot 25$ . 1

**Question 6 (12 Marks) Begin a SEPARATE sheet of paper**

- |   | <b>Marks</b> |
|---|--------------|
| (a) Boat $A$ is 15km from point $P$ on a bearing of $055^{\circ}$ T.<br>Boat $B$ is 25km from point $P$ on a bearing of $135^{\circ}$ T.  |              |
| (i) Draw a diagram showing the information above and find the angle $APB$ .   | <b>2</b>     |
| (ii) Calculate, to one decimal place, the distance that the two boats are apart.  | <b>2</b>     |
| <br>  |              |
| (b) Evaluate $\sum_{r=3}^7 2^r - 3r$ .  | <b>2</b>     |
| <br>  |              |
| (c) Water enters an empty container where the rate of filling is $R = 6(e^t + e^{-t})$ where $R = \frac{dV}{dt}$ in litres per minute. The volume ( $V$ ) of water in the vessel is in litres and time ( $t$ ) is in minutes. |              |
| (i) What is the initial rate of filling?  | <b>1</b>     |
| (ii) Find $V$ in terms of $t$ .   | <b>2</b>     |
| (iii) Show that, when the container holds 5 litres, then $6e^{2t} - 5e^t - 6 = 0$ .   | <b>1</b>     |
| (iv) Hence, find to the nearest second, the time taken for the volume to reach 5 litres.  | <b>2</b>     |

**Question 7 (12 Marks) Begin a SEPARATE sheet of paper**

**Marks**


(a) Find  $y$  if  $\frac{dy}{dx} = \frac{3\sec^2 x}{\tan x}$  and  $y = 4$  when  $x = \frac{\pi}{4}$ . **3**

(b) *Twinkle Finance* offers its investors the opportunity to have interest credited to their investment “as often as you wish”. Naturally, many investors decide for the “EVERY MINUTE” plan in which *Twinkle* offers 12% pa, compounded each minute.

(i) Tanya invests \$1000 for a year with *Twinkle* on the “EVERY MINUTE” plan. Theoretically, *Twinkle*’s computers multiply Tanya’s balance by approximately 1.000 000 228 every minute. Show why this is so. **2**

(ii) How much, to the nearest dollar, is Tanya’s investment worth after 1 year? **1**

(c)



In the diagram above, an equilateral triangle  $PQR$  is inscribed in a circle  $C_1$ , with centre  $O$ , and radius 5 units.  $\angle POQ = \angle QOR = \angle POR$ .

An arc of another circle  $C_2$ , with centre  $P$ , and radius  $5\sqrt{3}$  units, intersects  $C_1$  at the points  $Q$  and  $R$ , as shown in the diagram.

(i) Show  $\angle QPO = \frac{\pi}{6}$ , giving reasons. **1**

(ii) Find the area of the sector  $QPR$ , of the circle  $C_2$ . **1**

(iii) Find the area of the sector  $QOR$ , of the circle  $C_1$ . **1**

(iv) Hence, or otherwise, show that the area of the shaded region is  $25 \left( \frac{\sqrt{3}}{2} - \frac{\pi}{6} \right)$  units<sup>2</sup>. **3**


**Question 8 (12 Marks) Begin a SEPARATE sheet of paper**

- |  | <b>Marks</b> |
|--|--------------|
| (a) Let $A$ be the point $(-2, 0)$ and $B$ be the point $(6, 0)$ .<br>At $P(x, y)$ , $PA$ meets $PB$ at right angles.  |              |
| (i) Show that the gradient of $PA$ is $\frac{y}{x+2}$ .  | <b>1</b>     |
| (ii) Find the equation of the locus of $P$ <b>and</b> give a geometric description of the locus.   | <b>3</b>     |
|  |              |
| (b) The velocity of an object is given by the equation $v = 6t - 8 - t^2$ , where time ( $t$ ) is in seconds and velocity ( $v$ ) is in metres/second. Initially, the object is 5 metres to the right of the origin. |              |
| (i) Find an equation for the displacement of the object.   | <b>2</b>     |
| (ii) Find when the object is momentarily at rest.  | <b>1</b>     |
| (iii) Find the distance travelled by the object in the first four seconds.   | <b>2</b>     |
|  |              |
| (c) Two dice are biased so that, the probability of throwing a six face up is $\frac{3}{8}$ and the probability of throwing any other number is $\frac{1}{8}$ .  |              |
| Find the probability of:   |              |
| (i) Rolling a double six face up.  | <b>1</b>     |
| (ii) Rolling the dice so that neither is a six face up.  | <b>1</b>     |
| (iii) Only one six appearing face up when the two dice are rolled.   | <b>1</b>     |

**Question 9 (12 Marks) Begin a SEPARATE sheet of paper****Marks**

- (a) (i) Write  $x^2 - 6x + 8 = 2y$  in the form  $(x - h)^2 = 4a(y - k)$ . 2
- (ii) Hence, state the co-ordinates of focus for the parabola. 1

(b)




The graph of  $y = f'(x)$  is shown above. The zeros of  $f'(x)$  are  $x = -2$ ,  $0.5$ , and  $3$ .  
Point  $C$  has  $x$  co-ordinate of  $-0.9$  and point  $B$  has  $x$  co-ordinate of  $1.9$ .

- (i) For what range of values of  $x$  is  $f(x)$  increasing? 1
- (ii) Point  $C$  is the local maximum of  $f'(x)$ . 2  
What type of point occurs on  $y = f(x)$  when  $x = -0.9$ . Justify your answer.
- (iii) For what range of values of  $x$  is  $f(x)$  concave down? 1

**Question 9(c) continued on next page**

9(c)



The graphs of  $y = \sin \frac{\pi}{2} x$  and  $y = x^2$  are shown above.

The region bounded by these curves and the  $x$ -axis is shaded.

- (i) Show that the two curves meet where  $x = 1$ . 1
- (ii) Write the definite integral(s) which will give the volume of the solid of revolution when this region is rotated about the  $x$ -axis. **DO NOT EVALUATE THESE INTEGRALS** 1
- (iii) Use Simpson's Rule with five function values to approximate this volume, leaving your answer in terms of  $\pi$ . 3

**Question 10 (12 Marks) Begin a SEPARATE sheet of paper****Marks**

- (a) The percentage of Carbon14 in an organism falls exponentially after the death of the organism. After 1845 years 80% of the original concentration of Carbon14 remains. Using the equation  $C = C_0 e^{-kt}$  to represent the exponential fall of Carbon14:

- (i) Find the exact value of  $k$ . 2
- (ii) An artefact containing this organism has 65% of the original concentration of Carbon14. How long has this organism been dead? Give answer to the nearest year. 2
- (iii) A sea sponge containing this organism has been dead for 12 000 years. What percentage (to 1 decimal place) of the original Carbon14 concentration does it have? 2

- (b) Two sailors are paid to bring a motor launch back to Sydney from Gilligan's Island, a total distance of 1 200 km. They are each paid \$25 per hour for the time spent at sea.

The launch uses fuel at a rate of  $20 + \frac{v^2}{10}$  litres per hour where the speed of the launch is  $v$  km

per hour. Diesel fuel costs \$1.25 per litre.

- (i) Show that, to bring the launch back from Gilligan's Island, the total cost (\$ $C$ ) to the owners is 3

$$C = \frac{90\,000}{v} + 150v.$$

- (ii) Find the speed that minimises the cost and determine this cost, giving your answer to the nearest dollar. 3

**END OF EXAMINATION**

SOLUTIONS TO 2 UNIT TRIAL MATHEMATICS

Question 1 (12 marks)

a)  $\frac{e+1}{\pi} = 1.183565\dots$   
 $\frac{e+1}{\pi} = 1.184$  (3 dp)

b)  $\sin \theta = 0.500698846$   
 $\theta = 30^\circ 3'$   
 $\theta = 30^\circ$  (nearest degree)

c) Centre  $(-2, 3)$   
 radius =  $\sqrt{2.25} = 1.5$

d)  $\frac{3+5+7+x}{4} = 6.75$   
 $15+x = 27$   
 $x = 12$

e)  $|-3 - 4 \times 2.35| = |-12.4| = 12.4$

f)  $3x^2 - 5x - 2 = (3x+1)(x-2)$

g)  $2.32 \text{ rads} = 2.32 \times \frac{180^\circ}{\pi} = 132^\circ 56'$

h)  $x^2 - 9 \geq 0$   
 D:  $x \leq -3$  or  $x \geq 3$   
 R:  $y \geq 0$

Question 2 (12 marks)

a) (i) 0  
 (ii)  $\frac{2x-3}{x^2-3x}$   
 (iii)  $\frac{2 \cos x + (2x-1) \sin x}{\cos^2 x}$

b)  $-2 \cos \frac{x}{2} + c$


c)  $\int_0^1 \frac{5}{6} e^{3x} dx = \frac{5}{6} \left[ \frac{1}{3} e^{3x} \right]_0^1$   
 $= \frac{5}{18} (e^3 - 1)$

d) (i)  $\log pq = \log p + \log q = 1.75 + 2.25 = 4$

(ii)  $\log \frac{q}{p} = \log q - \log p = 2.25 - 1.75 = 0.5$

(iii) Let  $\sqrt[5]{pq^2} = y \Rightarrow \log_m \sqrt[5]{pq^2} = \log_m y$   
 $\frac{1}{5} (\log p + 2 \log q) = \log_m y$   
 $\frac{1}{5} (1.75 + 2 \times 2.25) = \log_m y$   
 $1.25 = \log_m y$   
 $\therefore m^{1.25} = y$

Question 3 (12 marks)



a) M is  $(1, 2)$   
 b) Sub.  $M(1, 2)$  into  $x+2y=5$ :  
 LHS =  $1 + 2 \times 2 = 5$   
 RHS = 5  
 $\therefore$  LHS = RHS  
 $\therefore M(1, 2)$  lies on  $x+2y=5$

c)  $m_{op} = \frac{4}{2} = 2$

d)  $m_{line n} = -\frac{2 \frac{1}{2}}{5} = -\frac{1}{2}$   
 $m_{op} \times m_{line n} = 2 \times -\frac{1}{2} = -1$

$\therefore OP \perp$  line  $n$   
 Also  $(1, 2)$  is the midpoint of  $OP \therefore$  bisector  
 $\therefore$  line  $n$  is the perpendicular bisector of  $OP$ .

e)  $Q(5, 0)$

f)  $m_{PQ} = -\frac{4}{3}$   
 $m_{OR} = m_{PQ} = -\frac{4}{3}$  ( $PQ \parallel OR$ )  
 By congruent triangles, and using diagram above,  
 $R$  is  $(-3, 4)$

g)  $OQ \parallel PR$  (gradients 0)  
 $PQ \parallel OR$  (from (f))  
 Diagonals meet at  $90^\circ$   
 $\therefore PQOR$  is a rhombus (2 pairs of opposite sides parallel & diagonals meet)

-1 if not properly set out.

NB: Other reasons are also possible.

(3)

Question 4 (12 marks)

a) (i) Required to prove  $(m^2-n^2)^2 + (2mn)^2 = (m^2+n^2)^2$

$$\begin{aligned} \text{LHS: } & (m^2-n^2)^2 + (2mn)^2 \\ &= m^4 - 2m^2n^2 + n^4 + 4m^2n^2 \\ &= m^4 + 2m^2n^2 + n^4 \\ &= (m^2+n^2)^2 \\ &= \text{RHS.} \end{aligned}$$

∴ LHS = RHS

(ii) If  $m=3, n=2$ : triad is 13, 5, 12

b) (i)  $f(x) = x - 2 \log_e x, x > 0$

$$f'(x) = 1 - \frac{2}{x} \text{ or } 1 - 2x^{-1}$$

$$f''(x) = 2x^{-2} \text{ or } \frac{2}{x^2}$$

(ii) For t.p. ( $f'(x) = 0$ ):  $1 - \frac{2}{x} = 0$

$$x - 2 = 0$$

$$x = 2$$

∴ Possible t.p. is  $(2, 2 - 2 \ln 2)$

To determine its nature use either 1st or 2nd derivative

Using 1st derivative:

$x$	1	2	3
$f'(x)$	-1	0	$\frac{1}{3}$

∴ local minimum at  $x=2$

Using 2nd derivative: at  $x=2$

$$f''(2) = 2 \times 2^{-2} > 0$$

∴ concave up

∴ local minimum at  $x=2$

(iii) For possible points of inflexion,  $f''(x) = 0$

$$\therefore \frac{2}{x^2} = 0$$

$$\text{but } \frac{2}{x^2} \neq 0$$

∴ there are no points of inflexion.


(4)

(iv) at  $x=1, f(1) = 1 - 2 \ln 1 = 1$

at  $x=5, f(5) = 5 - 2 \ln 5 \approx 1.78$


∴ Maximum value is  $5 - 2 \ln 5$

(v)  $y = x - 2 \ln x, 0 < x \leq 5$



Question 5 (12 marks)

a) (i)



(ii) at  $x = \frac{\pi}{6}, \sin \frac{\pi}{6} = \frac{1}{2}$

$$\cos 2 \times \frac{\pi}{6} = \cos \frac{\pi}{3} = \frac{1}{2}$$

∴  $\sin x = \cos 2x$  at  $x = \frac{\pi}{6}$

(iii)  $A = \int_{-\frac{\pi}{2}}^{\frac{\pi}{6}} (\cos 2x - \sin x) dx$

$$= \left[ \frac{1}{2} \sin 2x + \cos x \right]_{-\frac{\pi}{2}}^{\frac{\pi}{6}}$$

$$= \left( \frac{1}{2} \sin \frac{\pi}{3} + \cos \frac{\pi}{6} \right) - \left( \frac{1}{2} \sin(-\pi) + \cos\left(-\frac{\pi}{2}\right) \right)$$

$$= \frac{\sqrt{3}}{4} + \frac{\sqrt{3}}{2} = \frac{3\sqrt{3}}{4} \text{ units}^2$$

1/2  
1/2  
1

1/2 shape  
1/2 for minimum  
1/2 for (1,1) & (5, 5-2ln5)  
1/2 for scale

1 for shape  
1 for scale

1/2  
1/2

1  
1

Question 5 continued.

$$\begin{aligned}
 \text{b) LHS} &= \frac{\sec \theta - \sec \theta \cos^4 \theta}{1 + \cos^2 \theta} \\
 &= \frac{\sec \theta (1 - \cos^4 \theta)}{1 + \cos^2 \theta} \\
 &= \frac{\sec \theta (1 + \cos^2 \theta)(1 - \cos^2 \theta)}{1 + \cos^2 \theta} \\
 &= \frac{1}{\cos \theta} \cdot \sin^2 \theta \\
 &= \frac{\sin \theta}{\cos \theta} \cdot \sin \theta \\
 &= \tan \theta \cdot \sin \theta \\
 &= \text{RHS}
 \end{aligned}$$

[3]


$$\begin{aligned}
 \text{c) (i) } \Delta &= (2-k)^2 - 4 \cdot 1 \cdot (2 \cdot 25) \\
 &= (2-k)^2 - 9 \\
 \text{For equal roots, } \Delta &= 0 \\
 \therefore (2-k)^2 - 9 &= 0 \\
 (2-k)^2 &= 9 \\
 2-k &= \pm 3 \\
 \underline{k = 5 \text{ or } -1}
 \end{aligned}$$

[2]

$$\begin{aligned}
 \text{(ii) } y &= kx + 1 \\
 y &= x^2 + 2x + 3.25 \\
 \text{Solve: } x^2 + 2x + 3.25 &= kx + 1 \\
 x^2 + (2-k)x + 2.25 &= 0 \\
 \text{For line to be a tangent, } \Delta &= 0, \text{ then from (i)} \\
 k &= 5 \text{ or } -1.
 \end{aligned}$$

[1]

Question 6 (12 marks)



[2]

$$\begin{aligned}
 \text{(ii) } AB^2 &= 15^2 + 25^2 - 2 \cdot 15 \cdot 25 \cdot \cos 80^\circ \\
 &= 719.76 \\
 \underline{AB} &= \underline{26.8 \text{ km}}
 \end{aligned}$$

[2]

$$\begin{aligned}
 \text{b) } \sum_{r=3}^7 2^r - 3r &= 2^3 - 3 \times 3 + 2^4 - 3 \times 4 + 2^5 - 3 \times 5 \\
 &\quad + 2^6 - 3 \times 6 + 2^7 - 3 \times 7 \\
 &= (2^3 + 2^4 + 2^5 + 2^6 + 2^7) \\
 &\quad - 3(3 + 4 + 5 + 6 + 7) \\
 &= 248 - 75 = 173
 \end{aligned}$$

[2]

$$\text{c) (i) When } t=0, R = 6(e^0 + e^0) = 12 \text{ L/min}$$

[1]

$$\begin{aligned}
 \text{(ii) } V &= \int 6(e^t + e^{-t}) dt \\
 V &= 6(e^t - e^{-t}) + c \\
 \text{When } t=0, V=0 &\implies 0 = 6(e^0 - e^0) + c \\
 \underline{c} &= \underline{0}
 \end{aligned}$$

[2]

$$\begin{aligned}
 \text{(iii) } 5 &= 6(e^t - \frac{1}{e^t}) \\
 5e^t &= 6e^{2t} - 6 \\
 \therefore 6e^{2t} - 5e^t - 6 &= 0
 \end{aligned}$$

[1]

$$\begin{aligned}
 \text{(iv) } (2e^t - 3)(3e^t + 2) &= 0 \\
 e^t = \frac{3}{2} \text{ or } e^t = -\frac{2}{3} \\
 \text{but } e^t \neq -\frac{2}{3} \therefore \text{no sol}^n. \\
 \therefore t = \ln \frac{3}{2} \doteq \underline{24 \text{ seconds}}
 \end{aligned}$$

[2]

for diagram for  $\angle APB = 80^\circ$

-1/2 if not to 1 dp.

NB: It is not intended that rules for sum of AP & GP's be used here.

-1/2 if no units are provided.

-1/2 if not to nearest second.

Question 7 (12 marks)

a) dy/dx = 3 sec^2 x / tan x

y = 3 ln(tan x) + c

When x = pi/4, y = 4 => 4 = 3 ln(tan pi/4) + c

c = 4

y = 3 ln(tan x) + 4 [2]

b) (i) 1 year = 365 x 24 x 60 minutes = 525600 minutes

Incremental rate of interest = 12 / 525600 % per minute

= 0.12 / 525600 per minute

= 0.00002283 per min. 1/2

Since interest is added onto the amount invested then interest is multiplied by approximately

1.00002283 every minute [2] 1/2

(ii) 1000 (1.00002283)^525600 = \$1127 (nearest dollar) [1] 1

c) (i) LQOP = 2pi/3 (angle sum about a point is 360) LQPO = (2pi - 2pi/3) / 2 = pi/6 (angle sum of triangle is 360; angles opposite equal sides are equal) [2] 1

(ii) A\_sector QPR = 1/2 x 5sqrt(3) x 5sqrt(3) x pi/3 = 25pi/2 unit^2 [1] 1

(iii) A\_sector QOR = 1/2 x 5 x 5 x 2pi/3 = 25pi/3 units^2 [1] 1

(iv) There are several ways to do this which may involve areas of segments/triangles/sectors.

PTO.

Question 7 (continued)

Method 1

(iv) A\_segment QCR = 1/2 x (5sqrt(3))^2 (pi/3 - sin pi/3)

= 25 x 3/2 (pi/3 - sqrt(3)/2)

= 25 (pi/2 - 3sqrt(3)/4)

A\_segment QCR = 1/2 x 5^2 (2pi/3 - sin 2pi/3)

= 25 x 1/2 (2pi/3 - sqrt(3)/2)

= 25 (pi/3 - sqrt(3)/4)

A\_shaded = 25 (pi/3 - sqrt(3)/4) - 25 (pi/2 - 3sqrt(3)/4)

= 25 (sqrt(3)/2 - pi/6) units^2 [3] 1

Alt. Method 2


Shaded area = (iii) - (ii) + 2 A\_s

= 25pi/3 - 25pi/2 + 2 x 25sqrt(3)/4

= 25 (sqrt(3)/2 - pi/6) [3] 1

Question 8 (12 marks)

a)



m\_PA = (y-0)/(x-(-2)) = y/(x+2)

[1] 1

(ii) m\_PA x m\_PB = -1

y/(x+2) x y/(x-6) = -1

y^2 = -(x+2)(x-6)

= -x^2 + 4x + 12

x^2 - 4x + 4 + y^2 = 12 + 4

(x-2)^2 + y^2 = 16

∴ Locus is a circle, centre (2,0), radius 4 units. [2] 1

9

Question 8 (Continued)

b)  $v = 6t - 8 - t^2 = \frac{dx}{dt}$

(i)  $x = 3t^2 - 8t - \frac{t^3}{3} + c$

When  $t=0, x=+5 \Rightarrow 5 = c$

$\therefore x = 3t^2 - 8t - \frac{t^3}{3} + 5$

(ii) At rest,  $v=0 \Rightarrow 0 = 6t - 8 - t^2$

$0 = (t-4)(t-2)$

$\therefore$  at rest when  $t=2$  and  $t=4$

(iii)  $x(0) = 5$

$x(2) = -1\frac{2}{3}$   $\therefore$  In first 2 secs it travels  $5 + 1\frac{2}{3} = 6\frac{2}{3}m$

$x(4) = x = -\frac{1}{3}$   $\therefore$  from  $t=2$  to  $t=4$  it travels  $1\frac{1}{3}$  units.

$\therefore$  distance travelled =  $5 + 1\frac{2}{3} + 1\frac{1}{3} = 8$  units

c) (i)  $P(6,6) = \frac{3}{8} \times \frac{3}{8} = \frac{9}{64}$

(ii)  $P(\tilde{6}, \tilde{6}) = \frac{5}{8} \times \frac{5}{8} = \frac{25}{64}$

(iii)  $P(6, \tilde{6}) + P(\tilde{6}, 6) = \frac{3}{8} \times \frac{5}{8} \times 2 = \frac{30}{64} = \frac{15}{32}$

Question 9 (12 marks)

a) (i)  $x^2 - 6x + 8 = 2y$

$x^2 - 6x + 9 - 1 = 2y$


$(x-3)^2 = 2y + 1$

$= 2(y + \frac{1}{2})$

$(x-3)^2 = 4(\frac{1}{2})(y - (-\frac{1}{2}))$

(ii) Vertex =  $(3, -\frac{1}{2})$   $a = \frac{1}{4}$

Focus =  $(3, 0)$



10

Question 9 (Continued)

b) (i) For increasing function,  $f'(x) > 0$

ie  $\{x: -2 < x < \frac{1}{2}\}, \{x: x > 3\}$

(ii)  $f'(-0.9)$  is a maximum, so  $f''(-0.9) = 0$

$\therefore C$  is a possible point of inflexion.

for  $x < -0.9, f'(x) > 0$

for  $x > -0.9, f'(x) > 0$

$\therefore$  In the neighbourhood of  $-0.9$ , gradients  $> 0$ .

$\therefore C$  is a point of inflexion.

(iii) For concave down,  $f'(x)$  is decreasing.

$\therefore \{x: -0.9 < x < 1.9\}$

c) (i) at  $x=1, \sin \frac{\pi}{2} \times 1 = \sin \frac{\pi}{2} = 1$

$1^2 = 1$

$\therefore \sin \frac{\pi}{2} x = x^2$  at  $x=1$

(ii)  $V = \pi \int_0^1 (x^2)^2 dx - \pi \int_1^2 (\sin \frac{\pi}{2} x)^2 dx$

OR

$V = \pi \int_0^1 x^4 dx - \pi \int_1^2 \sin^2 \frac{\pi}{2} x dx$

(iii) 

x	Weight	$f(x)^2$
0	1	0
$\frac{1}{2}$	4	$\frac{1}{16}$
1	2	1
$1\frac{1}{2}$	4	$\sin^2 \frac{3\pi}{4} = \frac{1}{2}$
2	1	$\sin^2 \pi = 0$

0

$\frac{1}{2}$

1

$1\frac{1}{2}$

2

1

0

$V = \pi \times \frac{2-0}{12} [0 + 0 + 4(\frac{1}{16} + \frac{1}{2}) + 2(1)]$

$= \frac{\pi}{6} [\frac{17}{4}]$

$= \frac{17\pi}{24} u^3$

-1/2 if either is missing

(11)

## Question 10 (12 Marks)

a) (i) When  $t=0$ ,  $A=A_0$ When  $t=1845$ ,  $A=0.8(A_0)$ 

$$\therefore 0.8 A_0 = A_0 e^{-1845k}$$

$$0.8 = e^{-1845k}$$

$$\ln 0.8 = -1845k$$

$$k = -\frac{\ln 0.8}{1845} \text{ or } \frac{\ln 1.25}{1845} \quad [2]$$

(ii)  $A=0.65A_0$  when  $t=T$ 

$$\therefore 0.65 A_0 = A_0 e^{+\frac{\ln 0.8}{1845} T}$$

$$\ln 0.65 = \frac{\ln 0.8}{1845} T$$

$$T = \frac{1845 \times \ln 0.65}{\ln 0.8}$$

$$T = 3562 \text{ years} \quad [2]$$

(iii) When  $t=12000$ :

$$A = A_0 e^{+12000 \times \frac{\ln 0.8}{1845}}$$

$$A \doteq 0.234 A_0$$

$\therefore A \doteq 23.4\%$  of original amount is left [2]

b)  $S = \frac{D}{T}$ (i)  $T = \frac{D}{v} = \frac{1200}{v}$  hours

$$\text{Cost of wages} = 25 \times 2 \times \frac{1200}{v} = \frac{60000}{v}$$

$$\text{Cost of fuel} = \left(20 + \frac{v^2}{10}\right) \times \frac{1200}{v} \times 1.25$$

$$= \left(20 + \frac{v^2}{10}\right) \times \frac{1500}{v}$$

$$= \frac{30000}{v} + 150v$$

$$\text{Total cost, } C = \frac{60000}{v} + \frac{30000}{v} + 150v$$

$$C = \frac{90000}{v} + 150v \quad [3]$$

(12)

## Question 10 (continued)

$$(ii) C = \frac{90000}{v} + 150v$$

$$\frac{dC}{dv} = 150 - 90000v^{-2}$$

For min. cost,  $\left(\frac{dC}{dv} = 0\right)$ :

$$150 - \frac{90000}{v^2} = 0$$

$$150v^2 - 90000 = 0$$

$$v^2 - 600 = 0$$

$$v = \sqrt{600} \quad (v > 0)$$

$$\therefore v = 10\sqrt{6}$$

check  $C(10\sqrt{6})$  is a minimum:

$$\frac{d^2C}{dv^2} = 180000v^{-3}$$

$$\text{at } v=10\sqrt{6}, \quad \frac{d^2C}{dv^2} = \frac{180000}{(10\sqrt{6})^3} > 0$$

 $\Rightarrow$  concave up $\therefore$  local min at  $v=10\sqrt{6}$ .

(May use 1st derivative to test nature of stationary point.)

$$\therefore \text{Minimum cost} = \frac{90000}{\sqrt{600}} + 150 \times \sqrt{600}$$

$$= \underline{\underline{\$7348}} \quad [3]$$

END OF SOLUTIONS