



# THE KING'S SCHOOL

---

2004  
Higher School Certificate  
Trial Examination

## Mathematics

### General Instructions

- Reading time – 5 minutes
- Working time – 3 hours
- Write using black or blue pen
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- All necessary working should be shown in every question

### Total marks – 120

- Attempt Questions 1-10
- All questions are of equal value



# THE KING'S SCHOOL

---

## 2004 Higher School Certificate Trial Examination

### Mathematics

Question	Algebra and Number	Geometry	Functions	Trigonometry	Differential Calculus	Integral Calculus	Total
1	(c), (d), (f)		(b)	(a)		(e)	12
2			(b)		(a)		12
3		(c)	(b)			(a)	12
4			(c)	(a)			12
5					(a)	(b)	12
6			(c)	(a)		(b)	12
7			(a)		(c)	(b)	12
8	(a)				(b)		12
9	(a)	(b)					12
10	(i)			(ii), (iii), (v)	(iv)		12
Marks	20	10	26	20	26	18	120

**Total marks – 120**  
**Attempt Questions 1-10**  
**All questions are of equal value**

Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.

---

**Marks**

**Question 1 (12 marks)** Use a SEPARATE writing booklet.

- (a) Find the value of  $\sin 2x$  if  $x = 0.006$ , correct to 2 significant figures. **2**
- (b) State the domain and range of the function  $y = \log_e x$  **2**
- (c) Find  $\lim_{x \rightarrow 0} \frac{x^3 - 3x}{6x}$  **2**
- (d) Solve the equation  $(2x + 3)^2 = 4$  **2**
- (e) Find a primitive function of  $(2x + 3)^4$  **2**
- (f) For what value of  $x$  do  $y = 1 - 2x$  and  $2y = 7 + 6x$  hold simultaneously? **2**


**End of Question 1**

**Question 2 (12 marks)** Use a SEPARATE writing booklet.

**Marks**

- (a) Find the gradient of the normal to the curve  $y = \frac{x}{x^2 + 1}$  at the point  $(0, 0)$  **3**

(b)



In the diagram, ABCD is a rectangle where  $A = (-4, 8)$  and  $B = (2, 0)$ .  $P(3, 7)$  is a point on the side CD.

- (i) Find the gradient of line AB. **1**
- (ii) Deduce that the equation of line AB is  $4x + 3y - 8 = 0$  **2**
- (iii) Hence, or otherwise, show that the length of side BC is 5 units. **2**
- (iv) Write down the mid-point of side AB and deduce that  $P(3, 7)$  is the mid-point of side CD. **2**
- (v) Write down the point of intersection of the diagonals AC and BD. **1**
- (vi) Find the coordinates of point D. **1**

**End of Question 2**

(a) Find the exact value of


(i)  $\int_0^{\frac{\pi}{6}} \sin x \, dx$  2

(ii)  $\int_0^1 \frac{6x^2}{x^3 + 1} \, dx$  2

(b) For what values of  $k$  is  $x^2 + 2x + k$

(i) concave upward? 1

(ii) positive for all values of  $x$ ? 2



In the diagram,  $AB \parallel CD$  and  $ECP$  is a straight line.

$$\angle ABC = 50^\circ, \angle ECB = 170^\circ, \angle QPC = 140^\circ$$

(i) Find  $\angle BCP$ , giving reasons. 1

(ii) Find  $\angle ECD$ , giving reasons. 2

(iii) Prove that  $PQ \parallel AB$  2

**End of Question 3**

**Question 4 (12 marks)** Use a SEPARATE writing booklet.

**Marks**

---

- (a) (i) Sketch the curve  $y = \tan \frac{x}{2}$  for  $-2\pi \leq x \leq 2\pi$  **2**
- (ii) State the period of  $y = \tan \frac{x}{2}$  **1**
- (iii) Solve the equation  $\tan \frac{x}{2} = 1$  for  $-2\pi \leq x \leq 2\pi$  **2**
- (b) Consider the two arithmetic series
- $A = 11 + 13 + 15 + \dots$
- and  $B = -14 - 11 - 8 - \dots$
- (i) Find the sum of the first 40 terms of series  $A$  **1**
- (ii) The two series have the same number of terms and the same sum. How many terms are in the series? **3**
- (c) Find the values of  $a$ ,  $b$ ,  $c$  if
- $a(x+1)^2 + b(x+1)(x-1) + c(x-1) \equiv x^2 + 7x$  **3**

**End of Question 4**

(a) Consider the function  $f(x) = x^4 - 6x^2 + 8x$

(i) Show that  $f'(x) = 4(x+2)(x-1)^2$  **2**


(ii) For what values of  $x$  is the function increasing? **2**

(iii) Show that there is a minimum turning point at  $x = -2$  **2**

(iv) Show that there is a horizontal point of inflection at  $x = 1$  **2**

(b) (i) Briefly explain why Simpson's Rule gives the exact value of  $\int_a^b f(x) dx$  if  $f(x)$  is a quadratic function. **1**


(ii)



A parabola  $y = f(x)$  passes through the points  $(0,0)$ ,  $(1,-2)$  and  $(2,5)$ . Find the value of  $\int_0^2 f(x) dx$  **3**

**End of Question 5**


(a)



In the diagram,  $P(3,4)$  and  $Q(0,-5)$  are points on the circle  $x^2 + y^2 = 25$ .  
Let  $\angle POQ = \theta$ , as marked on the diagram.

- (i) Find the length of chord  $PQ$ . **1**
  
- (ii) Show that  $\cos \theta = -\frac{4}{5}$  **2**
  
- (iii) Find the area of minor sector  $POQ$  correct to 1 decimal place. **2**

(b)



In the diagram, the shaded region is bounded by the curve  $y = 3x^2 + \frac{3}{x^2}$ ,  $x > 0$ ,  
and the two lines  $y = 6$  and  $x = 3$ . Find the area of this shaded region. **4**

**Question 6 (continued)**

---

(c) The directrix of a parabola is the  $x$  axis and the focus is the point  $(0,4)$ .

(i) Write down the focal length of the parabola.


**1**

(ii) Find the equation of the parabola.

**2**


**End of Question 6**

- (a)  $A(-2,0)$  and  $B(0,4)$  are two points in the number plane.  $P(x,y)$  is any point such that  $AP$  is perpendicular to  $BP$ .



- (i) Prove that the equation of the locus of  $P(x,y)$  is  $x(x+2)+y(y-4)=0$  **3**
- (ii) Deduce that the equation in (i) represents a circle and find its centre and radius. **3**

(b)



Consider the region bounded by the curve  $y = \frac{2}{1+x^2}$  and the line  $y = 1$  as shown in the diagram.

The region is revolved about the  $y$  axis. Find the volume of the solid of revolution generated. **4**

**Question 7 continues next page**

- (c) The population,  $P$ , of a rural town is growing exponentially according to the equation  $P = 5000e^{0.1t}$ ,  $t$  measured in years. Currently, i.e.  $t = 0$ , the population is increasing at a rate of 500 people/year.

What rate of increase, correct to the nearest hundred, is expected after 20 more years?

**2****End of Question 7**

- (a) Simplify the expression  $x^{-1}y^2\left(x^{\frac{1}{2}} - y^{-1}\right)\left(x^{\frac{1}{2}} + y^{-1}\right)$ , giving your answer with positive indices. **2**
- (b) A particle moves on a straight line so that its velocity,  $v$  m/s, at any time  $t$  seconds is given by  $v = (t-1)^4 + \frac{t}{2}$ ,  $t \geq 0$
- (i) Find the initial velocity and show that the particle never stops. **2**
- (ii) Find the initial acceleration of the particle. **2**
- (iii) Find the least value of the velocity. **2**
- (iv) Sketch the velocity-time graph. **2**
- (v) Find the distance travelled by the particle in the first 2 seconds. **2**


**End of Question 8**

- (a) In the land of Welf the interest on investments is paid continuously. If an interest rate of  $100k\%$  p.a. is given then it can be shown that the amount in the fund after  $t$  years,  $t \geq 0$ , is  $A(t)$  where

$$A(t) = Pe^{kt}, \quad P \text{ is the initial investment.}$$

- |       |   |          |
|-------|---|----------|
| (i)   | A particular fund, $F$ , in Welf pays 10% p.a. interest. Show that $k = 0.1$  | <b>1</b> |
| (ii)  | Sally invests \$5000 into fund $F$ for 20 years. Find, correct to the nearest dollar, the amount Sally would have after the 20 years.   | <b>1</b> |
| (iii) | Sally wants to be more wealthy in Welf and decides not to terminate her investment of \$5000 until it has grown to at least \$100 000. For how many years will she need to wait?                        | <b>2</b> |
| (iv)  | Lucy also invests in fund $F$ . She decides to deposit \$1500 into the fund at the start of each year for 20 years. Find, correct to the nearest dollar, the amount Lucy would have after the 20 years. | <b>3</b> |


- (b)



In the diagram,  $ABCD$  is a parallelogram. The diagonals  $AC$  and  $BD$  meet at  $P$ .  $M$  is the mid-point of  $AB$  and  $MC$  meets  $BD$  at  $Q$ .

- |       |   |          |
|-------|---|----------|
| (i)   | Show that $\triangle CDQ$ is similar to $\triangle MBQ$ | <b>1</b> |
| (ii)  | Deduce that $DQ = 2BQ$                                  | <b>2</b> |
| (iii) | Prove that $BQ = 2QP$                                   | <b>2</b> |

**End of Question 9**



In the diagram,  $O$  is the centre of a circle of radius  $\sqrt{6}$  cm and  $PQ$  is a chord of length 4 cm.  $ABCD$  is a rectangle constructed in the minor segment cut off by chord  $PQ$ .  $OM$  is drawn perpendicular to  $PQ$  so that  $M$  is the mid-point of both chord  $PQ$  and side  $AB$ .  $N$  is the mid-point of side  $CD$ .

Let  $\angle CON = \theta$ ,  $\theta$  in radians.

- (i) Show that  $OM = \sqrt{2}$  cm **1**
  
- (ii) Show that  $0 < \theta < 1$  **2**
  
- (iii) Show that the area,  $a$ , of rectangle  $ABCD$  is given by  $a = 4\sqrt{3} \sin \theta (\sqrt{3} \cos \theta - 1)$  **3**
  
- (iv) Show that  $\frac{da}{d\theta} = 4\sqrt{3} (2\sqrt{3} \cos^2 \theta - \cos \theta - \sqrt{3})$  **3**
  
- (v) Find the maximum area of the rectangle. **3**

**End of Examination**

TKS MATHEMATICS TRIAL 2004 SOLUTIONS

Qn 1 (a)  $\sin 0.012 = 0.012$ , 2 sig. figs

(b) Domain  $x > 0$ , Range all real values for  $y$

(c)  $\lim_{x \rightarrow 0} \frac{x(x^2-3)}{6x} = \lim_{x \rightarrow 0} \frac{x^2-3}{6} = -\frac{1}{2}$

or  $\lim_{x \rightarrow 0} \left( \frac{x^2}{6} - \frac{3}{6} \right) = -\frac{1}{2}$

(d)  $2x+3=2$  or  $2x+3=-2$

$\therefore x = -\frac{1}{2}$  or  $-\frac{5}{2}$

(e)  $\frac{(2x+3)^5}{5 \times 2} = \frac{(2x+3)^5}{10}$

(f)  $2(1-2x) = 7+6x$

$2-4x = 7+6x$

$\therefore 10x = -5$

$x = -\frac{1}{2}$

Qn 2

$$(a) \frac{dy}{dx} = \frac{(x^2+1) - x(2x)}{(x^2+1)^2} = \frac{1-x^2}{(x^2+1)^2}$$

$$\text{At } (0,0), \frac{dy}{dx} = 1$$

$\therefore$  gradient of normal at  $(0,0)$  is  $-1$

$$(b) (i) \text{ gradient } AB = \frac{0-8}{2--4} = -\frac{8}{6} = -\frac{4}{3}$$

$$(ii) \text{ line } AB \text{ is } y = -\frac{4}{3}(x-2)$$

$$\text{or } 3y = -4x + 8$$

$$\text{ie } 4x + 3y - 8 = 0$$

(iii)  $BC$  = perpendicular distance from  $P$  to line  $AB$

$$\therefore BC = \frac{12 + 21 - 8}{\sqrt{4^2 + 3^2}} = \frac{25}{5} = 5$$

$$(iv) M_{AB} = \left(-\frac{2}{2}, \frac{8}{2}\right) = (-1, 4)$$

$\therefore$  From (iii), if  $PM_{AB} = 5$  then  $P$  is mid-point of  $CD$

$$PM_{AB} = \sqrt{4^2 + 3^2} = 5 \quad \therefore \text{result.}$$

(v) Diagonals meet at mid-point of  $PM_{AB}$ , from (iv)

$$\text{ie at } \left(\frac{2}{2}, \frac{11}{2}\right) = \left(1, \frac{11}{2}\right) = Q, \text{ say}$$

(vi) [LOTS OF WAYS]

$$\text{e.g. } B \rightarrow Q = Q \rightarrow D$$

$$\Rightarrow D = (0, 11)$$

Qn 3

$$(a) (i) \left[-\cos x\right]_0^{\frac{\pi}{6}} = -\frac{\sqrt{3}}{2} - -1 = 1 - \frac{\sqrt{3}}{2}$$

$$(ii) = 2 \int_0^1 \frac{3x^2}{x^3+1} dx = 2 \left[ \ln(x^3+1) \right]_0^1 \\ = 2 \ln 2$$

(b) (i)  $\cup$   $\therefore$  all values of  $k$

$$(ii) \text{ Need } \Delta < 0 \Rightarrow 4 - 4k < 0 \\ \text{or } 4k > 4 \text{ or } k > 1$$

(c) (i)  $\angle ECP = 180^\circ$ ,  $ECP$  a straight line

$$\therefore \angle BCP = 10^\circ$$

(ii)  $\angle BCD = \angle ABC = 50^\circ$ , alternate  $\angle$ s in  $\parallel$  lines  $AB, CD$

$$\therefore \angle ECD = 360^\circ - (170^\circ + 50^\circ), \text{ angles at point } C \\ = 140^\circ$$

(iii) From (ii),  $\angle QPC = \angle ECD = 140^\circ$  and these  
- are corresponding angles


$$\therefore PQ \parallel CD$$

But  $AB \parallel CD$

$$\therefore PQ \parallel AB$$

Qn 4

(a) (i)



(ii) period =  $2\pi$

(ii)  $\therefore \frac{x}{2} = \frac{\pi}{4}$  or  $-\frac{3\pi}{4}$  for  $-\pi \leq \frac{x}{2} \leq \pi$

[Sketch Helps]

$$\Rightarrow x = \frac{\pi}{2} \text{ or } -\frac{3\pi}{2}$$

(b) (i)  $S_{40} = \frac{40}{2} [22 + 39 \times 2] = 20 \times 100 = 2000$

(ii)  $\therefore \frac{n}{2} (22 + (n-1)2) = \frac{n}{2} (-28 + (n-1)3)$

$$\Rightarrow 22 + 2n - 2 = -28 + 3n - 3$$

$\therefore n = 51$  i.e. there's 51 terms

(c) Put  $x=1$ ,  $\therefore 4a = 1+7 \Rightarrow a=2$

$\therefore$  Equating coefficients of  $x^2$ ,  $2+b=1$

$$\Rightarrow b=-1$$

Put  $x=-1$ ,  $\therefore -2c = 1-7$

$$\therefore c=3$$

[LOTS OF WAYS, of course]

Qn 5

(a) (i)  $f'(x) = 4x^3 - 12x + 8 = 4(x^3 - 3x + 2)$

Now,  $(x+2)(x-1)^2 = (x+2)(x^2 - 2x + 1)$   
 $= x^3 - 2x^2 + x + 2x^2 - 4x + 2$   
 $= x^3 - 3x + 2$

$\therefore f'(x) = 4(x+2)(x-1)^2$

(ii) A test for increasing function is  $f'(x) > 0$

$4(x+2)(x-1)^2 > 0 \Rightarrow x+2 > 0$  since  $4(x-1)^2 \geq 0$  for all  $x$

$\therefore f(x)$  is increasing for  $x > -2$

[Note If  $x=1$ ,  $f'(x)=0$  but the function is still increasing.  
A curve is increasing if as  $x$  increases,  $f(x)$  increases]

(iii)  $f'(-2) = 0$   $\therefore$  at  $x=-2$  there's a stationary point

but for  $x < -2$ ,  $f'(x) < 0$   
&  $x > -2$ ,  $f'(x) > 0 \Rightarrow$  

$\therefore$  at  $x=-2$  there's a minimum turning point

(iv)  $f'(x) = 4(x^3 - 3x + 2)$


$\therefore f''(x) = 4(3x^2 - 3) = 12(x^2 - 1)$

$\therefore f''(1) = 0$  and  $f''(0) < 0$ ,  $f''(2) > 0$   
change in concavity  
and, as well,  $f'(1) = 0$

$\therefore$  at  $x=1$  there's a horiz. pt. of inflection


Qn 6

(a) (i)  $PQ = \sqrt{3^2 + 9^2} = \sqrt{90}$

(ii)   $\therefore \cos \theta = \frac{25 + 25 - 90}{2 \times 5 \times 5} = \frac{-40}{50} = -\frac{4}{5}$

(iii) Area =  $\frac{1}{2} \times 5^2 \times \theta$  where  $\cos \theta = -\frac{4}{5}$   
 $= \frac{1}{2} \times 25 \times 2.498 \dots = 31.2 \text{ u}^2, 1 \text{ d.p.}$

(b)  $A = \int_1^3 3x^2 + \frac{3}{x^2} - 6 \, dx$   
 $= \left[ x^3 - \frac{3}{x} - 6x \right]_1^3$   
 $= 27 - 1 - 18 - (1 - 3 - 6) = 16 \text{ u}^2$



(i) focal length  $a = 2$

(ii) Vertex =  $(0, 2)$

$\therefore$  equation is  $(x-0)^2 = 4 \times 2 (y-2)$

i.e.  $x^2 = 8(y-2)$

Qn 7

(a) (i) Since  $AP \perp BP$  the product of their gradients is  $-1$

$$\therefore \frac{y}{x+2} \times \frac{y-4}{x} = -1$$

$$\text{or } y(y-4) = -x(x+2)$$

$$\text{or } x(x+2) + y(y-4) = 0$$

$$(ii) \therefore x^2 + 2x + y^2 - 4y = 0$$

$$\Rightarrow (x+1)^2 - 1 + (y-2)^2 - 4 = 0$$

$$\text{or } (x+1)^2 + (y-2)^2 = 5$$

is a circle, centre  $(-1, 2)$ , radius  $\sqrt{5}$

$$(b) V = \pi \int_1^2 x^2 dy \text{ where } y = \frac{2}{1+x^2}$$

$$\therefore 1+x^2 = \frac{2}{y} \text{ or } x^2 = \frac{2}{y} - 1$$

$$\therefore V = \pi \int_1^2 \left( \frac{2}{y} - 1 \right) dy$$

$$= \pi [2 \ln y - y]_1^2$$

$$= \pi (2 \ln 2 - 2 - (0 - 1)) = \pi (2 \ln 2 - 1) \text{ m}^3$$

$$(c) \frac{dP}{dt} = 5000 \times 0.1 e^{0.1t} = 500 e^{0.1t}$$

$$\therefore \text{when } t = 20, \frac{dP}{dt} = 500 e^2 \approx 3700 \text{ people/yr,}$$

nearest hundred.

Qn 8

$$(a) \quad x^{-1} y^2 (x - y^{-2}) = y^2 - x^{-1} = y^2 - \frac{1}{x}$$
$$\left[ \text{or } \frac{y^2 x - 1}{x} \right]$$

$$(b) \quad (i) \quad t=0, \quad v = (-1)^4 + 0 = 1 \text{ m/s}$$

$$\text{Now, } (t-1)^4 \geq 0 \text{ for all } t \geq 0$$

$$\text{and } \frac{t}{2} > 0 \text{ for all } t > 0$$

$\therefore v > 0$  for all  $t \geq 0$  i.e. particle never stops

$$(ii) \quad \ddot{x} = \frac{dv}{dt} = 4(t-1)^3 + \frac{1}{2}$$

$$\therefore \text{ when } t=0, \quad \ddot{x} = -4 + \frac{1}{2} = -3\frac{1}{2} \text{ m/s}^2$$


(iii) From (i) and (ii), least  $v$  occurs when  $\ddot{x} = 0$

$$\Rightarrow 4(t-1)^3 + \frac{1}{2} = 0$$

$$\text{or } (t-1)^3 = -\frac{1}{8}$$

$$\therefore t-1 = -\frac{1}{2} \text{ or } t = \frac{1}{2}$$

$$\therefore \text{ least } v = \frac{1}{16} + \frac{1}{4} = \frac{5}{16} \text{ m/s}$$



$$(v) \quad x = \int_0^2 (t-1)^4 + \frac{t}{2} dt = \left[ \frac{(t-1)^5}{5} + \frac{t^2}{4} \right]_0^2$$
$$= \frac{1}{5} + 1 - \left( -\frac{1}{5} + 0 \right) = 1\frac{2}{5} \text{ m}$$

Qn 9

(a) (i)  $\therefore 100k = 10 \Rightarrow k = 0.1$

(ii)  $A(20) = 5000 e^{0.1 \times 20} = 5000 e^2$   
 $= \$36945$ , nearest dollar

(iii) Hence,  $A(t) = 5000 e^{0.1t} \geq 100000$

$$\therefore e^{0.1t} \geq 20$$

$$\text{or } 0.1t \geq \ln 20$$

$$\therefore t \geq \frac{\ln 20}{0.1} = 29.957 \dots$$

$\Rightarrow$  Sally needs to wait approx 30 years

(iv) Lucy would have

$$1500 e^{0.1 \times 20} + 1500 \times e^{0.1 \times 19} + \dots + 1500 e^{0.1 \times 1}$$

$$= 1500 e^1 + 1500 e^2 + 1500 e^3 + \dots + 1500 e^2,$$

is a geometric series where  $a = 1500 e^1$   
 $r = e^1$

$$\therefore \text{Lucy would have } \frac{1500 e^1 ((e^1)^{20} - 1)}{e^1 - 1}$$

$$= \frac{1500 e^1 (e^2 - 1)}{e^1 - 1}$$

$$= \$100707$$
, nearest dollar

Q9 (a)

(i) In  $\Delta$ s  $CDQ$ ,  $MBQ$

$\angle CDQ = \angle MBQ$ , alternate  $\angle$ s in  $\parallel$  lines  $CD, BA$

$\angle DCQ = \angle BMQ$ , "

$\therefore \Delta CDQ \parallel \Delta MBQ$ , 2 angles equal

(ii) From (i),  $\frac{DQ}{BQ} = \frac{CD}{BM}$ , ratios of corresponding sides

$= 2$ , since  $M$  is the mid-point of  $AB$ ,  $AB = CD$

$\therefore DQ = 2BQ$


(iii) Now,  $BQ + QP = PD$ , diagonals of  $\parallel$  gram bisect each other

$= DQ - QP$

$= 2BQ - QP$ , (ii)

$\therefore BQ = 2QP$

Qn 10




$$\therefore OM^2 + 4 = 6$$

$$\Rightarrow OM = \sqrt{2} \text{ cm}$$

(ii) as C approaches the vertical position above N then  $\theta \rightarrow 0$ , otherwise  $\theta > 0$


max  $\theta$  occurs as C approaches Q



$$\sin \theta = \frac{2}{\sqrt{6}}$$

$$\Rightarrow \theta = 0.955 \dots < 1$$

$$\therefore 0 < \theta < 1$$



$$\text{Then } \sin \theta = \frac{NC}{\sqrt{6}}, \quad NC = \sqrt{6} \sin \theta$$

$$\therefore DC = 2\sqrt{6} \sin \theta$$

$$\text{and } \cos \theta = \frac{ON}{\sqrt{6}} \quad \therefore ON = \sqrt{6} \cos \theta$$

$$\therefore MN = \sqrt{6} \cos \theta - \sqrt{2} = CB$$

$$\therefore \text{Area, } a = 2\sqrt{6} \sin \theta (\sqrt{6} \cos \theta - \sqrt{2})$$

$$= 2\sqrt{6}\sqrt{2} \sin \theta (\sqrt{3} \cos \theta - 1)$$

$$= 4\sqrt{3} \sin \theta (\sqrt{3} \cos \theta - 1)$$

$$\begin{aligned}
 \text{(iv)} \quad \frac{da}{d\theta} &= 4\sqrt{3} \left( \sin\theta (-\sqrt{3}\sin\theta) + (\sqrt{3}\cos\theta - 1)\cos\theta \right) \\
 &= 4\sqrt{3} \left( -\sqrt{3}\sin^2\theta + \sqrt{3}\cos^2\theta - \cos\theta \right) \\
 &= 4\sqrt{3} \left( -\sqrt{3}(1-\cos^2\theta) + \sqrt{3}\cos^2\theta - \cos\theta \right) \\
 &= 4\sqrt{3} \left( 2\sqrt{3}\cos^2\theta - \cos\theta - \sqrt{3} \right)
 \end{aligned}$$

$$\text{(v)} \quad \frac{da}{d\theta} = 4\sqrt{3} \left( (\sqrt{3}\cos\theta + 1)(2\cos\theta - \sqrt{3}) \right)$$

= 0 only if  $2\cos\theta - \sqrt{3} = 0$  since  $0 < \theta < 1$

and since as  $\theta \rightarrow 0$  or  $\theta \rightarrow 1$ , then  $a \rightarrow 0$ ,  
the value of  $\theta$  from  $2\cos\theta - \sqrt{3} = 0$  must produce  
a maximum turning point  $\Rightarrow$  maximum area

$\therefore$  max area occurs when  $\cos\theta = \frac{\sqrt{3}}{2}$  and  $\therefore \sin\theta = \frac{1}{2}$

$$\begin{aligned}
 \therefore \text{max } a &= 4\sqrt{3} \cdot \frac{1}{2} \left( \sqrt{3} \cdot \frac{\sqrt{3}}{2} - 1 \right) \text{ cm}^2 \\
 &= \sqrt{3} \text{ cm}^2
 \end{aligned}$$