


Name: _____

Teacher: _____



PYMBLE LADIES' COLLEGE
MATHEMATICS
TRIAL HSC EXAMINATION
2006

Reading Time: 5 minutes
Working Time: 3 hours

Instructions to students:

- Write using blue or black biro.
- All questions may be attempted.
- Diagrams are not to scale.
- All necessary working should be shown in every question.
- Your name and your teacher's name may be written before you begin the assessment.
- Start each question in a new booklet.
- Marks may be deducted for careless or untidy work.
- Board-approved calculators may be used.
- A table of standard integrals is provided.


Question 1 (12 Marks) Use a SEPARATE writing booklet

- (a) Evaluate $\frac{4.26 + 3.81}{3\sqrt{6.27}}$. Give the answer correct to three significant figures. Marks 2

- (b) Factorise $x^3 - 8$. 1

- (c) Find the values of a and b such that $\frac{8}{\sqrt{5} - 3} = a - \sqrt{b}$. 3

- (d) The diagram below represents the graph of $y = f(x)$. (2)




State its domain and range. (2)

- (e) Find the values of x for which $|5 - 2x| \geq 1$. (2)

- (f) Differentiate $\frac{\sqrt{x}}{4}$ with respect to x . (2)

Question 2 (12 Marks) Use a SEPARATE writing booklet

In the diagram, the points P and Q have coordinates $(0,1)$ and $(5,6)$ respectively. The line through T and R has equation $y = \frac{5-2x}{2}$.
Copy the diagram onto your answer booklet.



- (i) Find the size of the angle which the line PQ makes with the positive direction of the x -axis.

- (ii) Show that the equation of the line PQ is $x - y + 1 = 0$.

- (iii) Given M is the point where the line PQ intersects the line RT, find the coordinates of M.

- (iv) Find the perpendicular distance from P to the line RT.

- (v) If $\angle PMR$ is a right-angle, then find the area of $\triangle PRM$.

- (vi) On your diagram shade the region which satisfies $x - y + 1 \geq 0$, $2x + 2y \leq 5$ and $y \geq 0$ simultaneously.

Marks

2

1

2

2

3

2

Question 3 (12 Marks) Use a SEPARATE writing booklet

Marks

2

- (a) Differentiate $2x \tan x$ with respect to x .

2

- (b) Given $f(x) = x^3 - 4x^{-1}$, find the value of $f'(\sqrt{2})$.


2

- (c) Evaluate $\int_0^{\frac{\pi}{9}} \cos 3x \, dx$.

2

- (d) Find $\int \frac{3}{1+2x} \, dx$.

- (e) A cone is formed by folding the sector ABO so that the edges OA and OB coincide.



Find:

- (i) the exact area of the sector ABO.

1


- (ii) the exact length of the arc AB.

1

- (iii) the radius of the base of the cone formed.

2

Question 4 (12 Marks) Use a SEPARATE writing booklet

(a)  Marks

PQR is a triangle with $PQ = 48 \text{ mm}$, $QR = 6.5 \text{ cm}$ and $\angle QPR = 20^\circ$.
Find the size of $\angle PQR$ correct to the nearest degree.

3

(b) Consider the function $f(x) = -x^5 + 15x^3$.

(i) Find the coordinates of the stationary points of the curve $y = f(x)$ and determine their nature.

4

(ii) Sketch the curve showing all important features including the x -intercepts and points of inflection.

4

(iii) State the values of x for which the curve $y = f(x)$ is concave down.

1

Question 5 (12 Marks) Use a SEPARATE writing booklet

Marks

(a) Consider the parabola $(y - 4)^2 = 8(x + 2)$.

- (i) Write down the coordinates of the vertex.
- (ii) Find the focus.
- (iii) Find the y -intercepts.

1

2

2

(b) If α and β are the roots of the quadratic equation $3x^2 - 4x - 1 = 0$, find the value of:

- (i) $\alpha + \beta$.
- (ii) $\alpha\beta$.
- (iii) $\alpha^2 + \beta^2$.

1

1

2

(c) Given the sequence $\ln 4, \ln 16, \ln 64, \dots$

- (i) Show that it is arithmetic.


1

- (ii) Hence, find the sum of the first 20 terms in exact form.

2

Question 6 (12 Marks) Use a SEPARATE writing booklet

	Marks
(a) The velocity of a particle v cm/s moving in a straight line is given by $v = 1 + 2t - 3t^2$.	
(i) If the initial displacement is 3cm to the right of 0, calculate the displacement after 2 seconds.	2
(ii) When is the particle at rest?	2
(iii) How far does the particle travel in the third second?	2
(iv) Describe the motion of the particle.	1




PQRS is a parallelogram. M is the midpoint of SR.

PM produced meets QR produced at T.

- | | |
|--|---|
| (i) Prove that $\triangle PMS \cong \triangle TMR$. | 3 |
| (ii) Prove that PRTS is a parallelogram. | 2 |

Question 7 (12 Marks) Use a SEPARATE writing booklet

	Marks
(a) (i) Sketch the graph of $y = 3 \sin 2x$ for $0 \leq x \leq 2\pi$.	2
(ii) What is the period of the curve?	1
(iii) State the amplitude.	1
(iv) Find the area between the curve and the x-axis if $\frac{\pi}{4} \leq x \leq \frac{\pi}{2}$.	3
(b) (i) Copy and complete the table of values for $y = 4^x$.	1




- | | |
|---|---|
| (ii) Hence, using these three values and the trapezoidal rule, find an approximation for $\int_{-1}^1 4^x dx$. | 2 |
| (iii) (a) Find the derivative of 4^x with respect to x . | 1 |

(b) Hence, or otherwise, find the exact value of $\int_{-1}^1 4^x dx$.

Question 8 (12 Marks) Use a SEPARATE writing booklet

- | | Marks |
|--|-------|
| (a) Find the coordinates of the point on the curve $y = e^{3x}$ where the tangent is perpendicular to the line $y = 4 - \frac{x}{6}$. | 3 |
| (b) (i) Show that $\sin^2 x \cos x = \cos x - \cos^3 x$. | 1 |
| (ii) Hence, find $\frac{d}{dx} \left(\sin x - \frac{1}{3} \sin^3 x \right)$. | 3 |



The diagram above is of the exponential curve $y = e^{\frac{x}{2}}$ and the parabola $y = \frac{e}{4}x^2$. The point A is the first point where the two graphs meet on the right hand side of the y-axis.

- | | |
|---|---|
| (i) Show that A is the point $(2, e)$. | 2 |
| (ii) Show that the shaded area is $\frac{4e}{3} - 2$ units ² . | 3 |

Question 9 (12 Marks) Use a SEPARATE writing booklet

- | | Marks |
|---|-------|
| (a) The rate of increase of a population P(t) of people in a certain city is governed by the equation $\frac{dP}{dt} = kP$ where k is a constant and t is the time in years. The population of the city doubles every twenty years. | |
| (i) Show that $k = \frac{1}{20} \ln 2$. | 2 |
| (ii) In which year will the city reach a population three times that which it had at the beginning of 2006? | 2 |
| (iii) If at the beginning of 2010 the population is 20 million, what will be the population at the beginning of the year 2060? | 2 |

- (b) Sarah wishes to buy a car. She has worked out that she can afford repayments of \$400 a month for 5 years.

The interest rate on offer is 24% pa (reducible) calculated monthly.

Let A_n be the amount owing after n months based on a monthly repayment of \$400 and P being the amount borrowed.

- | | |
|--|---|
| (i) Give an expression for A_2 . | 1 |
| (ii) Show that $A_n = 1.02^n P - 20000(1.02^n - 1)$. | 3 |
| (iii) Hence, determine how much money Sarah is able to borrow? | 2 |

Question 10 (12 Marks) Use a SEPARATE writing booklet

Marks

- (a) Differentiate $\ln(x + \sqrt{x^2 + a^2})$ and hence find $\int \frac{1}{2\sqrt{x^2 + a^2}} dx$. 3

- (b) The region bounded by the curve $y = \ln x$, the axes and the line $y = \ln 2$, is rotated about the y -axis. 3

Find the volume of the solid formed.

- (c) ABCDE is a pentagon of fixed perimeter P cm. Its shape is such that ABE is an equilateral triangle and BCDE is a rectangle.

If the length AB is x cm:

- (i) Show that the length BC is $\frac{P-3x}{2}$ cm. 1

- (ii) Show that the area of the pentagon is given by
$$A = \frac{1}{4} [2Px - (6 - \sqrt{3})x^2]$$
. 2

- (iii) Find the value of $\frac{P}{x}$ for which the area of the pentagon is a maximum. 3

End of paper

2006 HSC TRIALS - MATHEMATICS SOLUTIONS AND

Q1

$$\text{a) } 4.26 + 3.81 \\ = 3\sqrt{6.27}$$

$$= 1.07428 \dots \\ = 1.07 \text{ (3 s.f.)}$$

$$\text{b) } x^3 - 8 \\ = (x-2)(x^2 + 2x + 4)$$

$$\text{c) } 8$$

$$\frac{8}{\sqrt{5} - 3} \times \frac{\sqrt{5} + 3}{\sqrt{5} + 3} = \frac{1}{2} \\ = \frac{8(\sqrt{5} + 3)}{-4} = \frac{-2(\sqrt{5} + 3)}{2}$$

$$= -6 - 2\sqrt{5} = -6 - \sqrt{20} \frac{1}{2}$$

$$\therefore a = -6 \frac{1}{2}, b = 20 \frac{1}{2}$$

d) Domain: all real x ; $x \neq 4$

Range: $y \geq -2$. L.R.W. (2)

$$\text{e) } |5-2x| \geq 1$$

$$5-2x \geq 1 \text{ OR } -5+2x \geq 1$$

$$-2x \geq -4$$

$$x \leq 2$$

$$2x \geq 6$$

$$x \geq 3$$

$$\text{f) } \frac{d}{dx} \left(\frac{\sqrt{x}}{4} \right)$$

$$= \frac{1}{4} \frac{d}{dx} \left(\frac{1}{4} x^{\frac{1}{2}} \right)$$


$$= \frac{1}{4} \left(\frac{1}{2} x^{-\frac{1}{2}} \right)$$

$$= \frac{1}{8} x^{-\frac{1}{2}}$$

$$= \frac{1}{8\sqrt{x}}$$

MARKING GUIDELINES

Q2



$$\text{iis } m_{PQ} = \frac{6-1}{5-0} = 1$$

Angle line PQ with x-axis = 45°

ii) Eq. of line PQ

$$\Rightarrow 1 = \frac{y-1}{x-0}$$

$$x = y - 1$$

$$x - y + 1 = 0$$

$$\text{iii) } \begin{cases} x - y + 1 = 0 \\ y = \frac{5-x}{2} \end{cases}$$

$$x - \left(\frac{5-x}{2} \right) + 1 = 0$$

$$2x = \frac{3}{2}$$

$$x = \frac{3}{4}$$

$$y = 1 \frac{3}{4}$$

$$M = \left(\frac{3}{4}, 1 \frac{3}{4} \right)$$

$$\text{iv) Eq. of RT} \Rightarrow 2y = 5-2x$$

$$2x + 2y - 5 = 0$$

Perpendicular distance from P to R.T.

$$= \frac{|2(0) + 2(1) - 5|}{\sqrt{2^2 + 2^2}}$$

$$= \frac{3}{2\sqrt{2}}$$

$$= \frac{3\sqrt{2}}{4}$$

$\frac{1}{2}$ for labelling RT
 $\frac{1}{2}$ for not colouring below
1 for correct region

$$v) 2x + 2y - 5 = 0$$

$$y = 0 \Rightarrow 2x = 5$$

$$x = \frac{5}{2} = \frac{1}{2}$$

$$R\left(\frac{5}{2}, 0\right)$$

$$\begin{aligned}\text{Length of RM} &= \sqrt{\left(\frac{5}{2} - \frac{3}{4}\right)^2 + (0 - 1\frac{3}{4})^2} \\ &= \sqrt{\frac{3}{4} \times 2} \\ &= \frac{7\sqrt{2}}{4} = \frac{1}{2}\end{aligned}$$

i) Area of $\triangle PRM$

$$\begin{aligned}&= \frac{1}{2} \times \frac{7\sqrt{2}}{4} \times \frac{3}{2\sqrt{2}} = \frac{1}{2} \\ &= \frac{1}{16} \text{ sq. units}\end{aligned}$$

Q3

$$a) \frac{d}{dx}(2x \tan x)$$

$$= 2 \tan x + 2x \sec^2 x$$

$$\text{Let } u = 2x \quad v = \tan x \\ \frac{du}{dx} = 2 \quad \frac{dv}{dx} = \sec^2 x \quad \frac{1}{2}$$

(2)

$$b) f(x) = x^3 - 4x^{-1}$$

$$f'(x) = 3x^2 + 4x^{-2}$$

$$f'(\sqrt{2}) = 3(\sqrt{2})^2 + 4(\sqrt{2})^{-2} =$$

$$= 6 + 2$$

$$= 8 = \frac{1}{2}$$

(2)

$$c) \int_0^{\frac{\pi}{4}} \cos 3x \, dx$$

$$= \frac{1}{3} \sin 3x \Big|_0^{\frac{\pi}{4}}$$

$$= \frac{1}{3} \sin \frac{\pi}{3} - \frac{1}{3} \sin 0 = \frac{1}{3}$$

$$= \frac{\sqrt{3}}{6}$$

(2)

$$d) \int_{\frac{\pi}{2}}^{\pi} \frac{3}{1+2x} \, dx$$

$$= \frac{3}{2} \ln(1+2x) + C \Big|_{\frac{\pi}{2}}$$

(2)

e) is Area of sector ABO

$$\begin{aligned}&= \frac{1}{2} \times 8^2 \times \frac{2\pi}{3} \\ &= \frac{64\pi}{3} \text{ cm}^2\end{aligned}$$

(1)

$$\text{iii) Arc AB} = \frac{8 \times 2\pi}{3} = \frac{16\pi}{3} \text{ cm}$$

(1)

iv) Circumference of base = length of arc AB.

$$2\pi r = \frac{16\pi}{3}$$

$$r = \frac{8}{3} = \frac{1}{2}$$

v) Radius of base of cone

$$= 2\frac{2}{3} \text{ cm.}$$

(2)

Q4


$$\text{a) } \frac{\sin \theta}{4.8} = \frac{\sin 20^\circ}{6.5}$$

$$\sin \theta = \frac{\sin 20^\circ}{6.5} \times 4.8$$

$$\sin \theta = \frac{\sin 20^\circ}{6.5} \times 4.8 \quad \frac{P}{2}$$

$$\theta = 14.629 \dots \frac{1}{2}$$

$$\angle PQR = 180^\circ - 20^\circ - 14.629 \dots \\ = 145^\circ 22' \\ = 145^\circ \frac{1}{2}$$



$$\text{b) } f(x) = -x^5 + 15x^3$$

$$\text{i) } f'(x) = -5x^4 + 45x^2 = 0 \quad \frac{1}{2} \\ = -5x^2(x^2 - 9) = 0$$

$$\begin{cases} x = 0 \\ y = 0 \end{cases} \quad \begin{cases} x = 3 \\ y = 162 \end{cases} \quad \begin{cases} x = -3 \\ y = -162 \end{cases}$$

$$f''(x) = -20x^3 + 90x$$

$$f''(0) = 0 \quad \frac{1}{2} \quad \frac{x+1}{y+1} \quad \frac{0+1}{0+1} \quad \frac{1}{2}$$

$$f''(3) = -20 \times 3^3 + 90 \times 3 < 0 \quad \frac{1}{2}$$

$$f''(-3) = -20(-3)^3 + 90(-3) > 0 \quad \frac{1}{2}$$

i) $(0, 0)$ is a horizontal point of inflection


$(3, 162)$ is a max. turning point $\frac{1}{2}$ per mistake

and $(-3, -162)$ is a min. turning point. ④

$$\text{ii) } f(x) = -x^5 + 15x^3 = 0$$

$$x^3(-x^2 + 15) = 0$$

$$x = 0 \quad \text{OR} \quad x = \pm \sqrt{15} \quad \frac{1}{2}$$



$$f''(x) = -20x^3 + 90x = 0$$

$$-10x(2x^2 - 9) = 0$$

$$x = 0, x = \pm \frac{3}{\sqrt{2}}$$

$$\text{When } x = \frac{3}{\sqrt{2}}, y = \frac{567}{450}$$

$$\text{and when } x = -\frac{3}{\sqrt{2}}, y = -\frac{567}{450}$$

$$x \mid \frac{3}{\sqrt{2}} \quad \frac{3}{\sqrt{2}} \quad \frac{3}{\sqrt{2}} \quad x \mid \frac{3}{\sqrt{2}} \quad \frac{3}{\sqrt{2}} \quad \frac{3}{\sqrt{2}} \quad \frac{3}{\sqrt{2}}$$

$$f''(x) \mid + \quad 0 \quad - \quad f''(x) \mid + \quad 0 \quad -$$

i) Points of inflection at $(\frac{3}{\sqrt{2}}, \frac{567}{450})$ and $(-\frac{3}{\sqrt{2}}, -\frac{567}{450})$ as well. ④

iii) Concave down when $\frac{-3}{\sqrt{2}} < x < 0$ and $x > \frac{3}{\sqrt{2}}$. ①


QUESTION 5

a. $(y-4)^2 = 8(x+2)$

i. Vertex is $(-2, 4)$

1 R/W

① MARK



$$4a = 8$$

$$a = 2$$

Focus $(0, 4)$

② MARKS

iii. y -intercept: sub $x=0$

$$(y-4)^2 = 8(0+2)$$

1

② MARKS

$$(y-4)^2 = 16$$

$$y-4 = \pm 4$$

$$y = 0, 8$$

$$\frac{1}{2} + \frac{1}{2}$$

b. $3x^2 - 4x - 1 = 0$

$$a = 3, b = -4, c = -1$$

① MARK

$$\alpha + \beta = -\frac{b}{a}$$

$$= \frac{4}{3}$$

1 R/W

① MARK

$$\alpha\beta = \frac{c}{a}$$

$$= -\frac{1}{3}$$

1 R/W

iii. $\alpha^2 + \beta^2$

② MARKS

$$= (\alpha + \beta)^2 - 2\alpha\beta$$

$$= \left(\frac{4}{3}\right)^2 - 2 \times -\frac{1}{3}$$

$$= \frac{22}{9}$$

$$= 2^{\frac{4}{9}}$$

c. i. $\ln 4, \ln 16, \ln 64, \dots$

① MARK

To prove $T_2 - T_1 = T_3 - T_2$

$$LHS = T_2 - T_1$$

$$RHS = T_3 - T_2$$

$$= \ln 16 - \ln 4$$

$$= \ln 64 - \ln 16$$

$$= \ln\left(\frac{16}{4}\right)$$

$$= \ln\left(\frac{64}{16}\right)$$

$$= \ln 4$$

$$= \ln 4$$

$$= LHS$$

As $T_2 - T_1 = T_3 - T_2$

then the sequence $\ln 4, \ln 16, \ln 64, \dots$
is arithmetic.

ii. $a = \ln 4, d = \ln 4, n = 20$

② MARKS

$$S_n = \frac{n}{2} [2a + (n-1)d]$$

$$S_{20} = \frac{20}{2} [2\ln 4 + 19\ln 4]$$

$$= 10 \times 21 \ln 4$$

$$= 210 \ln 4$$

$$= 210 \ln 2^2$$

$$= 420 \ln 2$$

QUESTION 6

a. $v = 1 + 2t - 3t^2$

i. (2) $\frac{dx}{dt} = 1 + 2t - 3t^2$
MARKS

$$x = t + t^2 - t^3 + C$$

Sub $t=0, x=3$

$\therefore C = 3$

$$x = t + t^2 - t^3 + 3$$

Sub $t=2$

$$x = 2 + 2^2 - 2^3 + 3$$

$x = 1$

\therefore Particle is 1cm to the right of O after 2 seconds

ii. Particle is at rest when $v=0$

(2) MARKS $1 + 2t - 3t^2 = 0$

$$3t^2 - 2t - 1 = 0$$

$$(3t+1)(t-1) = 0$$

$$t = -\frac{1}{3}, 1$$

rejected as $t \geq 0$

Particle is at rest after 1 second.

iii. $x = t + t^2 - t^3 + 3$

Sub $t=3$

$$x = 3 + 3^2 - 3^3 + 3$$

$$= -12$$

Particle has travelled $= 1 + 12$

from part (i)

$$= 13 \text{ cm}$$

iv. The particle starts 3cm to the right of O.

(1) It moves to the right and stops after 1 second
MARK when it is now 4cm to the right of O.

It then moves to the left through O and carries on in that direction.

b. i. In $\triangle PMS$ and $\triangle TMR$

(3) MARKS $\angle PMS = \angle TMR$ vertically opposite angles
 $SM = RM$ given M is the midpoint of SR
 $\angle PSM = \angle MRT$ alternate angles, $PS \parallel QT$
given PQRS is a parallelogram
 $\therefore \triangle PMS \cong \triangle TMR$ (ASA)
R $\frac{1}{2}$

ii. $PS = RT$ corresponding sides of congruent Δ 's

(2) MARKS $PS \parallel RT$ given PQRS is a parallelogram
then $PS \parallel QR$ and QR is produced to T.


Hence PRTS is a parallelogram as one pair of opposite sides are equal and parallel.

ALTERNATIVE METHOD

PM = TM corresponding sides of congruent Δ 's
SM = RM from part (i)

Hence PRTS is a parallelogram as both diagonals PT and SR are bisected.

QUESTION 7



MARKS

$\frac{1}{2}$ mark for correct shape (no arrow-heads)

$\frac{1}{2}$ mark for correct range

$\frac{1}{2}$ mark for correct x-intercepts

$\frac{1}{2}$ mark for MAX/MIN points labelled (x-values)

ii. PERIOD = $\frac{2\pi}{2}$

MARK

= π R/W

iii. AMPLITUDE = 3 UNITS R/W ignore units

MARK

$\frac{1}{2}$

Area = $\int_{\frac{\pi}{4}}^{\frac{\pi}{2}} 3 \sin 2x \, dx$

MARKS

$\frac{\pi}{4}$

= $3 \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \sin 2x \, dx$

= $-\frac{3}{2} [\cos 2x]_{\frac{\pi}{4}}^{\frac{\pi}{2}}$

= $-\frac{3}{2} \{ \cos \pi - \cos \frac{\pi}{2} \}$

= $-\frac{3}{2} \{ -1 - 0 \}$

= $\frac{3}{2}$ units² or $1\frac{1}{2}$ units²

$\leftarrow \frac{1}{2}$

Alternative Method: Area = $3 \int_0^{\frac{\pi}{4}} \sin 2x \, dx$

0

$y = 4^x$

b. i.

	x	-1	0	1
①	y	4^{-1}	1	4

MARKS

ii.

MARKS

$$\int_{-1}^1 4^x \, dx \div \frac{1}{2} \left(\frac{1}{4} + 4 + 2 \right) = 3\frac{1}{8}$$

$$\int_{-1}^1 4^x \, dx \div \frac{1}{2} \left(\left(\frac{1}{4}+1\right) + (4+1) \right) = 3\frac{1}{8}$$

iii. $\alpha \frac{d}{dx} 4^x$

MARKS

MARKS

① β. $\int_{-1}^x 4^u \, du$

MARKS

$$= \left[\frac{4^u}{\ln 4} \right]_{-1}^x \text{ from part a. } 4 = \frac{1}{\ln 4} \frac{d}{dx} 4^x$$

$$= \frac{1}{\ln 4} [4^x]_{-1}^x$$

$$= \frac{1}{\ln 4} (4 - \frac{1}{4})$$

$$= \frac{15}{4 \ln 4}$$

QUESTION 8

a.

$$y = e^{3x} \quad (1)$$

$$\frac{dy}{dx} = 3e^{3x}$$

MARLS

$$M = -\frac{1}{6}$$

$$\frac{1}{2} + \frac{1}{2}$$

\therefore Gradient of Perpendicular = 6

Want $3e^{3x} = 6$

$$e^{3x} = 2$$

$$3x = \ln 2$$

$$x = \frac{1}{3} \ln 2 \text{ or } \frac{\ln 2}{3}$$

Sub into (1)

$$y = e^{\frac{1}{3} \ln 2}$$

$$= e^{\ln 2}$$

$$= 2$$

\therefore Coordinates of required point is $(\frac{1}{3} \ln 2, 2)$

b. i. To prove $\sin^2 x \cos x = \cos x - \cos^3 x$

(1)

MARLS

$$\text{LHS} = \sin^2 x \cos x$$

$$= (1 - \cos^2 x) \cos x$$

$$= \cos x - \cos^3 x$$

$$= RHS$$

$$\therefore \sin^2 x \cos x = \cos x - \cos^3 x$$

ii.

$$\frac{d}{dx} (\sin x - \frac{1}{3} \sin^3 x)$$

(3)

MARLS

$$= \cos x - \frac{1}{3} \cdot 3 (\sin x)^2 \cdot \cos x$$

$$= \cos x - \sin^2 x \cdot \cos x$$

$$= \cos x - (\cos x - \cos^3 x) \text{ from part (i)}$$

$$= \cos^3 x$$

c. i.

$$y = e^{x/2}$$

$$\text{Sub } (2, e)$$

MARLS

$$e = e^{x/2}$$

$$e = e^1$$

$$e = e^{\frac{1}{4}}$$

True

$\therefore (2, e)$ lies on the

$$\text{line } y = e^{x/2}$$

$e = e^1$ \therefore

True

$\therefore (2, e)$ lies on the

$$\text{line } y = \frac{e}{4} x^2$$

As $(2, e)$ lies on both lines and $x=2 > 0$ (1st Quadrant)
then this must be the coordinates of the point A.

ii.

MARLS

$$\text{Shaded Area} = \int_0^{x/2} (e^{x/2} - \frac{e}{4} x^2) dx$$

$$= \left[2e^{x/2} - \frac{e}{12} x^3 \right]_0^{x/2}$$

$$= \left(2e^{\frac{x}{2}} - \frac{e}{12} \cdot \frac{x^3}{2} \right) = (2e^0 - 0) - \frac{1}{2}$$

$$= 2e - \frac{8e}{12} = 2$$

$$= 2e - \frac{2e}{3} - 2$$

$$= e(2 - \frac{2}{3}) - 2$$

$$= \frac{4e}{3} - 2 \text{ units}^2$$

Q.E.D.

QUESTION 9

a. i. $\frac{dp}{dt} = kp$

MARKS
1
 $\Rightarrow p = p_0 e^{kt}$

Sub $p = 2p_0, t = 20$

$2p_0 = p_0 e^{20k}$

$e^{20k} = 2$

$20k = \ln 2$

$k = \frac{1}{20} \ln 2$

ii. $P = P_0 e^{(\frac{1}{20} \ln 2)t}$

Sub $P = 3P_0$

$3P_0 = P_0 e^{(\frac{1}{20} \ln 2)t}$

$e^{(\frac{1}{20} \ln 2)t} = 3$

$(\frac{1}{20} \ln 2)t = \ln 3$

$t = 20 \ln 3$

$\ln 2$

≈ 31.7 years

During $(2006 + 31)$ 2037 the population will
beginning during

be 3 times that which it had at the beginning of 2006.

iii. $P = P_0 e^{(\frac{1}{20} \ln 2)t}$

MARKS
2
 $\text{Sub } P_0 = 20, t = 50$

$P = 20 e^{(\frac{1}{20} \ln 2)^{50}}$
 $= 20 e^{\frac{5}{2} \ln 2}$

$= 113.137085$ million

or 113.137.085

REFER TO NEXT PAGE FOR ALTERNATIVE SOLUTION

b. i. 24% per annum

MARKS
1
 $A_1 = P \times 1.02 - 400$

$A_2 = A_1 \times 1.02 - 400$

$= (P \times 1.02 - 400) \times 1.02 - 400$

$\therefore A_2 = P \times 1.02^2 - 400(1+1.02)$

ii. $A_3 = A_2 \times 1.02 - 400$

$= P \times 1.02^3 - 400(1+1.02+1.02^2)$

MARKS
3
Similarly

$A_n = P \times 1.02^n - 400(1+1.02+1.02^2+\dots+1.02^{n-1})$

This is a Geometric Series

$a=1, r=1.02, n=n$

$S_n = a \frac{(r^n - 1)}{r - 1}$

$= \frac{1(1.02^n - 1)}{1.02 - 1}$

$\therefore A_n = P \times 1.02^n - 400 \frac{(1.02^n - 1)}{0.02}$

$= P \times 1.02^n - 20000(1.02^n - 1)$

Q.E.D.

$$\text{iii. } n = 5 \times 12$$

$$= 60$$

MARKS

$$A_{60} = P \times 1.02^{60} - 20000 \left(1.02^{60} - 1 \right)^{\frac{1}{2}}$$

$$\text{But } A_{60} = 0 \quad [\text{LOAN IS REPAYED}]$$

$$\therefore P \times 1.02^{60} - 20000 \left(1.02^{60} - 1 \right) = 0$$

$$P = \frac{20000 \left(1.02^{60} - 1 \right)}{1.02^{60}}^{\frac{1}{2}}$$

$$= \$13904.35^{\frac{1}{2}}$$

Sarah is able to borrow \$13904 to the nearest dollar.

Part (a) (iii) ALTERNATIVE SOLUTION

In 2006, population P_0

2010, population 2 Million, $\therefore t=4$

$$20 = P_0 e^{4k}$$

$$P_0 = \frac{20}{e^{(\frac{1}{20} \ln 4) \times 4}} = 17.44101127 \text{ Million} \quad (1)$$

From 2006 to 2060 — 54 years. $(\frac{1}{2})$

$$\therefore P = P_0 e^{(\frac{1}{20} \ln 4) \times 54}$$

$$= 113137085 \quad (\frac{1}{2})$$

QUESTION 10

$$\text{a. } \frac{d}{dx} \ln(x + \sqrt{x^2 + a^2})$$

$$\text{MARKS} = \frac{d}{dx} \ln[x + (x^2 + a^2)^{\frac{1}{2}}]$$

$$= \frac{1}{x + \sqrt{x^2 + a^2}} \times \left[1 + \frac{1}{2} (x^2 + a^2)^{-\frac{1}{2}} \cdot 2x \right]^{\frac{1}{2} + \frac{1}{2}}$$

$$= \frac{1}{x + \sqrt{x^2 + a^2}} \times \left[1 + \frac{x}{\sqrt{x^2 + a^2}} \right]$$


$$= \frac{1}{x + \sqrt{x^2 + a^2}} \times \frac{\sqrt{x^2 + a^2} + x}{\sqrt{x^2 + a^2}}$$

$$= \frac{1}{\sqrt{x^2 + a^2}} \quad \frac{1}{2}$$

$$\int \frac{1}{2\sqrt{x^2 + a^2}} dx = \frac{1}{2} \int \frac{1}{\sqrt{x^2 + a^2}} dx$$

$$= \frac{1}{2} \int \left[\frac{d}{dx} \ln(x + \sqrt{x^2 + a^2}) \right] dx$$

$$= \frac{1}{2} \ln(x + \sqrt{x^2 + a^2}) + C \quad \frac{1}{2} + \frac{1}{2} \text{ for } +C$$



b.

MARKS

$$\sqrt{b} = \pi \int_a^b x^2 dy$$

$$= \pi \int_{\ln 2}^{2y} e^{-y} dy$$

$$\therefore V = \frac{\pi}{2} [e^{2y}]_0^{\frac{1}{2}}$$

$$= \frac{\pi}{2} \{ e^{2 \cdot \frac{1}{2}} - e^0 \}$$

$$= \frac{\pi}{2} \{ e^4 - 1 \}$$

$$= \frac{\pi}{2} (4 - 1)$$


$$= \frac{3\pi}{2} \text{ units}^3$$

[N.B. without π then
2 marks in total]

c.i.

①

MARK



$ED = BC$ opposite sides of a rectangle $BCDE$

Perimeter = $AB + BC + CD + DE + EA$

$AB = AE = x = EB$ Equal sides of Equilateral triangle AEB

$EB = DC = x$ Opposite sides of rectangle $BCDE$

$$P = x + BC + x + BC + x$$

$$P = 3x + 2BC$$

$$2BC = P - 3x$$

$$BC = \frac{P - 3x}{2}$$

ii. Area = Area of $\triangle ABE$ + Area of Rectangle $BCDE$

②

MARKS

$$= \frac{1}{2} ab \sin C + LB$$

$$= \frac{1}{2} \cdot x \cdot x \cdot \sin 60^\circ + x \cdot \frac{P-3x}{2}$$

$$= \frac{1}{2} x^2 \cdot \frac{\sqrt{3}}{2} + \frac{Px - 3x^2}{2}$$

$$= \frac{1}{4} [\sqrt{3}x^2 + 2(Px - 3x^2)]$$

$$= \frac{1}{4} [\sqrt{3}x^2 + 2Px - 6x^2]$$

$$= \frac{1}{4} [2Px - x^2(6 - \sqrt{3})]$$

$$\therefore A = \frac{1}{4} [2Px - (6 - \sqrt{3})x^2] \quad \text{Q.E.D.}$$

iii. Maximum Area occurs when $\frac{dA}{dx} = 0$

③

MARKS

$$\frac{dA}{dx} = \frac{1}{4} [2P - 2(6 - \sqrt{3})x]$$

$$= \frac{1}{2} [P - (6 - \sqrt{3})x]$$

$$\text{Hence want } \frac{1}{2} [P - (6 - \sqrt{3})x] = 0$$

$$P - (6 - \sqrt{3})x = 0$$

$$P = (6 - \sqrt{3})x$$

$$\frac{P}{x} = \frac{6 - \sqrt{3}}{2}$$

$$\frac{d^2A}{dx^2} = -\frac{1}{2}(6 - \sqrt{3})$$

$\frac{d^2A}{dx^2} < 0$ for all values of x \therefore Max Area occurs when $\frac{P}{x} = 6 - \sqrt{3}$.