



**SYDNEY BOYS HIGH SCHOOL**  
MOORE PARK, SURRY HILLS

**2004**  
TRIAL HIGHER SCHOOL  
CERTIFICATE EXAMINATION

# Mathematics

## *General Instructions*

- Reading time – 5 minutes.
- Working time – 3 hours.
- Write using black or blue pen.
- Board approved calculators may be used.
- All *necessary* working should be shown in every question if full marks are to be awarded.
- Marks may **NOT** be awarded for messy or badly arranged work.
- Hand in your answer booklets in **5** sections.  
Section A (Questions 1 & 2),  
Section B (Questions 3 & 4),  
Section C (Questions 5 & 6),  
Section D (Questions 7 & 8) and  
Section E (Questions 9 & 10).
- Start each **NEW** section in a separate answer booklet.

## **Total Marks - 120 Marks**

- Attempt Sections A - E
- All questions are of equal value.

Examiner: *P. Parker*

This is an assessment task only and does not necessarily reflect the content or format of the Higher School Certificate.

**Total marks – 120**  
**Attempt Questions 1 – 10**  
**All questions are of equal value**

Answer each section in a SEPARATE writing booklet. Extra writing booklets are available.

**SECTION A (Use a SEPARATE writing booklet)**

---

Question 1 (12 marks)	Marks
(a) Evaluate $\log_e \left( \tan \frac{5\pi}{12} \right)$ leaving your answer correct to 3 significant figures	2
(b) Differentiate $\sqrt{5x}$	2
(c) Solve $2t^2 - t - 15 = 0$	2
(d) Find a primitive of $3 - 2x$	2
(e) Solve the pair of simultaneous equations	2
$y = 2x$ $3x + 2y = 14$	
(f) Solve $3 - 4x < 1$ and graph the solution on a number line	2

Question 2 (12 marks)

Marks

(a) Differentiate

(i)  $(1 + \cos 2x)^3$  2


(ii)  $x^2 e^{x+2}$  2

(b) Find:

(i)  $\int \frac{\cos x}{\sin x} dx$  1

(ii)  $\int_{\frac{1}{2}}^2 \left(1 - \frac{1}{x^2}\right) dx$  2

(c)



In the diagram above  $\triangle ABC$  is isosceles, with  $AB = AC$ .  
 $M$  is the midpoint of the line  $NY$  and  $XM \parallel AB$ .

(i) By using similar triangles, or otherwise, show that  $\frac{MX}{NB} = \frac{1}{2}$  2

(ii) Hence show that  $\frac{MC}{NB} = \frac{1}{2}$  1

(d) The graph of  $y = g(x)$  passes through the point  $(2, 4)$  and  $g'(x) = 4 - 3x^2$ . 2

Find  $g(x)$ .


**SECTION B (Use a SEPARATE writing booklet)**

Question 3 (12 marks)

Marks

In the diagram below  $A$ ,  $B$  and  $D$  have coordinates  $(0, -3)$ ,  $(3, 1)$  and  $(-5, 5)$  respectively.

The angle  $\theta$  is the angle the line  $AD$  makes with the positive direction of the  $x$  axis.




- |       |  |   |
|-------|--|---|
| (i)   | Find the gradient of the line $AD$ .<br>Hence find $\theta$ to the nearest degree.   | 2 |
| (ii)  | Find the coordinates of $M$ , the midpoint of $BD$ .   | 1 |
| (iii) | Find the coordinates of $C$ , so that $ABCD$ is a parallelogram.   | 1 |
| (iv)  | Show that the line $AB$ has equation $4x - 3y - 9 = 0$ .   | 2 |
| (v)   | Find the perpendicular distance between $D$ and $AB$ .   | 1 |
| (vi)  | Find the area of parallelogram $ABCD$ .  | 2 |
| (vii) | The line $BD$ has equation $x + 2y - 5 = 0$ and meets the $y$ axis at $P$ .<br>Write down the three inequalities that define the region inside $\triangle ABP$ . | 3 |

Question 4 (12 marks)

Marks

(a) Solve  $\cos 2x^\circ = -\frac{1}{2}$  for  $0^\circ \leq x^\circ \leq 360^\circ$

2




(i) Find the coordinates of  $P$ .

2

(ii) Find the area of the shaded region bounded by  $y = \sqrt{x-5}$  and  $y = \frac{1}{5}(x-5)$ .

3

(c)



2


$ABC$  is a right-angled triangle in which  $\angle ACB = 90^\circ$ .

$\triangle CDB$  is isosceles, in which  $CB = DB$ .

$\angle AEC = 68^\circ$  and  $\angle EBD = 90^\circ$ .

Find  $\angle DCB$ , giving reasons.

(d)



The diagram shows a quadrilateral  $ABCD$  with  $\angle ABC = 60^\circ$ .

$AB = 2$  m,  $BC = 4$  m and  $DC = 1$  m and  $AB \parallel DC$ .

(i) Using the cosine rule, calculate  $AC$ .

1

(ii) Hence find  $AD$ , correct to 3 significant figures

3

**SECTION C (Use a SEPARATE writing booklet)**

Question 5 (12 marks)


**Marks**

- (a) A curve  $\mathcal{C}$  has equation  $y = x^3 - 5x^2 + 7x - 14$ .
- (i) Show  $\frac{dy}{dx} = (3x - 7)(x - 1)$  **1**
- (ii) Find the coordinates of the stationary points and determine their nature. **3**
- (iii) Sketch the graph of  $\mathcal{C}$ , given that an  $x$  intercept occurs in the interval  $4 \leq x \leq 5$ . **2**
- (iv) Find the values of  $x$  for which  $\mathcal{C}$  is concave down. **1**
- 
- (b) A polygon has 40 sides.
- The lengths of the sides, starting with the smallest, form an arithmetic series.
- The perimeter of the polygon is 495 cm and the length of the longest side is twice that of the shortest side.
- For this series:
- (i) Find the first term. **3**
- (ii) The common difference. **2**

Question 6 (12 marks)

Marks

- (a) The diagram below shows a sector of a circle of radius 10 cm. 2  
 Find the value of  $\theta$  to the nearest degree.



- (b) Consider the series  $\cos^2 x + \cos^4 x + \cos^6 x + \dots$  for  $0 < x < \frac{\pi}{2}$
- (i) Explain why a limiting sum exists. 1
- (ii) Find the limiting sum, expressing the answer in simplest form. 2
- (c) The rate at which people,  $N$ , are admitted to Homebake, a music festival in the Domain, is given by
- $$\frac{dN}{dt} = 450t(8-t)$$
- where  $t$  is measured in hours.
- (i) Find the maximum rate of people being admitted to the festival. 1
- (ii) If initially  $N = 0$ , find an expression for the amount of people present at time  $t$ . 2
- (iii) The festival *lasted* as long as there was a person there. How long did the festival last for? 1
- (d) For the parabola  $(y-1)^2 = 16-8x$
- (i) State the coordinates of the vertex and the focus. 2
- (ii) Sketch the graph of the parabola showing the information above 1


**SECTION D (Use a SEPARATE writing booklet)**

Question 7 (12 marks)

Marks

(a)

3



The diagram above shows the shaded region bounded by the curve  $y = x^2 + 3$ , the lines  $x = 1$ ,  $x = 0$  and the  $x$  axis. This region is rotated  $360^\circ$  about the  $y$  axis.

Find the volume generated.

(b)

At time  $t$ , the mass  $M$  of a material decaying radioactively is given by  $M = 5e^{-0.1t}$ .

- (i) If at time  $t_1$ , the mass is  $M_1$  and at time  $t_2$  the mass is  $\frac{1}{2}M_1$ , show that


2

$$t_2 - t_1 = 10 \ln 2$$

- (ii) Calculate the time taken for the initial mass to reduce to a mass of  $\frac{5}{32}$ .

2

(c)



The spinner above is used in a game. *Once spun*, it is equally likely to stop at any one of the letters **A**, **E**, **I**, **O** or **U**.

- (i) If the spinner is spun twice, find the probability that it stops on the same letter twice.
- (ii) How many times must the spinner be spun for it to be 99% certain that it will stop on the letter **E** at least once?

2

3

Question 8 (12 marks)

Marks


- (a) The velocity  $v$  (in km/min) of a train travelling from Olympack Park to Lydcome, non-stop, is given by  $v = 20t^2(3-t)$ , where  $t$  is the time (in minutes) during which the train has been in motion between the two stations.  
Find:
- (i) The acceleration of the train at the end of the second minute. 1
- (ii) Find an expression for the displacement  $x$  km of the train from Olympic Park. 2
- (iii) Hence calculate the distance travelled from Olympic Park to Lidcombe. 1
- (iv) Where and when, between the two stations, was the train travelling the fastest? 2
- (b) (i) Show  $\int_0^1 \frac{dx}{1+x} = \ln 2$  1
- (ii) By using Simpson's rule with five function values, find an approximation to  $\ln 2$ . 2
- (c) Yddap is given on his 18<sup>th</sup> birthday a present of \$500 from his grandparents. 3
- Yddap immediately deposits this into his Credit Union account. His Credit Union gives him a return of 4% pa, compounded annually.
- Each birthday from then on, Yddap decides to deposit \$500 into the same account. He does this up until his 39<sup>th</sup> birthday.
- His last deposit of \$500 is on his 39<sup>th</sup> birthday and when Yddap turns 40 he decides to transfer the total of this investment to another account.
- How much does Yddap transfer?

**SECTION E (Use a SEPARATE writing booklet)**

Question 9 (12 marks)

Marks

(a)



The above graph shows the velocity,  $v \text{ ms}^{-1}$ , of a particle moving on a straight line, for  $0 \leq t \leq 9$ .

- (i) State all the times, or intervals of time, for which the particle
- |              |                                      |   |
|--------------|--------------------------------------|---|
| ( $\alpha$ ) | is at rest,                          | 1 |
| ( $\beta$ )  | is moving in the positive direction, | 1 |
| ( $\gamma$ ) | the acceleration is positive,        | 1 |
| ( $\delta$ ) | is slowing down.                     | 1 |
- (ii) Using the graph, determine whether the particle has returned to its starting point when  $t = 9$ . Justify your answer. 2


Question 9 continues on page 11

Question 9 continued

Marks

(b) (i)


2



The diagram above shows triangle  $ABC$ .  
 $X$  is a point on  $AB$  such that  $AX = XB = XC$ .

Prove  $\angle ACB = 90^\circ$

(ii)



$AB$  is a diameter of the circle  $ABC$  whose centre is  $O$ .

$C$  is equidistant from  $A$  and  $B$ .

The arc  $AB$  is drawn with  $C$  as centre.

( $\alpha$ ) If the radius of the circle is  $r$ , using (i) show that  $AC = r\sqrt{2}$ . 1

( $\beta$ ) Hence show that the shaded area is equal to the area of the triangle  $ABC$ . 3

Question 10 starts on page 12

**SECTION E continued**

Question 10 (12 marks)

Marks

A derrick crane is used to lift and move a heavy block across flat ground.

The crane consists of a fixed vertical mast of height  $m$ , a boom of fixed length  $b$  hinged at the base of the mast, and a hoist rope.




Figure 1

Figure 1 above shows the block is at  $H_1$  on the ground. The hoist rope, anchored at  $E$ , passes over pulleys at  $F$  and  $G_1$ , then reaches vertically downwards and is attached to the block.




Figure 2

The length of the rope remains constant during the subsequent manoeuvre.

Figure 2 above shows that as the boom is lowered to  $G_2$ , the block moves outwards to  $H_2$ .

Let  $\theta = \angle FEG_2$ ,  $0 < \theta < \frac{\pi}{2}$  and let  $y$  be the height of the block above the ground.

Assume that  $b < 2m$  and that the ground is level. Ignore the size of the pulleys.

- (i) If  $R$  is the length of the rope, show that 2

$$y = m + b \cos \theta + \sqrt{b^2 + m^2 - 2bm \cos \theta} - R$$

- (ii) Show that 3

$$\frac{dy}{d\theta} = \frac{bm \sin \theta}{\sqrt{b^2 + m^2 - 2bm \cos \theta}} - b \sin \theta,$$

Question 10 continues on page 13

Question 10 continued

Marks

(iii) Show that when  $\frac{dy}{d\theta} = 0$  then either  $\cos \theta = \frac{b}{2m}$  or  $\theta = 0$ . 4

(iv) Assume that  $y$  is a maximum when  $\cos \theta = \frac{b}{2m}$ . 3

The horizontal distance from the mast to the vertical rope is called the *lifting radius* of the crane.

Find the lifting radius in terms of  $m$  and  $b$  when  $y$  is a maximum.

**End of paper**

**STANDARD INTEGRALS**

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax,$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left( x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left( x + \sqrt{x^2 + a^2} \right)$$

NOTE:  $\ln x = \log_e x, \quad x > 0$



**SYDNEY BOYS HIGH SCHOOL**  
MOORE PARK, SURRY HILLS

**2004**

**TRIAL HIGHER SCHOOL  
CERTIFICATE EXAMINATION**

# Mathematics

## Sample Solutions

<b>Section</b>	<b>Marker</b>
<b>A</b>	Mr Bigelow
<b>B</b>	Mr Hespe
<b>C</b>	Mr Choy
<b>D</b>	Mr Fuller
<b>E</b>	Mr Gainford

QUESTION 1.


(a)  $\log(\tan 514) = \boxed{1.32}$  ✓✓

(b)  $y = (5x)^{\frac{1}{2}}$   
 $y' = \frac{1}{2}(5x)^{-\frac{1}{2}} \times 5$   
 $= \boxed{\frac{5}{2\sqrt{5x}}} \checkmark \checkmark \left( \frac{\sqrt{5}}{2} x^{-\frac{1}{2}} \right)$

(c)  $2x^2 - x - 15 = 0$   
 $(2x+5)(x-3) = 0$   
 $\boxed{x = 3, -\frac{5}{2}} \checkmark \checkmark$

(d)  $y' = 3 - 2x$   
 $\boxed{y = 3x - x^2 + c} \checkmark \checkmark$

(e)  $y = 2x \quad \text{--- (1)}$   
 $3x + 2y = 14 \quad \text{--- (2)}$   
Sub (1) into (2)  
 $3x + 4x = 14$   
 $7x = 14$   
 $x = 2$   
Sub in (1)  
 $y = 4$   
 $\therefore \text{solv. is } \boxed{(2, 4)} \checkmark \checkmark$

(f)  $3 - 4x < 1$   
 $-4x < -2$   
 $\boxed{x > \frac{1}{2}} \checkmark \checkmark$   


QUESTION 2.

(a)(i)  $f(x) = (1 + \cos 2x)^3$

$\therefore f'(x) = 3(1 + \cos 2x)^2 \times -2 \sin 2x$   
 $= \boxed{-6 \sin 2x (1 + \cos 2x)^2}$  ✓✓

(ii)  $f(x) = x^2 e^{x+2}$

$f'(x) = x^2 e^{x+2} + 2x e^{x+2}$   
 $= \boxed{x e^{x+2} (x+2)}$  ✓✓

(b)(i)  $\int \frac{\cos x}{\sin x} dx = \boxed{\ln|\sin x| + c}$  ✓

(ii)  $\int_{\frac{1}{2}}^2 \left(1 - \frac{1}{x^2}\right) dx = \int_{\frac{1}{2}}^2 (1 - x^{-2}) dx$   
 $= \left[ x + x^{-1} \right]_{\frac{1}{2}}^2$   
 $= \left(2 + \frac{1}{2}\right) - \left(\frac{1}{2} + 2\right)$   
 $= \boxed{0}$  ✓✓

(c)(i)  $\widehat{NBY} = \widehat{MXC}$  (vertically opposite angles)

$\angle$  is common

$\therefore \triangle NBY \sim \triangle MXY$  (equiangular)

$\therefore \left[ \frac{MX}{NB} = \frac{1}{2} \right]$  (ratio of corresponding sides equal)

$\left[ \frac{NB}{NY} = \frac{MY}{NY} = \frac{1}{2} \right]$  (M is the midpt of NY)

(ii)  $\widehat{MCX} = \widehat{ABC}$  (base angles of an isosceles triangle)

$\widehat{MXC} = \widehat{ABC}$  (vertically opposite angles)

$\therefore \widehat{MCX} = \widehat{MXC}$

$\therefore \triangle XMC$  is isosceles  $\therefore MX = MC$   $\therefore \left[ \frac{MX}{NB} = \frac{MC}{NB} = \frac{1}{2} \right]$  ✓

$$(d) \quad g'(x) = 4 - 3x^2$$

$$\therefore g(x) = 4x - x^3 + C.$$

$$\text{Now } 4 = 4 \times 2 - 8 + C.$$

$$\therefore C = 4$$

$$\therefore \boxed{g(x) = 4x - x^3 + 4} \quad \checkmark \checkmark$$

$$3(i) \quad m_{AD} = \frac{-3-5}{0-5}$$

$$= -8/5 \quad \checkmark$$

$$\theta = \tan^{-1}(-8/5)$$

$$= 180^\circ - 58^\circ$$

$$= 122^\circ \quad \checkmark$$

$$(ii) \quad M = \left( \frac{-5+3}{2}, \frac{1+5}{2} \right)$$

$$= (-1, 3) \quad \checkmark$$

$$(iii) \quad C = (3 + (-5 - 0), 1 + (5 - (-3)))$$

$$= (-2, 9) \quad \checkmark$$

$$(iv) \quad m_{AB} = \frac{1-3}{3-0}$$

$$= 4/3 \quad \checkmark$$

$$\therefore \text{Eqn of AB is } y = \frac{4x}{3} - 3 \quad \checkmark$$

$$\text{or } 4x - 3y - 9 = 0.$$

$$(v) \quad \text{Distance} = \frac{|4 \times (-5) - 3 \times 5 - 9|}{\sqrt{4^2 + 3^2}}$$


$$= \frac{44}{5} \quad \checkmark \quad (\text{or } 8.8).$$

3 (vi) Length  $AB = \sqrt{9+16}$   
 $= 5 \checkmark$

$\therefore \text{Area} = \frac{4}{5} \times 5$   
 $= 4 \checkmark$

(vii)  $x+2y < 5 \checkmark \cap x > 0 \cap 4x-3y < 9 \checkmark$

4(a)  $2x^\circ = 120^\circ, 240^\circ, 480^\circ, 600^\circ$   
 $x^\circ = 60^\circ, 120^\circ, 240^\circ, 300^\circ \checkmark$



(b)(i)  $\sqrt{x-5} = \frac{x-5}{5}$

$25x - 125 = x^2 - 10x + 25$

$x^2 - 35x + 150 = 0$

$(x-5)(x-30) = 0 \checkmark$

$x = 5, 30$


So P is  $(30, 5) \checkmark$

(ii) Area =  $\int_5^{30} \left\{ (x-5)^{1/2} - \left( \frac{x-5}{5} \right) \right\} dx \checkmark$


$= \left[ \frac{2}{3} (x-5)^{3/2} - \frac{x^2}{10} + x \right]_5^{30} \checkmark$

$= \frac{2}{3} (125 - 0) - \left( \frac{900}{10} - \frac{25}{10} \right) + 30 - 5$

$= \frac{125}{6} \checkmark$  or  $20 \frac{5}{6}$  or  $20.83$



$\hat{B}ED = 68^\circ$  (vert. opp.  $\angle$ s)  $\checkmark$   
 $\hat{B}DE = 22^\circ$  (angle sum of  $\Delta$ )  $\checkmark$   
 $\hat{D}CB = 22^\circ$  (base angle of isosceles  $\Delta$ )  $\checkmark$



(i)  $AC^2 = 4^2 + 2^2 - 2 \times 4 \times 2 \times \cos 60^\circ \text{ m}^2$

$= 16 + 4 - 16 \times \frac{1}{2} \text{ m}^2$

$= 12 \text{ m}^2$

$AC = 2\sqrt{3} \text{ m.} \checkmark$

(3.464101615...)

$$4(d)(ii) \frac{\sin \hat{BCA}}{2} = \frac{\sin 60^\circ}{2\sqrt{3}}$$

$$\sin \hat{BCA} = \frac{1}{2}$$

$$\hat{BCA} = 30^\circ \quad \checkmark$$

$$\hat{ACD} = 180^\circ - 60^\circ - 30^\circ$$

$$= 90^\circ \quad \checkmark$$

$$\therefore AD^2 = 12 + 1$$

$$\therefore AD = \sqrt{13} \quad (3.605551275 \text{ calculator})$$

$$= 3.61 \text{ m} \quad (3 \text{ sig. fig.})$$

5  $y = x^3 - 5x^2 + 7x - 14$


(a) (i)  $\frac{dy}{dx} = 3x^2 - 10x + 7$  [1]  
 $= (3x-7)(x-1)$

(ii)  $\frac{dy}{dx} = 0, x = \frac{7}{3}, 1$

When  $x = 1, y = -11$  [1]  
 $x = \frac{7}{3}, y = -12\frac{5}{27}$  (-12.185) [1]

3  $\frac{d^2y}{dx^2} = 6x - 10$

$f''(1) = -4 < 0$  (1, -11) max [1]  
 $f''(\frac{7}{3}) = 4 > 0$  ( $\frac{7}{3}, -12\frac{5}{27}$ ) min (-12.2 (1dp))



2

(iv)  $f''(x) < 0, 6x - 10 < 0, \begin{cases} 6x < 10 \\ x < 5/3 \end{cases}$

(b)  $a, a+d, a+2d, \dots, a+39d$

$S_{40} = \frac{40}{2} [2a + 39d]$

$\therefore 495 = 20(2a + 39d)$

$99 = 4(2a + 39d)$  — (1)

$a + 39d = 2a$  — (2)

$d = 39d \therefore d = \frac{a}{39}$  — (3)

Subst (3) into (1)

$99 = 4(2a + 39 \times \frac{a}{39})$

$\therefore 99 = 4(3a) \Rightarrow a = \frac{99}{4}$  (8.25)

$\Rightarrow d = \frac{11}{52} \text{ cm}$  (0.212 cm) [2]

$\frac{495}{2340}$


Q (6). (a)  $\lambda = 10\theta$   
 $\therefore 12 = 10\theta$   $\theta = 1.2^\circ$

$l^c = \frac{180}{\pi} \therefore 1.2^c = \frac{180 \times 1.2}{\pi} \approx 69^\circ$

(b) (i)  $t = \cos^2 x$  ( $< 1$ ). [1]

(ii)  $s = \frac{\cos^2 x}{1 - \cos^2 x} = \frac{\cos^2 x}{\sin^2 x} = \cot^2 x$ . [2]

(c).  $\frac{dN}{dt} = 450t(8-t)$



(i)  $\frac{dN}{dt} (\text{max})$   
 $= 450 \times 4 \times 4$   
 $= 7200$  [1] When  $t=4$

$\frac{dN}{dt} = 3600t - 450t^2$

$N = 1800t^2 - 150t^3 + c$  ✓

When  $t=0, N=0, \Rightarrow c=0$


(ii)  $\therefore N = 150t^2(12-t)$ . [2]

(iii)  $N=0$ , when  $t=0$ , or  $12$ . [1]  
 $\therefore$  festival last for 12 hrs.

(d)  $(y-1)^2 = -8(x-2)$

$4a = 8, a = 2$

3



**SECTION D**

**Question 7**

(a) when  $x=0$   $x=1$   
 $y=3$   $y=4$

$$V = \pi r^2 h - \pi \int_3^4 x^2 dy$$

$$= \pi(1)^2(4) - \pi \int_3^4 (y-3) dy$$

$$= 4\pi - \pi \left[ \frac{1}{2}y^2 - 3y \right]_3^4$$

$$= 4\pi - \pi \left\{ \left[ \frac{1}{2}(4)^2 - 3(4) \right] - \left[ \frac{1}{2}(3)^2 - 3(3) \right] \right\}$$

$$= 4\pi - \frac{\pi}{2}$$

$$= \frac{7\pi}{2} \text{ units}^3$$

(b)(i) when  $t = t_1$ ,  $M = M_1$ , ie  $M_1 = 5e^{-0.1t_1}$   
 when  $t = t_2$ ,  $M = \frac{1}{2}M_1$ , ie  $\frac{1}{2}M_1 = 5e^{-0.1t_2}$   
 $M_1 = 10e^{-0.1t_2}$

$$5e^{-0.1t_1} = 10e^{-0.1t_2}$$

$$e^{-0.1t_1} = 2e^{-0.1t_2}$$

$$-0.1t_1 = \ln(2e^{-0.1t_2})$$

$$-0.1t_1 = \ln(2) + \ln(e^{-0.1t_2})$$

$$-0.1t_1 = \ln(2) + -0.1t_2$$

$$t_1 = -10\ln(2) + t_2$$

$$t_2 - t_1 = 10\ln(2)$$

(ii)  $\frac{5}{32} = 5e^{-0.1t}$   
 $e^{-0.1t} = \frac{1}{32}$   
 $-0.1t = \ln\left(\frac{1}{32}\right)$   
 $t = -10\ln\left(\frac{1}{32}\right)$   
 $t = 34.657$  correct to 3 decimal places

(c)(i) P(same letter twice) =  $1 \times \frac{1}{5}$

(ii) P(E at least once) =  $1 - P(\text{no E})$

$$= 1 - \left(\frac{4}{5}\right)^n$$

$$P(\text{E at least once}) \geq \frac{99}{100}$$

$$1 - \left(\frac{4}{5}\right)^n \geq \frac{99}{100}$$

$$\left(\frac{4}{5}\right)^n \leq \frac{1}{100}$$

$$n \ln\left(\frac{4}{5}\right) \leq \ln\left(\frac{1}{100}\right)$$

$$n \geq \frac{\ln\left(\frac{1}{100}\right)}{\ln\left(\frac{4}{5}\right)}$$

$$n \geq 20.6377\dots$$

$$\therefore n = 21$$

**Question 8**

(a)  $v = 20t^2(3-t)$   $v = 60t^2 - 20t^3$

(i)  $a = \frac{dv}{dt}$   
 $a = \frac{d(60t^2 - 20t^3)}{dt}$

$$a = 120t - 60t^2$$

when  $a = 2$ ,  $a = 120(2) - 60(2)^2$   
 $a = 0$

(ii)  $x = \int \frac{dx}{dt} dt$

$$x = 20t^3 - 5t^4 + C$$

Note:  $t = 0, x = 0 \therefore C = 0$

Hence,  $x = 20t^3 - 5t^4$

(iii) Let  $v = 0$ .

$$0 = 20t^2(3-t)$$

$$t = 0 \text{ or } t = 3$$

when  $t = 3$ ,  $x = 20(3)^3 - 5(3)^4$



SECTION E

Question 9

(a) (i) (a) At rest  $t=0$   
 $t=5$   
 $t=9$


(b)  $0 < t < 5$  [1]

(c)  $0 < t < 2, 7 < t < 9$  [1]

\* (d)  $2 < t < 5, 7 < t < 9$  [1]

(ii) No, it has not returned.

The area under the curve is the measure of distance travelled. The negative area (5, 9) is less than the positive area (0, 5).



Aim To prove  $\angle ACB = 90^\circ$

Proof let  $\angle ABC = \alpha$   
 $\angle BAC = \beta$

Now  $\angle BCX = \alpha$  (Isos  $\Delta$ )

and  $\angle ACX = \beta$  ( " )


But  $\angle BAC + \angle ACB + \angle CBA = 180^\circ$   
 (angles in a triangle)

$$\therefore \beta + (\beta + \alpha) + \alpha = 180^\circ$$

$$2(\alpha + \beta) = 180^\circ$$

$$\alpha + \beta = 90^\circ$$

$$\therefore \angle ACB = 90^\circ \quad [2]$$



(a) In  $\Delta ABC$ ,  $AO = BO = CO$   
 (radii)

$$\therefore \angle ACB = 90^\circ$$

Since  $AO = r$ , in  $\Delta ABC$ , using Pythagoras

$$AB^2 = 2AC^2$$

$$(2r)^2 = 2AC^2$$

$$4r^2 = 2AC^2$$

$$AC^2 = 2r^2$$

$$\therefore AC = r\sqrt{2} \quad [1]$$

$$(b) \Delta_{ABC} = \frac{1}{2} (r\sqrt{2})^2 = r^2$$

$$A_{\text{shaded}} = \frac{1}{2} \pi r^2 - A_{\text{segment}}$$


$$= \frac{1}{2} \pi r^2 - \frac{1}{2} (r\sqrt{2})^2 \left( \frac{\pi}{2} - \sin \frac{\pi}{2} \right)$$

$$= \frac{1}{2} \pi r^2 - r^2 \left( \frac{\pi}{2} - 1 \right)$$

$$= r^2$$

$$= \Delta_{ABC} \quad \text{Q.E.D.} \quad [3]$$

Question 10



$$(i) R = FG_2 + G_2K - y$$

$$\therefore y = FG_2 + G_2K - R$$

$$= \sqrt{b^2 + m^2 - 2bm \cos \theta} + b \cos \theta - R$$

as required. [2]

$$(ii) \frac{dy}{d\theta} = -b \sin \theta + \frac{1 \times 2bm \sin \theta}{2\sqrt{b^2 + m^2 - 2bm \cos \theta}}$$

$$= \frac{bm \sin \theta}{\sqrt{b^2 + m^2 - 2bm \cos \theta}} - b \sin \theta$$

[3]

(iii) When  $\frac{dy}{d\theta} = 0$

$$0 = \frac{bm \sin \theta - b \sin \theta \sqrt{b^2 + m^2 - 2bm \cos \theta}}{\sqrt{b^2 + m^2 - 2bm \cos \theta}}$$

$$\therefore \sin \theta = b \sin \theta (m - \sqrt{b^2 + m^2 - 2bm \cos \theta})$$

$\therefore$  Either  $\sin \theta = 0$   $\hat{=}$   $\theta = 0$

$$\text{or } m - \sqrt{b^2 + m^2 - 2bm \cos \theta} = 0$$

$$\hat{=} m = \sqrt{b^2 + m^2 - 2bm \cos \theta}$$

That is  $b^2 = 2bm \cos \theta$

$$\hat{=} \cos \theta = \frac{b}{2m}$$

[4]

(2)

(ii) [Clearly  $y$  is min when  $\theta = 0$ ]

$\therefore y$  is max when  $\cos \theta = \frac{b}{2m}$

Now hypotenuse radius is  $b \sin \theta$

$$= b \sqrt{1 - \cos^2 \theta}$$

$$= b \sqrt{1 - \frac{b^2}{4m^2}}$$

$$= \frac{b}{2m} \sqrt{4m^2 - b^2}$$

[3]