



SYDNEY GRAMMAR SCHOOL  
MATHEMATICS DEPARTMENT  
TRIAL EXAMINATIONS 2007

# FORM VI

# MATHEMATICS

## Examination date

Wednesday 1st August 2007

## Time allowed

3 hours (plus 5 minutes reading time)

## Instructions

- All ten questions may be attempted.
- All ten questions are of equal value.
- All necessary working must be shown.
- Marks may not be awarded for careless or badly arranged work.
- Approved calculators and templates may be used.
- A list of standard integrals is provided at the end of the examination paper.

## Collection

- Write your candidate number clearly on each booklet.
- Hand in the ten questions in a single well-ordered pile.
- Hand in a booklet for each question, even if it has not been attempted.
- If you use a second booklet for a question, place it inside the first.
- Keep the printed examination paper and bring it to your next Mathematics lesson.
- Bundle the separate sheet with Question 10.

## Checklist

- Folded A3 booklets: 10 per boy. A total of 1250 booklets should be sufficient.
- Candidature: 93 boys.

## Examiner

SJE/LYL

**QUESTION ONE** (12 marks) Use a separate writing booklet.


Marks

- (a) Evaluate  $\sqrt[3]{43.1 - 16.98}$  correct to one decimal place. 2
- (b) Solve  $x^2 = 3x$ . 2
- (c) Solve  $|x + 1| = 5$ . 2
- (d) Differentiate  $3x + \cos 2x$ . 2
- (e) Write down the exact value of  $\tan \frac{3\pi}{4}$ . 1
- (f) Find a primitive function of  $\frac{5}{x}$ . 1
- (g) Find  $a$  and  $b$  if  $a + b\sqrt{2} = (3 + \sqrt{2})(2 - \sqrt{2})$ . 2

**QUESTION TWO** (12 marks) Use a separate writing booklet.

Marks

(a)



In the diagram above  $ABCD$  is a square.

- (i) Find the gradient of  $AC$ . 1
- (ii) Show that the equation of  $BD$  is  $7x - 3y - 7 = 0$ . 2
- (iii) Find  $q$ , the  $y$ -coordinate of  $B$ . 1
- (iv) Find the length of  $AC$ . 1
- (v) Hence, or otherwise, find the area of  $ABCD$ . 1

- (b) (i) Evaluate  $\sum_{n=2}^4 \frac{n}{n+1}$ . 1
- (ii) Evaluate  $\int_1^2 e^x dx$ . 1
- (c) Find the limiting sum of the geometric series  $500 + 100 + 20 + 4 + \dots$ . 2
- (d) Find  $\int \frac{x}{2x^2 + 3} dx$ . 2

**QUESTION THREE** (12 marks) Use a separate writing booklet.


Marks

- (a) Differentiate with respect to  $x$ :
- (i)  $e^{x^2-9}$  1
- (ii)  $x^2 \tan 5x$  2
- (b) Consider the parabola  $(x - 1)^2 = -6(y + 4)$ .
- (i) Write down coordinates of the vertex. 1
- (ii) Find the equation of the directrix. 2
- (c) Evaluate  $\int_{\frac{\pi}{3}}^{\frac{3\pi}{2}} 2 \cos x dx$ , leaving your answer as an exact value. 2
- (d) Find the sum of the arithmetic series  $1 + 4 + 7 + \dots + 226$ . 2
- (e) Find  $k$  such that  $\int_1^k \frac{dx}{x} = 2$ . 2

**QUESTION FOUR** (12 marks) Use a separate writing booklet.

Marks

(a)



In the diagram above, triangle  $ABC$  has dimensions  $AB = 7$  cm and  $BC = 15$  cm. The point  $D$  lies on  $AC$  such that  $AD = 5$  cm and  $BD = 3$  cm.

- (i) Use the cosine rule to show that  $\angle ADB = 120^\circ$ . 2
- (ii) Show that  $\angle BCD = 10^\circ$  (rounded to the nearest degree). 2
- (iii) Find the length of  $DC$ , correct to the nearest millimetre. 2
- (b) The roots of the quadratic equation  $px^2 - x + q = 0$  are  $-1$  and  $3$ . Find  $p$  and  $q$ . 3
- (c) Find the equation of the normal to the curve  $y = (2 - x)^3$  at the point where  $x = 0$ . 3

**QUESTION FIVE** (12 marks) Use a separate writing booklet.

Marks

- (a) A particle moves in a straight line. At time  $t$  seconds, its displacement  $x$  metres from the origin is given by  $x = 3 \cos \frac{t}{2}$ , where  $0 \leq t \leq 4\pi$ .
  - (i) Sketch the graph of  $x$  as a function of  $t$ . 2
  - (ii) Find the times at which the particle is at rest. 2
  - (iii) What is the particle's initial displacement and acceleration? 3
  - (iv) Find the total distance travelled by the particle. 1
- (b) Consider the function  $f(x) = \frac{x^2}{1 + x^2}$ .
  - (i) Show that  $f''(x) = \frac{2(1 - 3x^2)}{(1 + x^2)^3}$ . 3
  - (ii) For what values of  $x$  is the function concave up? 1

**QUESTION SIX** (12 marks) Use a separate writing booklet.

Marks

- (a) The table below shows the value of a function  $f(x)$  for three values of  $x$ .

2

$x$	3	4	5
$f(x)$	$\sqrt{7}$	$\sqrt{14}$	$\sqrt{23}$

Use the trapezoidal rule with the three given function values to find an approximation of  $\int_3^5 f(x) dx$ . Give your answer correct to one decimal place.

- (b) A ball is rolled up an inclined plane and is subject to an acceleration of  $a = -6 \text{ m/s}^2$ . Initially the ball has a velocity of  $v = 12 \text{ m/s}$  and its displacement, measured from the bottom of the plane, is 36 m.

(i) Show that the velocity function is  $v = 12 - 6t$ .

1

(ii) Find the displacement as a function of time.

2

(iii) When does the ball reach the bottom of the plane and what is its speed then?

2

- (c) The fruit bat population in the Sydney Botanical Gardens has been increasing according to the equation  $P = Ae^{kt}$ , where  $A$  and  $k$  are constants. On 1st April 2005 there were 4800 bats, and by 1st April 2007 there were 10 800.

(i) Show that  $k = \log_e \frac{3}{2}$ .

2

(ii) If the trend continues without any intervention, how many bats will inhabit the gardens by 1st April 2010?

1


(iii) During what year is the bat population increasing at a rate of 6500 bats per year?

2

**QUESTION SEVEN** (12 marks) Use a separate writing booklet.

**Marks**

- (a) Find the area between the curves  $y = x^2$  and  $x = y^2$ . 3
- (b) Alex decides to invest \$30 000 into an investment fund offering 9% p.a. interest compounded monthly. How many months will it be before his money has doubled? Give your answer correct to the nearest month. 3
- (c) Gabriel's Horn is formed by rotating the area enclosed by the curve  $y = \frac{1}{x}$  and the  $x$ -axis, between  $x = 1$  and  $x = a$ , around the  $x$ -axis.
  - (i) Find the volume of the horn when  $a = 5$ . 2
  - (ii) Find the limiting value of the volume as  $a$  gets larger. 1
- (d)



In the diagram above,  $ABC$  is a Reuleaux Triangle. Its sides are equal arcs of congruent circles centred at  $A$ ,  $B$  and  $C$ . The radius of each circle is 12 cm. Find:

- (i) the perimeter of the Reuleaux Triangle, 1
- (ii) the exact area of the Reuleaux Triangle. 2

**QUESTION EIGHT** (12 marks) Use a separate writing booklet.

Marks

(a) Karen borrows \$15 000 from the bank. The loan plus interest and charges are to be repaid at the end of each month in equal monthly instalments of \$ $M$  over 5 years. Interest is charged at 6% p.a. and is calculated on the balance owing at the beginning of each month. Furthermore, at the end of each month a bank charge of \$15 is added to the account.

Let  $A_n$  be the amount owing after  $n$  months.

- (i) Write down expressions for  $A_1$  and  $A_2$  and show that the amount owing after three months is given by 3

$$A_3 = 15\,000 \times 1.005^3 - (M - 15)(1 + 1.005 + 1.005^2).$$

- (ii) Hence write an expression of  $A_n$ . 1

- (iii) Find the monthly instalment, correct to the nearest cent. 3

(b) The line  $x - 4y + 2 = 0$  is a tangent to the parabola  $x = Ay^2$ , where  $A$  is a constant.


- (i) Form a quadratic equation and hence show that  $A = 2$ . 2

- (ii) Draw a neat sketch of the parabola and the tangent, showing the point of contact. 3

**QUESTION NINE** (12 marks) Use a separate writing booklet.

Marks


- (a) (i) Show that  $\frac{d}{dx}(x \ln x - x) = \ln x$ . 1



NOT TO SCALE

The shaded region in the diagram above is bounded by the curve  $y = \log_e x$ , the line  $y = 3$  and the coordinate axes. Using the result in part (i), or otherwise, find the exact area of the region. 2

(b)



In the diagram above,  $ABCD$  and  $AEFD$  are parallelograms.  $AE$  bisects  $\angle DAB$  and  $DC$  bisects  $\angle FDA$ . Let  $\angle DAG = \alpha$  and  $\angle FDG = \beta$ .


(i) Show that  $\triangle AGD$  is equilateral, giving reasons. 3

(ii) If  $AD = 2$  units and  $AB = 7$  units, show that the area of the trapezium  $ABFD$  is  $\frac{77\sqrt{3}}{4}$  square units. 3

(c) Show that if  $y = \frac{e^x + e^{-x}}{2}$  then  $y'' = \sqrt{1 + (y')^2}$ . 3

**QUESTION TEN** (12 marks) Use a separate writing booklet. Marks

(a)



The square  $ABCD$  is shown above. The diagonals  $AC$  and  $BD$  intersect at  $M$ .  $P$  is a point on the diagonal  $AC$  between  $M$  and  $C$ , and  $P$  is joined to  $D$ . The point  $X$  is chosen on  $DP$  so that  $AX \perp DP$ , and  $AX$  intersects the diagonal  $DB$  at  $Q$ . Let  $\angle DAQ = \theta$ .


The diagram has been reproduced on a separate sheet which can be used for your solution to this question. Insert this sheet with the rest of Question 10.

(i) Show that  $\angle PDC = \theta$ . 1

(ii) Hence show that  $\triangle ADQ \cong \triangle DCP$ . 3

(iii) Deduce that  $\triangle DXQ \parallel \triangle AMQ$ . 2

(b)




The diagram above shows that the distance between a boy's home  $H$  and his school  $S$  is 6 km. A canal  $ABCD$  is 1 km from both his home and school. In winter the canal is frozen, so he take an alternate route  $HBCS$ , walking  $HB$ , skating  $BC$  and walking  $CS$ . His walking speed is 4 km/h and his skating speed is 12 km/h. Let  $\angle AHB = \angle DSC = \theta$ .

(i) Show that the time taken for this alternate route is  $T = \frac{1}{2 \cos \theta} + \frac{1}{2} - \frac{\tan \theta}{6}$ . 2

(ii) Find, to the nearest minute, the value of  $\theta$  which minimises the time taken for the journey to school. 4

**END OF EXAMINATION**

Candidate Number .....



Q1

a)  $\sqrt[3]{26 \cdot 12} \doteq 2.967$  (2.9, etc 1 mark)  
 $\doteq 3.0$  (1dp) ✓✓

b)  $x^2 = 3x$   
 $x^2 - 3x = 0$

$x(x-3) = 0$   
 $x = 0$  ✓  $x - 3 = 0$   
 $x = 3$  ✓

c)  $|x+1| = 5$   
 $x+1 = 5$  or  $x+1 = -5$   
 $x = 4$  ✓ or  $x = -6$  ✓

d)  $3 - 2 \sin 2x$

e)  $\tan \frac{3\pi}{4} = -1$  ✓

f)  $5 \log_e x + c$  ✓

g)  $(3 + \sqrt{2})(2 - \sqrt{2})$   
 $= 6 - 3\sqrt{2} + 2\sqrt{2} - 2$   
 $= 4 - \sqrt{2}$   
 $a = 4$   $b = -1$  ✓

Q2

a) i) gradient AC =  $\frac{2-5}{6--1}$   
 $= -\frac{3}{7}$  ✓

ii) gradient of BD =  $\frac{7}{3}$  ✓

equation of BD

$\frac{y-0}{x-1} = \frac{7}{3}$  ✓

$y = \frac{7(x-1)}{3}$

$7x - 3y - 7 = 0$

iii) Sub  $x=4$  into  $7x - 3y - 7 = 0$   
 $28 - 3y - 7 = 0$   
 $21 = 3y$   
 $y = 7$  ✓

iv)  $AC^2 = (6--1)^2 + (2-5)^2$   
 $= 49 + 9$   
 $= 58$   
 $AC = \sqrt{58}$  units ✓

v) Area ABCD =  $\frac{1}{2} \times (\sqrt{58})^2$  ✓  
 $= 29$  square units

b) i)  $\sum_{n=2}^4 \frac{n}{n+1} = \frac{2}{3} + \frac{3}{4} + \frac{4}{5}$   
 $= \frac{133}{60}$  or  $2\frac{13}{60}$  ✓

ii)  $\int_1^2 e^x dx = [e^x]_1^2$   
 $= 2e - e$   
 $= e$  ✓

c)  $a = 500$   
 $r = \frac{1}{5}$  } either ✓

$S_{\infty} = \frac{a}{1-r}$   
 $= \frac{500}{1-\frac{1}{5}}$   
 $= 625$  ✓

d)  $\int \frac{x}{2x^2+3} dx = \frac{1}{4} \int \frac{4x}{2x^2+3} dx$  ✓  
 $= \frac{1}{4} \log_e (2x^2+3) + c$  ✓

### Question 3

$$2) \text{ (i) } \frac{d}{dx} (e^{x^2-9}) = 2xe^{x^2-9} \quad \checkmark$$


$$\text{(ii) } \frac{d}{dx} (x^2 \tan 5x) = 2x \tan 5x + x^2 \cdot 5 \sec^2 5x \quad \checkmark \\ = 2x \tan 5x + 5x^2 \sec^2 5x \quad \checkmark$$

$$2) (x-1)^2 = -6(y+4)$$

(i) Vertex is  $(1, -4)$   $\checkmark$

(ii)  $4a = 6$   
 $a = \frac{3}{2}$   $\checkmark$

directrix is  $y = -4 + \frac{3}{2}$   $\checkmark$   
 $y = -\frac{5}{2}$   $\checkmark$



$$2) \int_{\frac{\pi}{3}}^{\frac{3\pi}{2}} 2 \cos x \, dx = \left[ 2 \sin x \right]_{\frac{\pi}{3}}^{\frac{3\pi}{2}} \quad \checkmark \\ = -2 - 2 \frac{\sqrt{3}}{2} \quad \checkmark \\ = -2 - \sqrt{3} \quad \checkmark$$

$$2) 1 + 4 + 7 + \dots + 226$$

$a = 1$   
 $d = 3$

$$T_n = a + (n-1)d$$

So  $226 = 1 + (n-1)3$   
 $n-1 = 75$   
 $n = 76$   $\checkmark$

$$\text{Sum of series is } = \frac{n}{2} (a+l) \quad \checkmark \\ = \frac{76}{2} (1+226) \quad \checkmark \\ = 8626 \quad \checkmark$$

$$2) \int_1^k \frac{dx}{x} = 2 \quad \checkmark \\ \left[ \log_e x \right]_1^k = 2 \quad \checkmark \\ \log_e k - \log_e 1 = 2 \quad \checkmark \\ \log_e k = 2 \quad \checkmark \\ \therefore k = e^2 \quad \checkmark$$

$$a) i) \cos \angle ADB = \frac{5^2 + 3^2 - 7^2}{2 \times 5 \times 3} \quad \checkmark \checkmark$$

$$\angle ADB = 120^\circ$$

$$ii) \frac{\sin BCD}{3} = \frac{\sin 60}{15} \quad \checkmark$$

$$\sin BCD = \frac{3 \times \sin 60}{15} \quad \checkmark$$

$$BCD \doteq 10^\circ$$

( $170^\circ$  impossible as  $170^\circ + 60^\circ > 180^\circ$ )

$$iii) \frac{DC}{\sin 110^\circ} = \frac{3}{\sin 10^\circ} \quad \checkmark$$

$$DC = \frac{3 \sin 110^\circ}{\sin 10^\circ}$$

$$\doteq 16.2 \text{ cm} \quad \checkmark$$

$$b) -3 = \frac{q}{p} \quad \checkmark$$

$$2 = \frac{1}{p}$$

$$p = \frac{1}{2} \quad \checkmark$$

$$q = -\frac{3}{2} \quad \checkmark$$

$$c) y = (2-x)^3$$

$$y' = -3(2-x)^2 \quad \checkmark$$

$$m = -12$$


$$\text{normal } m = \frac{1}{12} \quad \checkmark$$

$$\text{when } x=0 \quad y=8$$

$$y-8 = \frac{1}{12}(x)$$

$$12y - 96 = x$$

$$x - 12y + 96 = 0 \quad \checkmark$$



5. a) i)  $\checkmark$  correct shape, amplitude or period.

$\checkmark$  remaining information correct and presented on the graph.

ii) At rest when  $t=0$   $\checkmark$  one correct  
 $t=2\pi$  or  $t=4\pi$   $\checkmark$  all three correct

iii) Initial displacement  $x=3\text{m}$   $\checkmark$

$$x = 3 \cos \frac{t}{2}$$

$$\dot{x} = -\frac{3}{2} \sin \frac{t}{2}$$

$$\ddot{x} = -\frac{3}{4} \cos \frac{t}{2} \quad \checkmark$$

$\therefore$  at  $t=0$  acceleration =  $-\frac{3}{4} \text{ ms}^{-2}$   $\checkmark$

iv) Total distance travelled =  $12\text{m}$   $\checkmark$

$$b) f(x) = \frac{x^2}{1+x^2}$$

$$f'(x) = \frac{(1+x^2)2x - x^2(2x)}{(1+x^2)^2} \quad \checkmark$$

$$= \frac{2x(1+x^2-x^2)}{(1+x^2)^2}$$

$$= \frac{2x}{(1+x^2)^2}$$

$$f''(x) = \frac{(1+x^2)^2 \cdot 2 - 2x \cdot 2 \cdot 2x(1+x^2)}{(1+x^2)^4} \quad \checkmark$$

$$= \frac{2(1+x^2)[(1+x^2)-4x^2]}{(1+x^2)^3}$$

$$= \frac{2(1-3x^2)}{(1+x^2)^3} \quad \checkmark$$

ii) Concave up when  $f''(x) > 0$

$$1-3x^2 > 0$$

$$x^2 < \frac{1}{3} \quad \checkmark$$

$$-\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}}$$

26

1)  $\int_a^b f(x) dx \equiv \frac{b-a}{2} (f(a) + f(b))$

$\int_3^5 f(x) dx = \int_3^4 f(x) dx + \int_4^5 f(x) dx$   
 $\equiv \frac{1}{2} (f(3) + f(4)) + \frac{1}{2} (f(4) + f(5))$   
 $\equiv \frac{1}{2} (\sqrt{7} + 2\sqrt{14} + \sqrt{23})$   
 $\equiv 7.5 \text{ (1dp)}$  ✓

b)  $a = -6$

i)  $\frac{dv}{dt} = -6$

$v = -6t + c$

$t=0 \quad v=12$

$c = 12$  ✓

$v = -6t + 12$

ii)  $\frac{dx}{dt} = -6t + 12$

$x = \frac{-6t^2}{2} + 12t + k$  ✓

$t=0 \quad x=36$

$k = 36$  ✓

$x = -3t^2 + 12t + 36$

iii)  $v=0$

$-3t^2 + 12t + 36 = 0$

$3t^2 - 12t - 36 = 0$

$t^2 - 4t - 12 = 0$

$(t+2)(t-6) = 0$

$t \neq -2 \quad t = 6$  ✓

$t = 6$

$v = 12 - 36$

$= -24 \text{ m/s}$

$\therefore \text{speed} = 24 \text{ m/s}$  ✓

c) i)  $P = Ae^{kt}$

$A = 4800$

$t = 2 \quad P = 10800$

$10800 = 4800 e^{2k}$  ✓

$\frac{108}{48} = e^{2k}$

$\frac{9}{4} = e^{2k}$

$\log \left(\frac{9}{4}\right)^2 = \log e^{2k}$  ✓ must show

$2 \log \frac{3}{2} = 2k$

$k = \log \frac{3}{2}$

ii)  $t = 5$

$r = 4800 e^{k5}$

$= 36450$  ✓

iii) 6500 bats / year

$\frac{dP}{dt} = kP$

$6500 = kP$

$P = \frac{6500}{k}$

$P = Ae^{kt}$

$\frac{6500}{k} = 4800 e^{kt}$  ✓

$\frac{6500}{4800k} = e^{kt}$

$\log e \left(\frac{65}{48k}\right) = kt$

$t = \frac{\log e \left(\frac{65}{48k}\right)}{k}$

$\doteq 2.97$

$\therefore$  During 2008 the bat population is 6500 bats per year. ✓

$t = 0 \quad 2005 \quad \left( \begin{array}{l} \text{1st April} \\ \text{1st year} \end{array} \right)$


$t = 1 \quad 2006 \quad \left( \begin{array}{l} \text{2nd year} \end{array} \right)$

$t = 2 \quad 2007 \quad \left( \begin{array}{l} \text{3rd year} \end{array} \right)$

$t = 3 \quad 2008$

### Question 7

(a) Area =  $\int_0^1 \sqrt{x} dx - \int_0^1 x^2 dx$  ✓  
 $= \left[ \frac{2}{3} x^{3/2} \right]_0^1 - \left[ \frac{x^3}{3} \right]_0^1$  ✓  
 $= \frac{2}{3} - \frac{1}{3}$  ✓  
 $= \frac{1}{3}$  square unit. ✓




(b)  $A = P(1+r)^n$   
 $60000 = 30000 (1.0075)^n$  ✓  
 $1.0075^n = 2$  ✓  
 $n = \frac{\log_e 2}{\log_e 1.0075}$

$P = \$30\,000$   
 $r = 9\% \text{ p.a.}$   
 $= 0.0075$  ✓  
 $A = \$60\,000$  (doubled)

∴ His money has doubled after 93 months

(c) (i)  $V = \pi \int_1^5 \left(\frac{1}{x}\right)^2 dx$   
 $= \pi \int_1^5 x^{-2} dx$  ✓  
 $= \pi \left[ -x^{-1} \right]_1^5$   
 $= \pi \left[ -\frac{1}{5} - (-1) \right]$  ✓  
 $= \frac{4}{5} \pi$  cubic units




(ii)  $V = \pi \left( -\frac{1}{a} - (-1) \right)$   
 $= \pi \left( 1 - \frac{1}{a} \right)$

as  $a \rightarrow \infty$ ,  $V \rightarrow \pi$  ✓ ∴ limiting value is  $\pi$  cubic units.

(d) (i) Perimeter =  $3 \times r\theta$   
 $= 3 \times 12 \frac{\pi}{3}$   
 $= 12\pi$  cm ✓

(ii) Area = Area Equilateral Triangle + 3 segments. ✓  
 $= \frac{1}{2} r^2 \sin \theta + 3 \left( \frac{1}{2} r^2 (\theta - \sin \theta) \right)$   
 $= \frac{144}{2} \sin \frac{\pi}{3} + \frac{3}{2} \cdot 144 \cdot \frac{\pi}{3} - \frac{3}{2} (144) \sin \frac{\pi}{3}$   
 $= 72 \frac{\sqrt{3}}{2} + 72\pi - 3(72) \frac{\sqrt{3}}{2}$   
 $= 72\pi - 72\sqrt{3}$   
 $= 72(\pi - \sqrt{3})$  cm<sup>2</sup> ✓



### Question 8.

a) 6% p.a = 0.005 per month

(i)  $A_1 = 15000(1.005) - M + 15$  ✓

$A_2 = 15000(1.005)^2 - M(1.005) + 15(1.005) - M + 15$

$A_3 = 15000(1.005)^3 - M(1.005)^2 - M(1.005) - M + 15(1.005)^2 + 15(1.005) + 15$  ✓

$= 15000(1.005)^3 - M(1.005^2 + 1.005 + 1) + 15(1.005^2 + 1.005 + 1)$  ✓

$= 15000(1.005)^3 - (M-15)(1.005^2 + 1.005 + 1)$  ✓

$= 15000(1.005)^3 - (M-15)(1 + 1.005 + 1.005^2)$  as required. ✓

(ii)  $A_n = 15000(1.005)^n - (M-15)(1 + 1.005 + 1.005^2 + \dots + 1.005^{n-1})$  ✓

(iii)  $A_{60} = 0$

So  $(M-15)[1 + 1.005 + 1.005^2 + \dots + 1.005^{n-1}] = 15000(1.005)^{60}$  ✓

GP  $a=1, r=1.005$

$\therefore (M-15)\left[\frac{1.005^{60} - 1}{1.005 - 1}\right] = 15000(1.005)^{60}$  ✓

$M = \frac{15000(1.005)^{60} \times 0.005}{1.005^{60} - 1} + 15$

$= \$304.99$  ✓

b) (i)  $x - 4y + 2 = 0$

$x = 4y^2$

$4y^2 - 4y + 2 = 0$  ✓

For tangent  $\Delta = 0$

So  $(-4)^2 - 4A(2) = 0$

$16 - 8A = 0$  ✓

$A = 2$

(ii) At point of intersection

$2y^2 - 4y + 2 = 0$


$y^2 - 2y + 1 = 0$

$(y-1)^2 = 0$

$y = 1$  ✓


and  $x = 2$

Coords. of point of intersection (2, 1)



# Question 9

a) (i)  $\frac{d}{dx} (x \ln x - x) = x \cdot \frac{1}{x} + \ln x - 1 = \ln x$  ✓




(ii) Area =  $3e^3 - \int_1^{e^3} \ln x \, dx$   
 $= 3e^3 - [x \ln x - x]_1^{e^3}$  ✓  
 $= 3e^3 - [(e^3 \ln e^3 - e^3) - (0 - 1)]$   
 $= 3e^3 - [3e^3 - e^3 + 1]$   
 $= e^3 - 1$  square units. ✓


Alternate approach:

Area =  $\int_0^3 x \, dy$   
 $= \int_0^3 e^y \, dy$   
 $= [e^y]_0^3$   
 $= e^3 - e^0$   
 $= e^3 - 1$  sq. units

b) (i)  $\angle CAB = \alpha$   
 $\angle ADG = \beta$



$\angle AGD = \alpha$  (alternate angles,  $AB \parallel DC$ ) ✓  
 $\angle AGC = \beta$  (corresponding angles,  $DF \parallel AE$ ) ✓  
 $\angle AGD = \angle CGE$  (vertically opposite) ✓  
 $\therefore \alpha = \beta$   
 $\therefore \triangle ADG$  is equilateral.



(ii) Area = Area ABCD +  $\triangle DGE$   
 $= 7\sqrt{3} + \frac{1}{2} 7^2 \sin 60^\circ$  ✓  
 $= 7\sqrt{3} + \frac{49 \cdot \sqrt{3}}{2}$   
 $= \frac{28\sqrt{3} + 49\sqrt{3}}{2}$   
 $= \frac{77\sqrt{3}}{2}$  sq. units ✓

Height of parallelogram ABCD =  $\sqrt{3}$

Alternate Method 1


Area of trapezium =  $\frac{1}{2} h (a+b)$   
 $= \frac{1}{2} \frac{7\sqrt{3}}{2} (2+9)$   
 $= \frac{77\sqrt{3}}{4}$  sq. units

$h = \frac{7\sqrt{3}}{2}$

Question 9 (cont.)

Alternate Method 2

Consider area of equilateral triangle side lengths 9 units.



$$\begin{aligned} \text{Area} &= \frac{1}{2} 9^2 \sin 60^\circ - \frac{1}{2} 2^2 \sin 60^\circ \\ &= \frac{81}{2} \frac{\sqrt{3}}{2} - \frac{4}{2} \frac{\sqrt{3}}{2} \\ &= \frac{77\sqrt{3}}{4} \text{ sq. units.} \end{aligned}$$

(c)

$$y = \frac{e^x + e^{-x}}{2}$$

$$y' = \frac{1}{2} (e^x - e^{-x})$$

$$y'' = \frac{1}{2} (e^x + e^{-x})$$

We need to show  $y'' = \sqrt{1 + (y')^2}$

$$\text{LHS} = \frac{1}{2} (e^x + e^{-x})$$

$$\text{RHS} = \sqrt{1 + \left(\frac{1}{2}(e^x - e^{-x})\right)^2}$$

$$= \sqrt{1 + \frac{1}{4}(e^{2x} - 2 + e^{-2x})}$$

$$= \frac{1}{2} \sqrt{4 + e^{2x} - 2 + e^{-2x}}$$

$$= \frac{1}{2} \sqrt{e^{2x} + 2 + e^{-2x}}$$

$$= \frac{1}{2} \sqrt{(e^x + e^{-x})^2}$$

$$= \frac{1}{2} (e^x + e^{-x})$$

$$= \text{LHS}$$

Question 10

(a) (i)  $\angle ADX = 180^\circ - 90^\circ - \theta$  (angle sum of  $\triangle ADX$ )  
 $= 90^\circ - \theta$

$\therefore \angle POC = 90^\circ - (90^\circ - \theta)$  (property of a square)  
 $= \theta$  as required.

(ii) In triangles  $ADQ$  and  $DCP$   
 $AD = DC$  (sides of a square)

$\angle PDC = \angle QAD = \theta$  (from (i) above)

$\angle ADQ = \angle DCP = 45^\circ$  (diagonals of a square bisect vertex)

$\therefore \triangle ADQ \cong \triangle DCP$  (AAS)

(iii) In triangles  $DXQ$  and  $AMQ$

$\angle AMQ = 90^\circ$  (given)

$\angle DXQ = 90^\circ$  (diagonals of a square intersect at right angles)

$\angle AQM = \angle DQX$  (vertically opposite angles)

$\therefore \triangle DXQ \parallel \triangle AMQ$  (AA)

b) (i) In  $\triangle HAB$ ,  $\cos \theta = \frac{1}{HB}$  and  $\tan \theta = \frac{AB}{1}$

$HB = \frac{1}{\cos \theta}$

$AB = \tan \theta$

Distance walked =  $2HB = \frac{2}{\cos \theta}$ ,  $\therefore$  Time walked =  $\frac{\frac{2}{\cos \theta}}{4}$  hours

Distance skated =  $6 - 2\tan \theta$ ,  $\therefore$  Time skated =  $\frac{6 - 2\tan \theta}{12}$  hours

So  $T = \frac{1}{2\cos \theta} + \frac{6 - 2\tan \theta}{12}$   
 $= \frac{1}{2\cos \theta} + \frac{1}{2} - \frac{\tan \theta}{6}$  as required.

(ii)  $T = \frac{(\cos \theta)^{-1}}{2} + \frac{1}{2} - \frac{\tan \theta}{6}$

$\frac{dT}{d\theta} = \frac{-(-\sin \theta)(\cos \theta)^{-2}}{2} - \frac{\sec^2 \theta}{6}$

$= \frac{\sin \theta}{2\cos^2 \theta} - \frac{1}{6\cos^2 \theta}$

$= \frac{1}{\cos^2 \theta} (3\sin \theta - 1)$

For minimum  $\frac{dT}{d\theta} = 0$

So

$3\sin \theta - 1 = 0$

$\sin \theta = \frac{1}{3}$

$\theta \doteq 19^\circ 28'$

and

$\theta$	$10^\circ$	$19^\circ 28'$	$30^\circ$
$\frac{dT}{d\theta}$	-0.94	0	$\frac{1}{9}$

$\therefore \theta = 19^\circ 28'$  is a minimum.

Test boundaries:  
 $\theta = 0$   $T = \frac{1}{4} + \frac{6}{12} + \frac{1}{4} = 1$  hour  
 $\theta = 90^\circ$   $T = \frac{6}{4} = 1\frac{1}{2}$  hours  
 and  $\theta = 19^\circ 28'$   $T = \frac{1}{2\cos 19^\circ 28'} + \frac{1}{2} - \frac{\tan 19^\circ 28'}{6} \doteq 58.28$  minutes.