



NORTH SYDNEY  
GIRLS HIGH SCHOOL

2009

HIGHER SCHOOL CERTIFICATE

ASSESSMENT TASK #3

# Mathematics Extension 1

Student Name: \_\_\_\_\_ Teacher: \_\_\_\_\_

*General Instructions*

- Reading time – 2 minutes.
- Working time – 65 minutes.
- Write using black or blue pen.
- Board approved calculators may be used.
- All *necessary* working should be shown in every question if full marks are to be awarded.
- Marks may **NOT** be awarded for untidy or badly arranged work.
- Start each **NEW** question on a separate answer sheet.

**Total Marks: 57 Marks**

- Attempt Questions 1 - 6
- All questions are **NOT** of equal value.

	Q 1		Q 2		Q 3		Q4			Q5	Q6			Total
	acd	b	abcd	e	a	b	a	b	c		a	b	c	
<b>H 6</b>														/3
<b>H 8</b>														/10
<b>H 9</b>														/31
<b>HE 2</b>														/3
<b>HE 6</b>														/10
	/9		/10		/10		/9			/10	/9			/57

**Total marks – 57**

**Attempt Questions 1 - 6**

**All questions are NOT of equal value**

Start each question on a SEPARATE answer sheet.

---

<b>Question 1</b>	(9 marks)	<b>Marks</b>
(a)	Evaluate $\int_0^1 \frac{dx}{2x+1}$ , leaving your answer in the exact form.	<b>2</b>
(b)	Using the substitution $u = 4 - x^2$ , evaluate $\int \frac{x}{\sqrt{4-x^2}} dx$	<b>3</b>
(c)	Let $f(x) = \frac{1}{2}(e^x + e^{-x})$ and $F(x) = \frac{1}{2}(e^x - e^{-x})$ Prove that $[f(x) + F(x)]^n = f(nx) + F(nx)$	<b>2</b>
(d)	Evaluate $\int_0^1 \frac{e^x}{e^x+1} dx$	<b>2</b>

---

**Question 2** (10 Marks) Start a NEW answer sheet.**Marks**

- (a) Solve  $e^x = 5$ , leaving your answer correct to 3 decimal places **1**
- (b) Find a primitive of  $\frac{3x}{1+x^2}$  **2**
- (c) Find  $\frac{d}{dx}(3x \log_e x)$  **2**
- (d) Evaluate  $\int_0^3 3^x dx$  **2**
- (e) Using the substitution  $u = \log_e x$ , evaluate  $\int_1^e \frac{(1 + \log_e x)^2}{x} dx$  **3**


(a) (i) Show that  $\frac{5}{\sqrt{5x+3}-\sqrt{5x-2}} = \sqrt{5x+3} + \sqrt{5x-2}$  **2**

(ii) Hence find  $\int \frac{dx}{\sqrt{5x+3}-\sqrt{5x-2}}$  **2**

(b) (i) Show that  $\frac{d}{dx}(x \ln x - x) = \ln x$ . **1**


(ii) Hence, or otherwise, find  $\int \ln x^2 dx$ . **2**

(iii) The graph below shows the curve  $y = \ln x^2$  ( $x > 0$ ) which meets the line  $x = 5$  at  $Q$ . **3**  
Using your answers above, or otherwise, find the area of the shaded region.




(a) Find  $\int \frac{x+1}{x^2} dx$  **2**

(b) The following graph shows the gradient function  $y = f'(x)$ . **3**  
 The graph shows that  $f'(1) = f'(-1) = 0$ .  
 Sketch the graph of  $y = f(x)$ , given that  $f'(x)$  is continuous everywhere except at  $x = 2$  and that  $f(0) = 1$  and  $f(-1) = -2$



(c) The shaded region below is that bounded by  $y = \frac{1}{\sqrt{x}}$ , the coordinate axes and the lines  $x = e$  and  $y = e$ . **4**  
 Find the volume when the shaded region is rotated about the y-axis, correct to 2 significant figures.



Consider the function  $y = \frac{\ln x}{x}$

- (a) What is the domain of this function? **1**
- (b) Show that  $\frac{d}{dx}\left(\frac{\ln x}{x}\right) = -\left(\frac{\ln x - 1}{x^2}\right)$  **1**
- (c) Describe the behaviour of the function as  $x$
- (i) approaches zero. **1**
- (ii) increases indefinitely **1**
- (d) Find any stationary points and determine their nature. **2**
- (e) Sketch the curve of this function. **2**
- (f) Hence find the value(s) of  $k$  for which  $e^{kx} = x$  has no solutions. **2**

---

**Question 6** (9 Marks) Start a NEW answer sheet. **Marks**

- (a) Use mathematical induction to show that the following statement is true **3**

$$n^3 + 2n \text{ is a multiple of } 12$$

where  $n$  is an even positive integer

- (b) By use of an appropriate diagram and reasons, evaluate the following sum. **2**  
**Do NOT evaluate any primitive functions.**

$$\int_0^1 e^x dx + \int_1^e \ln x dx$$

- (c) (i) Show  $\frac{1}{u} - \frac{1}{u+1} = \frac{1}{u(u+1)}$  **1**

- (ii) Using the substitution  $x = \ln u$ , find  $\int \frac{dx}{1+e^x}$  **3**

**End of paper**

**THIS PAGE IS INTENTIONALLY LEFT BLANK**

### STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln(x + \sqrt{x^2 - a^2}), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln(x + \sqrt{x^2 + a^2})$$

NOTE:  $\ln x = \log_e x, \quad x > 0$



NORTH SYDNEY  
GIRLS HIGH SCHOOL

2009

HIGHER SCHOOL CERTIFICATE

ASSESSMENT TASK #3

Mathematics      Extension 1

**SAMPLE SOLUTIONS**

---

**Question 1**

---

- (a) Evaluate  $\int_0^1 \frac{dx}{2x+1}$ , leaving your answer in the exact form.

$$\begin{aligned}\int_0^1 \frac{dx}{2x+1} &= \frac{1}{2} \int_0^1 \frac{2dx}{2x+1} = \frac{1}{2} [\ln|2x+1|]_0^1 \\ &= \frac{1}{2} (\ln 3 - \ln 1) \\ &= \frac{1}{2} \ln 3\end{aligned}$$

- (b) Using the substitution  $u = 4 - x^2$ , evaluate  $\int \frac{x}{\sqrt{4-x^2}} dx$

$$\begin{aligned}u = 4 - x^2 &\Rightarrow du = -2x dx \\ \int \frac{x}{\sqrt{4-x^2}} dx &= -\frac{1}{2} \int \frac{-2x}{\sqrt{4-x^2}} dx = -\frac{1}{2} \int \frac{du}{\sqrt{u}} = -\frac{1}{2} \int u^{-\frac{1}{2}} du \\ &= -u^{\frac{1}{2}} + c = -\sqrt{4-x^2} + c\end{aligned}$$

- (c) Let  $f(x) = \frac{1}{2}(e^x + e^{-x})$  and  $F(x) = \frac{1}{2}(e^x - e^{-x})$

$$\text{Prove that } [f(x) + F(x)]^n = f(nx) + F(nx)$$

$$\begin{aligned}f(x) + F(x) &= \frac{1}{2}(e^x + e^{-x}) + \frac{1}{2}(e^x - e^{-x}) \\ &= 2 \times \left(\frac{1}{2} e^x\right) = e^x\end{aligned}$$

$$\text{LHS} = [f(x) + F(x)]^n = (e^x)^n = e^{nx}$$

$$\begin{aligned}\text{RHS} = f(nx) + F(nx) &= \frac{1}{2}(e^{nx} + e^{-nx}) + \frac{1}{2}(e^{nx} - e^{-nx}) \\ &= 2 \times \frac{1}{2} e^{nx} = e^{nx}\end{aligned}$$

- (d) Evaluate  $\int_0^1 \frac{e^x}{e^x+1} dx$

$$\int_0^1 \frac{e^x}{e^x+1} dx = [\ln(e^x+1)]_0^1 = \ln(e+1) - \ln(2) = \ln\left(\frac{e+1}{2}\right)$$

---

**Question 2**

---

- (a) Solve  $e^x = 5$ , leaving your answer correct to 3 decimal places  
 $e^x = 5 \Rightarrow x = \ln 5 \approx 1.609437912\dots$   
 $x = 1.609$  [3 dp]

- (b) Find a primitive of  $\frac{3x}{1+x^2}$   
$$\int \frac{3x}{1+x^2} dx = \frac{3}{2} \int \frac{2x}{1+x^2} dx = \frac{3}{2} \ln(1+x^2)$$

- (c) Find  $\frac{d}{dx}(3x \log_e x)$   
$$\frac{d}{dx}(3x \log_e x) = 3x \times \frac{1}{x} + 3 \times \ln x$$
$$= 3 + 3 \ln x$$

- (d) Evaluate  $\int_0^3 3^x dx$   
$$\int_0^3 3^x dx = \left[ \frac{3^x}{\ln 3} \right]_0^3 = \frac{1}{\ln 3} (3^3 - 3^0) = \frac{26}{\ln 3}$$

- (e) Using the substitution  $u = \log_e x$ , evaluate  $\int_1^e \frac{(1 + \log_e x)^2}{x} dx$
- |   |  |
|---|--|
| $x = 1 \Rightarrow u = \ln 1 = 0$         | $\int_1^e \frac{(1 + \log_e x)^2}{x} dx = \int_1^e (1 + \log_e x)^2 \frac{dx}{x}$ $= \int_0^1 (1 + u)^2 du$ $= \left[ \frac{1}{3} (1 + u)^3 \right]_0^1$ $= \frac{1}{3} (2^3 - 1^3) = \frac{7}{3}$ |
| $x = e \Rightarrow u = \ln e = 1$         |  |
| $u = \ln x \Rightarrow du = \frac{dx}{x}$ |  |

---

**Question 3**

---

(a) (i) Show that  $\frac{5}{\sqrt{5x+3}-\sqrt{5x-2}} = \sqrt{5x+3} + \sqrt{5x-2}$

$$\begin{aligned}\text{LHS} &= \frac{5}{\sqrt{5x+3}-\sqrt{5x-2}} \\ &= \frac{5}{\sqrt{5x+3}-\sqrt{5x-2}} \times \frac{\sqrt{5x+3}+\sqrt{5x-2}}{\sqrt{5x+3}+\sqrt{5x-2}} \\ &= \frac{5(\sqrt{5x+3}+\sqrt{5x-2})}{[(5x+3)-(5x-2)]} \\ &= \frac{5(\sqrt{5x+3}+\sqrt{5x-2})}{5} \\ &= \sqrt{5x+3} + \sqrt{5x-2} \\ &= \text{RHS}\end{aligned}$$

(ii) Hence find  $\int \frac{dx}{\sqrt{5x+3}-\sqrt{5x-2}}$

$$\begin{aligned}\int \frac{dx}{\sqrt{5x+3}-\sqrt{5x-2}} &= \int \frac{(\sqrt{5x+3}+\sqrt{5x-2})dx}{5} \\ &= \frac{1}{5} \int \left[ (5x+3)^{\frac{1}{2}} + (5x-2)^{\frac{1}{2}} \right] dx \\ &= \frac{1}{5} \left[ \frac{1}{5} \times \frac{2}{3} (5x+3)^{\frac{3}{2}} + \frac{1}{5} \times \frac{2}{3} (5x-2)^{\frac{3}{2}} \right] + C \\ &= \frac{2}{75} \left[ (5x+3)^{\frac{3}{2}} + (5x-2)^{\frac{3}{2}} \right] + C\end{aligned}$$

---

**Question 3 continued**

---

- (b) (i) Show that  $\frac{d}{dx}(x \ln x - x) = \ln x$ .


$$\begin{aligned}\frac{d}{dx}(x \ln x - x) &= x \times \frac{1}{x} + 1 \times \ln x - 1 \\ &= 1 + \ln x - 1 \\ &= \ln x\end{aligned}$$

- (ii) Hence, or otherwise, find  $\int \ln x^2 dx$ .

$$\int \ln x^2 dx = 2 \int \ln x dx = 2(x \ln x - x) + C$$

- (iii) The graph below shows the curve  $y = \ln x^2$  ( $x > 0$ ) which meets the line  $x = 5$  at  $Q$ .

Using your answers above, or otherwise, find the area of the shaded region.



$P$  has coordinates  $(0, \ln 25)$

$$\begin{aligned}\text{The required area} &= \text{area rectangle } OPQR - \int_1^5 \ln x^2 dx \\ &= 5 \times \ln 25 - \left[ 2(x \ln x - x) \right]_1^5 \\ &= 5 \ln 25 - 2 \left[ (5 \ln 5 - 5) - (\ln 1 - 1) \right] \\ &= 5 \ln 25 - 10 \ln 5 + 10 - 2 \\ &= 5 \ln 25 - 5 \ln 25 + 8 \\ &= 8 \text{ u}^2\end{aligned}$$

---

**Question 4**

---

- (a) Find  $\int \frac{x+1}{x^2} dx$  2


$$\begin{aligned}\int \frac{x+1}{x^2} dx &= \int \left( \frac{1}{x} + \frac{1}{x^2} \right) dx = \int \left( \frac{1}{x} + x^{-2} \right) dx \\ &= \ln|x| - x^{-1} + C \\ &= \ln|x| - \frac{1}{x} + C\end{aligned}$$

- (b) The following graph shows the gradient function  $y = f'(x)$ . 3

The graph shows that  $f'(1) = f'(-1) = 0$ .


Sketch the graph of  $y = f(x)$ , given that  $f'(x)$  is continuous everywhere except at  $x = 2$  and that  $f(0) = 1$  and  $f(-1) = -2$

A possible solution:



**Question 4 continued**

- (c) The shaded region below is that bounded by  $y = \frac{1}{\sqrt{x}}$ , the coordinate axes and the lines  $x = e$  and  $y = e$ . 4  
 Find the volume when the shaded region is rotated about the y-axis, correct to 2 significant figures.



The volume  $V$  is the sum of two volumes  $V_1$  and  $V_2$ .

$V_1$  is the volume formed by rotating the curve  $y = \frac{1}{\sqrt{x}}$  from  $y = \frac{1}{\sqrt{e}}$  to  $y = e$

about the y-axis.  $y = \frac{1}{\sqrt{x}} \Rightarrow x = \frac{1}{y^2} \Rightarrow x^2 = \frac{1}{y^4} = y^{-4}$

$V_2$  is the cylinder formed by rotating the line  $x = e$  about the y-axis.

It has radius  $e$  and height  $\frac{1}{\sqrt{e}}$ .

$$\begin{aligned} V_1 &= \pi \int_{\frac{1}{\sqrt{e}}}^e x^2 dy = \pi \int_{\frac{1}{\sqrt{e}}}^e y^{-4} dy \\ &= \pi \left[ -\frac{1}{3} y^{-3} \right]_{\frac{1}{\sqrt{e}}}^e = \frac{\pi}{3} \left[ -\frac{1}{y^3} \right]_{\frac{1}{\sqrt{e}}}^e \\ &\quad \left[ \text{NB } (\sqrt{e})^3 = e\sqrt{e} \right] \\ &= \frac{\pi}{3} \left[ -\frac{1}{e^3} + \frac{1}{\frac{1}{e\sqrt{e}}} \right] = \frac{\pi}{3} \left[ \frac{\sqrt{e}}{e^2} - \frac{1}{e^3} \right] \\ &= \frac{\pi}{3} \left( e\sqrt{e} - \frac{1}{e^3} \right) \end{aligned}$$

$$\begin{aligned} V_2 &= \pi \times e^2 \times \frac{1}{\sqrt{e}} \\ &= \pi e^{\frac{3}{2}} \end{aligned}$$

$$V = \frac{\pi}{3} \left( e\sqrt{e} - \frac{1}{e^3} \right) + \pi e^{\frac{3}{2}} \approx 19 \text{ u}^3$$

**Question 5**

Consider the function  $y = \frac{\ln x}{x}$

(a) What is the domain of this function?  $x > 0$

(b) Show that  $\frac{d}{dx}\left(\frac{\ln x}{x}\right) = -\left(\frac{\ln x - 1}{x^2}\right)$

$$\frac{d}{dx}\left(\frac{\ln x}{x}\right) = \frac{x \times \frac{1}{x} - \ln x \times 1}{x^2} = \frac{1 - \ln x}{x^2} = -\left(\frac{\ln x - 1}{x^2}\right)$$

(c) Describe the behaviour of the function as  $x$

(i) approaches zero.  $y \rightarrow -\infty$

(ii) increases indefinitely  $y \rightarrow 0$

(d) Find any stationary points and determine their nature.

$$y' = 0 \Rightarrow \ln x - 1 = 0 \Rightarrow \ln x = 1$$


$\therefore x = e \Rightarrow \left(e, \frac{1}{e}\right)$  is the stationary point

$x$	2	$e$	3
$y'$	0.3	0	-0.1

Only need to check  $(1 - \ln x)$  as  $x^2 > 0$ .

So  $(e, e^{-1})$  is a maximum turning point.

(e) Sketch the curve of this function.



(f) Hence find the value(s) of  $k$  for which  $e^{kx} = x$  has no solutions.

$$e^{kx} = x \Rightarrow kx = \ln x$$

$$\therefore k = \frac{\ln x}{x}$$

So the solutions to  $e^{kx} = x$  are found by intersecting the line  $y = k$  with

$$y = \frac{\ln x}{x}$$

So there will be no solutions when  $k > \frac{1}{e}$ .

---

**Question 6**

---

- (a) Use mathematical induction to show that the following statement is true

$$n^3 + 2n \text{ is a multiple of } 12$$

where  $n$  is an even positive integer

Test  $n = 2$

$$2^3 + 2 \times 2 = 12$$

Clearly  $n = 2$  is true.

Assume true for  $n = 2k$  i.e.  $(2k)^3 + 2(2k) = 12N$ ,  $N \in \mathbb{Z}$

$$\therefore 8k^3 + 4k = 12N.$$

NTP true for  $n = 2k + 2$  i.e.  $(2k + 2)^3 + 2(2k + 2) = 12M$ ,  $M \in \mathbb{Z}$


$$\begin{aligned} (2k + 2)^3 + 2(2k + 2) &= 8(k + 1)^3 + 4(k + 1) \\ &= 8(k^3 + 3k^2 + 3k + 1) + 4k + 4 \\ &= (8k^3 + 4k) + 24k^2 + 24k + 12 \\ &= 12(N + 2k^2 + 2k + 1) \\ &= 12M \quad \left[ \because N + 2k^2 + 2k + 1 \in \mathbb{Z} \right] \end{aligned}$$

So the statement is true for  $n = 2k + 2$  provided it is true for  $n = 2k$ .

So by the principle of mathematical induction it is true for all positive even integers.

- (b) By use of an appropriate diagram and reasons, evaluate the following sum.  
**Do NOT evaluate any primitive functions.**

$$\int_0^1 e^x dx + \int_1^e \ln x dx$$



By symmetry the integral  $\int_1^e \ln x dx$  produces the same area as that of  $e^x$  next to the  $y$ -axis for  $1 \leq y \leq e$ .

So  $\int_0^1 e^x dx + \int_1^e \ln x dx$  is the area of the rectangle with dimensions  $1 \times e$

$$\therefore \int_0^1 e^x dx + \int_1^e \ln x dx = e$$

---

**Question 6 continued**

---

(c) (i) Show  $\frac{1}{u} - \frac{1}{u+1} = \frac{1}{u(u+1)}$

$$\frac{1}{u} - \frac{1}{u+1} = \frac{u+1-u}{u(u+1)} = \frac{1}{u(u+1)}$$

(ii) Using the substitution  $x = \ln u$ , find  $\int \frac{dx}{1+e^x}$

$$\begin{array}{l} x = \ln u \Rightarrow dx = \frac{du}{u} \\ x = \ln u \Rightarrow u = e^x \end{array} \left| \begin{array}{l} \int \frac{dx}{1+e^x} = \int \frac{1}{1+e^x} \times dx = \int \left( \frac{1}{1+u} \right) \frac{du}{u} \\ = \int \left[ \frac{du}{u(u+1)} \right] = \int \left( \frac{1}{u} - \frac{1}{u+1} \right) du \\ = \ln u - \ln(u+1) + C \\ = \ln \left( \frac{e^x}{1+e^x} \right) + C \quad \left[ = x - \ln(1+e^x) + C \right] \end{array} \right.$$

**End of Solutions**