

**QUESTION 1.**Use a *separate* Writing Booklet**Marks**

- (a) Find a primitive of  $\frac{1}{\sqrt{1-x^2}}$ . 1
- (b) Evaluate  $\int_0^{\ln 5} e^{-x} dx$  2
- (c) Find  $\lim_{h \rightarrow 0} \frac{\sin \frac{h}{2}}{h}$  1
- (d) A box contains 5 red marbles and 4 white marbles.  
Three marbles are drawn in succession without replacement,  
What is the probability that any two marbles are red and one is white? 2
- (e) The sector  $OAB$  of a circle centre  $O$  and radius  $r$  has an area of  $\frac{3\pi}{4} \text{ cm}^2$ . 2  
If the arc  $AB$  subtends an angle of  $\frac{\pi}{6}$  at  $O$ , find the length of the arc  $AB$ .
- (f) If  $\cos^{-1}\left(\frac{\sqrt{3}}{2}\right) + \tan^{-1}(1) = k\pi$  (where  $k$  is rational) 2  
Find the value of  $k$ .
- (g) Find  $\int \frac{\ln 3x}{x} dx$ , using the substitution  $u = \ln 3x$ . 4

**QUESTION 2.**Use a *separate* Writing Booklet.**Marks**(a) Differentiate the following with respect to  $x$ .**8**

(i)  $x \cos x$

(ii)  $\log_e(\cos 5x)$

(iii)  $\tan^{-1}\left(\frac{x}{3}\right)$

(iv)  $e^{\tan x}$

(v)  $\left\{1 + \cos^{-1}(3x)\right\}^3$

(b) Find a primitive of:

**6**

(i)  $x^2 e^{2x^3+1}$

(ii)  $\frac{5}{4+x^2}$

(iii)  $\frac{x^2}{x^3+7}$

(iv)  $-\sin(\pi - x)$

**QUESTION 3.**Use a *separate* Writing Booklet.**Marks**

- (a) In choosing three letters from the word PROBING, and assuming each choice is equally likely, what is the probability of choosing just one vowel? 2
- (b) (i) In how many ways can the numbers 1, 2, 3, 4, 5, 6 be arranged around a circle? 3
- (ii) How many of these arrangements have at least two even numbers together?
- (c) (i) Show that the function  $f(x) = x^2 + e^{-\frac{1}{2}x} - 5$  has a root between  $-2$  and  $-1$ . 4
- (ii) Taking  $x = -2$  as a first approximation, apply Newton's method once, to show that the root of  $f(x) = 0$  is approximately  $-\frac{18}{e+8}$ .
- (d) (i) Show that  $\frac{1+x}{1-x} = -1 + \frac{2}{1-x}$ . 3
- (ii) Hence find  $\int \frac{1+x}{1-x} dx$
- (e) Write down the general solution of  $\sqrt{3} \tan \theta - 1 = 0$ . 2  
Leave answer in exact radian form.

**QUESTION 4.**Use a *separate* Writing Booklet**Marks**

- (a) Consider the curves  $y = \sin x$  and  $y = \cos 2x$ . **5**
- (i) Sketch the graphs of the two curves on the same axes, in the domain  $\frac{-\pi}{2} \leq x \leq \frac{\pi}{6}$ .
- (ii) Show that the curves intersect at  $x = -\frac{\pi}{2}$  and  $x = \frac{\pi}{6}$ .
- (iii) Hence find the exact area bounded by the two curves.
- (b) Consider the function  $f(x) = 3\sin^{-1}\left(\frac{x}{2}\right)$ . **3**
- (i) Evaluate  $f(2)$ .
- (ii) State the domain and range of  $y = f(x)$ .
- (iii) Sketch the graph of  $y = f(x)$ .
- (c) Find the exact value of  $\tan\left[2\tan^{-1}\left(-\frac{1}{2}\right)\right]$ . **3**
- (d) The function  $g(x)$  is given by  $g(x) = \cos^{-1}x + \sin^{-1}x$ ,  $0 \leq x \leq 1$ . **3**
- (i) Find  $g'(x)$ .
- (ii) Sketch the graph of  $y = g(x)$ .

**QUESTION 5.**Use a *separate* Writing Booklet**Marks**

(a) Consider the function  $f(x) = x \sin^{-1}(x^2)$ . **8**

- (i) State the domain and range of  $f(x)$ .
- (ii) Find  $f'(x)$ .
- (iii) Show that there is a horizontal inflexion at the origin.
- (iv) What is the slope of the tangent at  $x = 1$ .
- (v) Sketch the curve which represents  $f(x)$ .

(b) (i) Show that  $\int \frac{x}{\sqrt{x^2 + 2x - 3}} dx = \int \frac{x}{\sqrt{(x+1)^2 - 4}} dx$ . **3**

(ii) Using the substitution  $2 \sec u = x + 1$ , find  $\int \frac{x}{\sqrt{x^2 + 2x - 3}} dx$

(c) The curve  $y = \frac{1}{\sqrt{1+x^2}}$  is rotated about the  $x$ -axis. **3**

Find the volume of the solid enclosed between  $x = \frac{1}{\sqrt{3}}$  and  $x = \sqrt{3}$ .

**QUESTION 6.**Use a *separate* Writing Booklet**Marks**

(a) Evaluate  $\int_0^{\sqrt{2}} x \sqrt[3]{x^2 + 1} dx$  – using the substitution  $u = x^2 + 1$ . **3**  
Answer in exact form.

(b) Consider the equation  $\cos 3x = \sin x$  **3**

(i) Show that  $\cos\left(\frac{\pi}{2} - A\right) = \sin A$

(ii) Hence, or otherwise, find the general solution of the equation  $\cos 3x = \sin x$ .

(c) (i) Show that  $\frac{d}{dx} \tan^3 x = 3\sec^4 x - 3\sec^2 x$ . **4**

(ii) Using (i) or otherwise, evaluate  $\int_0^{\frac{\pi}{4}} \sec^4 x dx$  –

(d) Show that the exact value of  $\int_0^{\frac{\pi}{6}} \sin^2 x dx = \frac{2\pi - 3\sqrt{3}}{24}$  **4**

**THIS IS THE END OF THE PAPER**





(1) (a)  $\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1}x + c$

(b)  $\int_0^{\ln 5} e^{-x} dx = -e^{-x} \Big|_0^{\ln 5}$   
 $= -[e^{-\ln 5} - e^0]$   
 $= -[e^{\ln \frac{1}{5}} - 1]$   
 $= 1 - \frac{1}{5} = \frac{4}{5}$

$e^{\ln x} = x$

(c)  $\lim_{h \rightarrow 0} \frac{\sin \frac{h}{2}}{\frac{h}{2}} = \frac{1}{2} \lim_{h \rightarrow 0} \frac{\sin \frac{h}{2}}{\frac{h}{2}}$   
 $= \frac{1}{2} \times 1$   
 $= \frac{1}{2}$

$\left[ \lim_{u \rightarrow 0} \frac{\sin u}{u} = 1 \right]$

(d) 5R 4W

method 1:

$\frac{\binom{5}{2} \times \binom{4}{1}}{\binom{9}{3}} = \frac{10}{21}$


method 2: RRW

$\left( \frac{5}{9} \times \frac{4}{8} \times \frac{4}{7} \right)$

x3

$= \frac{10}{21}$

(e)



$\therefore \frac{1}{2} r^2 \theta = \frac{3\pi}{4}$

$\therefore \frac{1}{2} r^2 \times \frac{\pi}{6} = \frac{3\pi}{4}$

$\therefore r^2 = \frac{3}{4} \times 12 = 9$

$\therefore r = 3$

$\therefore l = r\theta$

$= 3 \times \frac{\pi}{6}$

$= \frac{\pi}{2} \text{ cm}$

- 2 -

$$(f) \quad \cos^{-1}\left(\frac{\sqrt{3}}{2}\right) + \tan^{-1}(1) = \frac{\pi}{6} + \frac{\pi}{4} = \frac{10\pi}{24} = \frac{5\pi}{12}$$

$$\boxed{R = \frac{5}{12}}$$

$$(g) \quad \int \frac{\ln 3x}{x} dx \quad u = \ln 3x$$
$$\therefore du = \frac{3}{3x} dx = \frac{dx}{x}$$

$$= \int \ln 3x \times \frac{dx}{x}$$

$$= \int u du = \frac{1}{2} u^2 + C$$

$$= \frac{1}{2} \ln^2 3x + C$$

$$(2) \quad (a) \quad (i) \quad \frac{d(x \cos x)}{dx} = x(-\sin x) + (1) \cos x$$
$$= \cos x - x \sin x$$

$$(ii) \quad \frac{d(\ln(\cos 5x))}{dx} = \frac{1}{\cos 5x} \times -5 \sin 5x$$
$$= -\frac{5 \sin 5x}{\cos 5x} = -5 \tan 5x$$

$$(iii) \quad \frac{d\left(\tan^{-1}\left(\frac{x}{3}\right)\right)}{dx} = \frac{1}{3} \times \frac{1}{1 + \left(\frac{x}{3}\right)^2} = \frac{1}{3} \times \frac{9}{9 + x^2}$$
$$= \frac{3}{9 + x^2}$$

$$(iv) \quad \frac{d(e^{\tan x})}{dx} = \sec^2 x e^{\tan x}$$

$$(v) \quad \frac{d\left(\left[1 + \cos^{-1}(3x)\right]^3\right)}{dx} = 3\left[1 + \cos^{-1}(3x)\right]^2 \times \frac{-3}{\sqrt{1-9x^2}} = -\frac{9\left(1 + \cos^{-1}(3x)\right)^2}{\sqrt{1-9x^2}}$$

2(b) (i)  $\int x^2 e^{2x^3+1} dx$

$$= \frac{1}{6} \int (6x^2) e^{2x^3+1} dx$$

$$\left[ \text{N.B. } \frac{d(2x^3+1)}{dx} = 6x^2 \right]$$

$$= \frac{1}{6} e^{2x^3+1} + C$$

(ii)  $\int \frac{5}{4+x^2} dx = \frac{5}{2} \int \frac{2}{4+x^2} dx$

$$= \frac{5}{2} \tan^{-1}\left(\frac{x}{2}\right) + C$$

(iii)  $\int \frac{x^2}{x^3+7} dx = \frac{1}{3} \int \frac{3x^2}{x^3+7} dx = \frac{1}{3} \ln|x^3+7| + C$

(iv)  $-\sin(\pi-x) = -\sin x$

$$\therefore \int -\sin x dx = \cos x + C$$

(3) (a) PRBN IG

$$3 \text{ letters} = \binom{6}{3} = 20$$

$$\frac{12}{20} = \frac{3}{5}$$

$$1 \text{ vowel} = \binom{4}{2} \times \binom{2}{1} = 6 \times 2$$

(b) (i) 6 numbers  $\Rightarrow 5! = 120$

(ii) 135 246

No even numbers together mean alternating



place an even number first

$$\therefore 2! \times 3! = 2 \times 3 = 12 \text{ ways}$$

$$\therefore \text{Prob} = \frac{12}{120} = \frac{1}{10}$$

$$\therefore \text{At least 2} = 1 - \frac{1}{10} = \frac{9}{10}$$

-4-

(c) (i)  $f(x) = x^2 + e^{-\frac{1}{2}x} - 5$  is continuous

$$f(-2) = 4 + e^{-1} - 5 = e^{-1} - 1 > 0 \quad (\because e > 3)$$

$$f(-1) = 1 + e^{-\frac{1}{2}} - 5 = e^{-\frac{1}{2}} - 4 < 0$$

$\therefore f(-2) \cdot f(-1) < 0$  and with  $f$  continuous

$\exists c$  s.t.  $f(c) = 0$ ,  $-2 < c < -1$

(ii)  $x_0 = -2$

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$\begin{cases} f(-2) = e^{-1} \\ f'(x) = 2x - \frac{1}{2}e^{-\frac{1}{2}x} \\ f'(-2) = -4 - \frac{1}{2}e \\ = \frac{-8 - e}{2} = -\frac{e+8}{2} \end{cases}$$

$$\therefore x_1 = -2 - \frac{e^{-1}}{-\left(\frac{e+8}{2}\right)}$$

$$= -2 + \frac{2(e^{-1})}{e+8} = \frac{-2(e+8) + 2(e^{-1})}{e+8}$$

$$= \frac{-2e - 16 + 2e^{-1}}{e+8}$$

$$= -\frac{16}{e+8}$$

$$(d) (i) \frac{1+x}{1-x} = \frac{-(1-x) + 2}{1-x}$$

$$= -\frac{(1-x)}{1-x} + \frac{2}{1-x}$$

$$= -1 + \frac{2}{1-x}$$

$$(ii) \int \frac{1+x}{1-x} dx = \int \left(-1 + \frac{2}{1-x}\right) dx$$


$$= -x + -2 \ln|1-x| + c$$

$$= -x - 2 \ln|1-x| + c$$

3(e)  $\sqrt{3} \tan \theta = 1$   
 $\tan \theta = \frac{1}{\sqrt{3}}$

$\therefore \theta = n\pi + \frac{\pi}{6}$

(4) (a) (i)



(ii)  $\sin\left(-\frac{\pi}{2}\right) = -1$

$\cos\left(2 \times -\frac{\pi}{2}\right) = \cos(-\pi) = -1$

$\sin\frac{\pi}{6} = \frac{1}{2}$

$\cos\left(2 \times \frac{\pi}{6}\right) = \cos\frac{\pi}{3} = \frac{1}{2}$

(iii) 
$$\text{Area} = \int_{-\pi/2}^{\pi/6} (\cos 2x - \sin x) dx = \left[ \frac{1}{2} \sin 2x + \cos x \right]_{-\pi/2}^{\pi/6}$$

$$= \left( \frac{1}{2} \sin \frac{\pi}{3} + \cos \frac{\pi}{6} \right) - (0)$$

$$= \frac{1}{2} \times \frac{\sqrt{3}}{2} + \frac{\sqrt{3}}{2}$$


$$= \frac{3}{2} \times \frac{\sqrt{3}}{2} = \frac{3\sqrt{3}}{4}$$

(b)  $f(x) = 3 \sin^{-1}\left(\frac{x}{2}\right)$

(i)  $f(2) = 3 \sin^{-1}(1) = 3 \times \frac{\pi}{2} = \frac{3\pi}{2}$

(ii)  $-1 \leq \frac{x}{2} \leq 1 \Rightarrow -2 \leq x \leq 2$

$-\frac{\pi}{2} \leq \frac{y}{3} \leq \frac{\pi}{2} \Rightarrow -\frac{3\pi}{2} \leq y \leq \frac{3\pi}{2}$



4(c)  $\tan \left[ 2 \tan^{-1} \left( -\frac{1}{2} \right) \right]$

$\left\| \begin{array}{l} -\frac{\pi}{2} < \tan^{-1} x < \frac{\pi}{2} \end{array} \right.$

let  $\alpha = \tan^{-1} \left( -\frac{1}{2} \right)$

$\therefore \tan \alpha = -\frac{1}{2}$


$\tan 2\alpha = \frac{2 \tan \alpha}{1 - \tan^2 \alpha} = \frac{2 \times \left( -\frac{1}{2} \right)}{1 - \left( -\frac{1}{2} \right)^2} = \frac{-1}{1 - \frac{1}{4}} = \frac{-1}{\frac{3}{4}} = -\frac{4}{3}$

(d)  $g(x) = \cos^{-1} x + \sin^{-1} x$

(i)  $g'(x) = \frac{-1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{1-x^2}} = 0$

(ii)  $\therefore g(x) = \text{constant for } -1 \leq x \leq 1$

$g(0) = \cos^{-1}(0) + \sin^{-1}(0) = \frac{\pi}{2} + 0 = \frac{\pi}{2}$



(5) (a)  $f(x) = x \sin^{-1}(x^2)$

(i)  $-1 \leq x^2 \leq 1 \Rightarrow -1 \leq x \leq 1$

(ii)  $f'(x) = \sin^{-1}(x^2) + x \left( \frac{2x}{\sqrt{1-x^4}} \right)$

$\Rightarrow -\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$

$= \sin^{-1}(x^2) + \frac{2x^2}{\sqrt{1-x^4}}$

= even function

(iii)

x	0 <sup>-</sup>	0	0 <sup>+</sup>
f'(x)	+	0	+

$\leftarrow$   $\therefore f' \left( -\frac{1}{2} \right) = f' \left( \frac{1}{2} \right)$

$\therefore$  HPOI

$\frac{+}{0} / +$


(iv)  $f'(1) = \sin^{-1} 1 + \frac{2}{\sqrt{0}}$

$\therefore$  vertical tangent

[OR:  $\sin^{-1}(x^2)$  has a double root at  $x=0$   $\therefore$  even

$\therefore x \sin^{-1}(x^2)$  has a triple root  $\therefore$  HPOI ]

5 (a) (v)  $f(x)$  is an odd function




(b) (i)  $x^2 + 2x - 3 = (x^2 + 2x + 1) - 4$   
 $= (x+1)^2 - 4$

$$\therefore \int \frac{x}{\sqrt{x^2 + 2x - 3}} dx = \int \frac{x dx}{\sqrt{(x+1)^2 - 4}}$$

(ii)  $2 \sec u = x+1$   
 $\therefore 2 \sec u \tan u du = dx$

$$\tan^2 u + 1 = \sec^2 u$$

$$[4 \sec^2 u - 4 = 4(\sec^2 u - 1) = 4 \tan^2 u]$$



$$\therefore \tan u = \frac{\sqrt{x^2 + 2x - 3}}{2}$$

$$\begin{aligned} & \int \frac{x dx}{\sqrt{(x+1)^2 - 4}} \\ &= \int \frac{(2 \sec u - 1) 2 \sec u \tan u du}{\sqrt{4 \sec^2 u - 4}} \\ &= \int \frac{2(2 \sec u - 1) \sec u \tan u du}{2 \tan u} \\ &= \int (2 \sec^2 u - \sec u) du \\ &= 2 \tan u - \int \sec u du \\ &= \sqrt{x^2 + 2x - 3} - \int \sec u du \end{aligned}$$

could be done reversing the substitution BUT worth more than 2 marks!

not an Ext 1 integral

$$\begin{aligned} &= \sqrt{x^2 + 2x - 3} - \ln |\sec u + \tan u| + C \\ &= \sqrt{x^2 + 2x - 3} - \ln \left| \frac{x+1}{2} + \frac{\sqrt{x^2 + 2x - 3}}{2} \right| + C \\ &= \sqrt{x^2 + 2x - 3} - \ln |x+1 + \sqrt{x^2 + 2x - 3}| + K \end{aligned}$$

$$(5) \quad (c) \quad y = \frac{1}{\sqrt{1+x^2}}$$

$$\therefore V = \pi \int_{\frac{1}{\sqrt{3}}}^{\sqrt{3}} y^2 dx$$

$$= \pi \int_{\frac{1}{\sqrt{3}}}^{\sqrt{3}} \frac{1}{1+x^2} dx$$

$$= \pi \left[ \tan^{-1}(x) \right]_{\frac{1}{\sqrt{3}}}^{\sqrt{3}}$$

$$= \pi \left[ \tan^{-1}(\sqrt{3}) - \tan^{-1}\left(\frac{1}{\sqrt{3}}\right) \right]$$

$$= \pi \left[ \frac{\pi}{3} - \frac{\pi}{6} \right]$$

$$= \pi \left[ \frac{\pi}{6} \right]$$

$$= \frac{\pi^2}{6} \text{ c.u.}$$

$$(6) \quad a) \quad \int_0^{\sqrt{2}} x^3 \sqrt{x^2+1} dx \quad \equiv \frac{1}{2} \int_0^{\sqrt{2}} \sqrt{x^2+1} (2x dx)$$

$$u = x^2 + 1 \quad \therefore du = 2x dx$$

$$x=0 \Rightarrow u=1$$

$$x=\sqrt{2} \Rightarrow u=3$$

$$= \frac{1}{2} \int_1^3 u^{\frac{1}{2}} du$$

$$= \frac{1}{2} \times \frac{3}{4} u^{\frac{4}{3}} \Big|_1^3$$

$$= \frac{3}{8} [3^{\frac{4}{3}} - 1]$$

$$= \frac{3}{8} [\sqrt[3]{81} - 1]$$

$$= \frac{3}{8} [3^3 \sqrt[3]{3} - 1]$$

$$27 \times 3 = 81$$

6(b)  $\cos 3x = \sin x$

(i)  $\cos\left(\frac{\pi}{2} - A\right)$   
 $= \cos\frac{\pi}{2} \cos A + \sin\frac{\pi}{2} \sin A$   
 $= 0 \times \sin A + 1 \times \sin A$   
 $= \sin A$

(ii)  $\cos 3x = \sin x$   
 $\Rightarrow \cos 3x = \cos\left(\frac{\pi}{2} - x\right)$

$\therefore 3x = 2n\pi \pm \left(\frac{\pi}{2} - x\right)$

$$\begin{array}{l} 3x = 2n\pi + \frac{\pi}{2} - x \\ 4x = (4n+1)\pi \\ x = \left(\frac{4n+1}{4}\right)\pi \end{array} \quad \left| \begin{array}{l} 3x = 2n\pi - \frac{\pi}{2} + x \\ 2x = \frac{(4n-1)\pi}{2} \\ x = \frac{(4n-1)\pi}{4} \end{array} \right.$$

$\therefore x = \left(\frac{4n \pm 1}{4}\right)\pi$

(c) (i)  $d\left(\frac{\tan^3 x}{dx}\right) = 3 \tan^2 x \cdot \sec^2 x$   
 $= 3(\sec^2 x - 1) \sec^2 x$   
 $= 3 \sec^4 x - 3 \sec^2 x$

(ii)  $\therefore \tan^3 x = \int (3 \sec^4 x - 3 \sec^2 x) dx$

$\therefore \tan^3 x \Big|_0^{\pi/4} = 3 \int_0^{\pi/4} \sec^4 x dx - 3 \int_0^{\pi/4} \sec^2 x dx$

$\therefore 3 \int_0^{\pi/4} \sec^4 x dx = \left[ \tan^3 \frac{\pi}{4} - \tan^3(0) \right] + 3 \tan x \Big|_0^{\pi/4}$

$= (1-0) + 3(1-0)$

$= 4$

$\therefore \int_0^{\pi/4} \sec^4 x dx = \frac{4}{3}$

$$(6)(d) \int_0^{\pi/6} \sin^2 x \, dx$$

$$= \frac{1}{2} \int_0^{\pi/6} 2 \sin^2 x \, dx$$

$$= \frac{1}{2} \int_0^{\pi/6} (1 - \cos 2x) \, dx$$

$$= \frac{1}{2} \left[ x - \frac{1}{2} \sin 2x \right]_0^{\pi/6}$$

$$= \frac{1}{2} \left[ \frac{\pi}{6} - \frac{1}{2} \sin \frac{\pi}{3} \right]$$

$$= \frac{\pi}{12} - \frac{1}{4} \times \frac{\sqrt{3}}{2}$$

$$= \frac{\pi}{12} - \frac{\sqrt{3}}{8}$$

$$= \frac{2\pi - 3\sqrt{3}}{24}$$

$$\cos 2A = 1 - 2\sin^2 A$$