



BARKER COLLEGE

**TRIAL HIGHER SCHOOL CERTIFICATE
2000**

**MATHEMATICS
3 UNIT (ADDITIONAL)
AND
3/4 UNIT (COMMON)**

BTP
AES
CFR
PJR
MRB
JGD*
JFH*

PM TUESDAY 1 AUGUST
LONDONDERRILLI
68 QUEENSTOWN ST
KILDARE 2061
(15)

100 copies

*TIME ALLOWED : TWO HOURS
(Plus 5 minutes reading time)*

DIRECTIONS TO STUDENTS:

- Write your Barker Student Number on EACH AND EVERY page.
- Students are to attempt ALL questions.
ALL questions are of equal value. [12 marks]
- The questions are not necessarily arranged in order of difficulty.
Students are advised to read the whole paper carefully at the start of the examination.
- ALL necessary working should be shown in every question.
Marks may be deducted for careless or badly arranged work.
- Begin your answer to each question on a NEW page. The answers to the questions in this paper are to be returned in SEVEN SEPARATE BUNDLES.
Write on ONLY ONE SIDE of each page.
- Approved calculators and geometrical instruments may be used.
- A table of Standard Integrals is provided at the end of the paper.

* * * *

QUESTION 1.

(a) Solve for x :

(i) $\frac{x + 4}{x - 2} > 5$ [3m]

(ii) $\left(x + \frac{1}{x}\right)^2 - 5\left(x + \frac{1}{x}\right) + 6 = 0$ [3m]

(b) Differentiate with respect to x :

(i) $\cos^3 2x$ [2m]

(ii) $e^{xt \ln x}$ [2m]

(c) AB is a variable interval. M and N divide AB in ratio $-2 : 1$ and $2 : 1$ respectively.

Draw a diagram and decide in what ratio B divides MN .

[2]

QUESTION 2.

(a) Evaluate: $\lim_{x \rightarrow 0} \frac{\sin 5x}{2x}$ [2m]

(b) (i) Sketch the curve $y = \sin^{-1}(2x)$

(ii) State the domain and range of this function. [3m]

(c) Evaluate: $\int_0^2 \frac{4}{\sqrt{4 - x^2}} dx$ [3m]

(d) Find the obtuse angle, to the nearest minute, between the lines

$3x - 4y + 8 = 0$ and $x + 2y + 1 = 0$ [4m]

QUESTION 3.

(a) Prove: $\frac{\sin \theta + \sin 2\theta}{1 + \cos \theta + \cos 2\theta} = \tan \theta$ [3 m]

(b) By using the substitution $u = \cos x$, or otherwise, evaluate $\int_0^{\frac{\pi}{4}} \tan x dx$ [4 m]

(c) If ${}^9C_4 + {}^9C_5 = {}^{10}C_m$, find the value of m . [1 m]

(d) Find the derivatives of:

(i) $\ln(\sec 3x)$

(ii) $\tan^{-1}(2 \tan x)$ [4 m]

QUESTION 4.

(a) $P(4p, 2p^2)$ is a point on the parabola $x^2 = 8y$ and S is the focus. The tangent to the parabola at P meets the y -axis in M . The perpendicular from the focus S to the tangent PM meets the tangent in N .

(i) Write down the equation of PM and hence show that M has coordinates $(0, -2p^2)$. [1 m]

(ii) Write down the equation of SN and hence find the coordinates of N . [4 m]

(iii) Find the coordinates of the midpoint of the interval MN . [1 m]

(iv) Find the equation of the locus of the midpoint MN as P varies. [1 m]

(b) Use the binomial theorem to find the term in x^5 in the expansion $(1 + 2x)^8$. [2 m]

(c) Give the exact value of $\cos^{-1}\left(\sin \frac{4\pi}{3}\right)$. [3 m]

QUESTION 5.


- (a) Prove, by mathematical induction, that $3^{2^n} - 1$ is divisible by 8 for all positive integers. [3m]
- (b) Rain is falling steadily and is collected in an inverted right cone so that the volume collected increases at a constant rate of 5 cm³/h. If the radius r cm of the surface of the water is one third its depth, y cm, find the rate in cm/h at which the depth is increasing when $y = 3.5$. [5m]
- (c) Find all angles θ with $0 \leq \theta \leq 2\pi$ for which $\cos 2\theta = \cos \theta$. [4m]

QUESTION 6.

- (a) Find the term independent of x in the expansion of $\frac{1}{x} \left(3x - \frac{1}{2x} \right)^7$. [3m]
- (b) A particle moves in a straight line and its position at any time t is given by:
$$x = 2\cos 3t - 5\sin 3t.$$
- (i) Find the acceleration in terms of position and hence show that the motion is simple harmonic.
- (ii) Find the greatest speed of the particle. [5m]
- (c) (i) Show that $\frac{d}{dx} [e^x (\sin x + \cos x)] = 2e^x \cos x$. [4m]
- (ii) Hence, evaluate: $\int_1^{\frac{\pi}{2}} e^x \cos x dx$ (correct to 3 significant figures). [4m]

QUESTION 7.

(a)



The angle of elevation of a tower PQ , of height h metres, at a point A due east of it, is 17° . From another point B , the bearing of the tower is $061^\circ T$ and the angle of elevation is 15° . The points A and B are 1000 metres apart and on the same level as the base Q of the tower.

- (i) Show that $\angle AQB = 151^\circ$.
- (ii) Consider the ΔAPQ and show that $AQ = h \tan 73^\circ$.
- (iii) Find a similar expression for BQ .
- (iv) Calculate h , using the cosine rule, in the ΔAQB .
(Answer to nearest metre).

[6m]

- (b) A cricket ball is projected from the ground with an initial velocity of 30 ms^{-1} at an angle of 40° to the horizontal. The equations of motion taken in the horizontal and vertical directions are $\ddot{x} = 0$, $\ddot{y} = -10$. (Use $g = 10 \text{ ms}^{-2}$).

- (i) Calculate the greatest height reached by the ball.
- (ii) What is the speed of the ball at the greatest height?
- (iii) How high is it after the ball has travelled 40 metres horizontally? [6m]

E N D O F E X A M


ANS (3u) Trial 2000.

Q1. (a) (i) Method 1: $x(x-2)^2$:
 $(x+4)(x-2) > 5(x-2)^2$
 $x^2 + 2x - 8 > 5(x^2 - 4x + 4)$
 $0 > 4x^2 - 22x + 28$

i.e. $2x^2 - 11x + 14 < 0$


$(2x - 7)(x - 2) < 0$

$\therefore \boxed{2 < x < 3\frac{1}{2}}$



(c)

Method 2: sketch $y = \frac{x-2+6}{x-2} = 1 + \frac{6}{x-2}$.



Inters: $x+4 = 5(x-2)$.
 $= 5x - 10$
 $4x - 10 = 0$
 $x = \frac{10}{4} = \frac{5}{2} = 3\frac{1}{2}$.

Method 3: cases:

For $x > 2$: $x+4 > 5(x-2) \Rightarrow x+4 > 5x-10$.
 $i.e. 4x-14 < 0 \therefore x < 3\frac{1}{2}$

$\therefore 2 < x < 3\frac{1}{2}$ is part sol.

For $x < 2$: $x+4 < 5(x-2) \Rightarrow x > 3\frac{1}{2}$
no part sol. here

$\therefore \boxed{2 < x < 3\frac{1}{2}}$

(ii) $y^2 - 5y + 6 = 0 \Rightarrow (y-2)(y-3) = 0$
 $\therefore x + \frac{1}{x} = 2, 3$

$x^2 - 2x + 1 = 0$ or $x^2 - 3x + 1 = 0$
 $(x-1)^2 = 0$ $x = \frac{3 \pm \sqrt{9-4}}{2}$


$\therefore x = 1, \frac{3 \pm \sqrt{5}}{2}$

Q1. (b) (i) $y = \cos^3 2x$

$y' = 3 \cos^2 2x \cdot -\sin 2x \cdot 2$ ← 2 marks, 1 off each mistake.
 $= -6 \sin 2x \cdot \cos^2 2x$

(ii) $y = e^{x \ln x}$

$y' = (1 \cdot \ln x + x \cdot \frac{1}{x}) e^{x \ln x}$ ← 2 marks, 1 off each mistake.
 $= (1 + \ln x) e^{x \ln x}$




B divides MN in ratio 3:1

Q2. (a) $I = \lim_{x \rightarrow 0} \frac{\sin 5x}{2x} = \lim_{5x \rightarrow 0} \frac{\sin 5x}{5x} \cdot \frac{5}{2}$

$$= 1 \times \frac{5}{2} = \frac{5}{2}$$

(b) (i) $-1 \leq 2x \leq 1 \Rightarrow -\frac{1}{2} \leq x \leq \frac{1}{2}$



(ii) Domain $-\frac{1}{2} \leq x \leq \frac{1}{2}$

Range $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$

(c) $I = \int_0^2 \frac{4 dx}{\sqrt{4-x^2}} = 4 \left[\sin^{-1} \frac{x}{2} \right]_0^2$

$$= 4 \left\{ \sin^{-1} 1, -\sin^{-1} 0 \right\}$$

$$= 4 \left\{ \frac{\pi}{2}, 0 \right\}$$

$$= 2\pi$$

(d) $m_1 = \frac{3}{4}, m_2 = -\frac{1}{2}$

$$\tan \theta = \left| \frac{\frac{3}{4} - -\frac{1}{2}}{1 + \frac{3}{4}(-\frac{1}{2})} \right| = \frac{5/4}{5/8} = 2$$

acute.

$$\therefore \theta = 180^\circ - 63^\circ 26' = 116^\circ 34'$$

Q3.

(a) LHS = $\frac{\sin \theta + 2 \sin \theta \cos \theta}{1 + \cos \theta + 2 \cos^2 \theta - 1}$

$$= \frac{\sin \theta (1 + 2 \cos \theta)}{\cos \theta}$$

$$= \tan \theta$$

$$= RHS.$$

(b) $u = \cos x, x = \frac{\pi}{3} \Rightarrow u = \frac{1}{2}$

$$du = -\sin x dx, x = 0 \Rightarrow u = 1$$

$$I = - \int_0^{\frac{1}{2}} \frac{-\sin x dx}{\cos x}$$

$$= - \int_1^{\frac{1}{2}} \frac{du}{u}$$

$$= \int_{\frac{1}{2}}^1 \frac{du}{u}$$

$$= [\ln u]_{\frac{1}{2}}^1$$

$$= \ln 1 - \ln \frac{1}{2}$$

$$= 0 - -\ln 2$$

$$= \ln 2$$

(c) LHS = $\frac{9!}{5!4!} + \frac{9!}{4!5!} = \frac{2 \times 9!}{5!4!} \times \frac{5}{5} = \frac{10!}{5!5!} = {}^{10}C_5$

$$\therefore m=5 \quad [\text{note bald answer OK}]$$

(d) (i) $\frac{d}{dx} (\ln(\sec 3x)) = \frac{3 \times \sec 3x \tan 3x}{\sec 3x}$

$$= 3 \tan 3x$$

(ii) $\frac{d}{dx} (\tan^{-1}(\tan x)) = \frac{2 \sec^2 x}{1 + 4 \tan^2 x}$


Q4.

$$(a) (i) \text{ PM is } x+4p = 4(y+2p^2) \quad a=2$$

$$\text{ie. } px = y + 2p^2$$

$$\text{cuts y-axis: } x=0 \quad \therefore y = -2p^2$$

$$\text{ie. M is } (0, -2p^2)$$



$$(ii) \text{ gr. PM} = p$$

$$\therefore \text{SN is } y-2 = -\frac{1}{p}(x-0)$$

$$\text{ie. } y = 2 - \frac{x}{p}.$$

$$N: px = (2 - \frac{x}{p}) + 2p^2$$

$$\text{ie. } p^2x = 2p - x + 2p^3$$

$$x(p^2+1) = 2p(p^2+1)$$

$$\therefore x = 2p, \text{ since } p^2+1 > 0$$

$$\text{ie. } y = 2 - \frac{2p}{p} = 0$$

$$\therefore N \text{ is } (2p, 0)$$

$$(iii) \text{ mid pt of MN: } (\frac{0+2p}{2}, \frac{-2p^2+0}{2})$$

$$\text{ie. } (p, -p^2)$$

$$(iv) \text{ locus: } y = -p^2 = -x^2$$

$$\text{ie. } y = -x^2$$

$$(b) (1+2x)^8 = \binom{8}{0} + \binom{8}{1}(2x) + \dots + \binom{8}{5}(2x)^5 + \dots + (2x)^8$$

$$\therefore \text{coeff of } x^5 \text{ is } \binom{8}{5} \times 2^5 = \frac{8 \times 7 \times 6}{1 \times 2 \times 3} \times 32 = 7 \times 256$$

$$= 1792.$$

$$(c) \sin \frac{4\pi}{3} = -\sin \frac{\pi}{3} = -\frac{\sqrt{3}}{2}, \quad \cos^{-1}(-\frac{\sqrt{3}}{2}) \quad \text{Anis} = \pi - \pi/6 = \frac{5\pi}{6}$$

$$QS.(a) n=1 \Rightarrow 3^{2^1}-1 = 9-1=8 \quad \therefore \text{div by 8 when } n=1$$

$$\text{Say } (n=k), \quad 3^{2^k}-1 = 8P \text{ for some pos. int. } k, P$$

$$\text{then } 3^{2(k+1)}-1 = 3^{2k+2}-1$$

$$= 9 \times (3^{2k}-1) + 8$$

$$= 9 \times 8P + 8$$

$$= 8(9P+1) \quad \& (9P+1) \text{ is an int.}$$

\therefore If div by 8 for some value of n then div by 8 for next value of n , and shown true for $n=1 \therefore$ true for all pos. int. n .



$$V = \frac{1}{3}\pi r^2 y$$

$$\frac{dV}{dt} = 5 \text{ cm}^3/\text{s}$$

$$\text{want } \frac{dy}{dt} \text{ when } y=3.5$$

$$\therefore V = \frac{\pi}{3} \cdot \frac{4}{3}^3 = \frac{\pi 4^3}{9}$$

$$\frac{dV}{dy} = \frac{\pi 4^2}{3}$$

$$\text{And. } \frac{dV}{dt} = \frac{dV}{dy} \cdot \frac{dy}{dt}$$

$$5 = \frac{\pi 4^2}{3} \cdot \frac{dy}{dt}$$

$$\therefore \frac{dy}{dt} = 5 \times \frac{3}{\pi \times 8.5^2} \text{ at } y=3.5$$

$$\therefore 1.2 \text{ cm/l.}$$

$$(c) 2\cos^2\theta - \cos\theta - 1 = 0$$

$$(2\cos\theta + 1)(\cos\theta - 1) = 0$$

$$\cos\theta = -\frac{1}{2}, 1$$

$$\theta = \pi - \pi/3, \pi + \pi/3, 0, 2\pi$$

$$\therefore \theta = \frac{2\pi}{3}, \frac{4\pi}{3}, 0, 2\pi$$

$$Q6(a) \quad \frac{1}{x} \cdot \binom{7}{k} (3x)^{7-k} \left(-\frac{1}{2x}\right)^k$$

$$\text{we want } x^{-1} \times x^{7-k} \times x^{-k} = 1 = x^0$$

$$\therefore 6 - 2k = 0$$

$$\therefore k = 3$$

$\therefore \left(\frac{7}{3}\right) 3^{7-3} \left(-\frac{1}{2}\right)^3$ is the req. term

$$\therefore - \frac{7 \times 4 \times 5}{1 \times 2 \times 3} \times \frac{3^4}{2^3} = - \frac{35 \times 81}{8}$$

$$\therefore -354 \frac{3}{8}$$

$$(b) \quad x = 2 \cos 3t - 5 \sin 3t$$

$$\dot{x} = -6 \sin 3t - 15 \cos 3t$$

$$\ddot{x} = -18 \cos 3t + 45 \sin 3t$$

$$(i) \therefore \ddot{x} = -9 \text{ m/s} \quad \text{which is SHM}$$

(ii) max speed is when $\ddot{x} = 0$

$$\text{i.e. } x=0 \quad \therefore 2 \cos 3t = 5 \sin 3t$$

$$\therefore \frac{2}{5} = \tan 3t$$

$$\& \text{max speed} = \left| -6 \times \frac{2}{\sqrt{29}} - 15 \times \frac{5}{\sqrt{29}} \right|$$

$$= \frac{87}{\sqrt{29}} \div 16 \approx 155 \div 16 \text{ speed units}$$

$$(c)(i) \quad \frac{d}{dx} [e^x (\sin x + \cos x)] = e^x (\sin x + \cos x) + e^x (\cos x - \sin x) \\ = 2e^x \cos x. \quad \text{answ.}$$

$$(ii) I = \int_{1}^{\pi/2} e^x \cos x dx$$


$$= \frac{1}{2} \int_{1}^{\pi/2} 2e^x \cos x dx.$$

$$= \frac{1}{2} \left[e^x (\sin x + \cos x) \right]_{1}^{\pi/2}$$

$$= \frac{1}{2} \left\{ e^{\pi/2} (1+0) - e^1 (\sin 1 + \cos 1) \right\}$$

$$\div 0.527$$

Q7.(a)



$$(i) \quad \angle AQB = 61^\circ + 90^\circ = 151^\circ$$

$$(ii) \quad \text{in } \triangle APQ: \tan 73^\circ = \frac{AQ}{h} \quad \therefore AQ = h \tan 73^\circ$$

$$(iii) \quad \text{in } \triangle BPQ: \tan 75^\circ = \frac{BQ}{h} \quad \therefore BQ = h \tan 75^\circ$$


(iv) in $\triangle ABQ$:

$$1000^2 = (h \tan 73^\circ)^2 + (h \tan 75^\circ)^2 - 2(h \tan 73^\circ)(h \tan 75^\circ) \cos 151^\circ \\ = h^2 [\tan^2 73^\circ + \tan^2 75^\circ - 2 \tan 73^\circ \tan 75^\circ \cos 151^\circ]$$

$$= h^2 \times 45.9796 \dots$$

$$\therefore h = \frac{1000}{\sqrt{45.9796 \dots}} = 147.47 \dots$$

$$= 147 \text{ m.}$$



$$\begin{cases} \ddot{x} = 0 & y = -10 \\ \dot{x} = 30 \cos 40^\circ & y = -10t + 30 \sin 40^\circ \\ x = 30t \cos 40^\circ & y = -5t^2 + 30t \sin 40^\circ \end{cases}$$

(i) max ht: when $\dot{y} = 0$: $t = 3 \sin 40^\circ$

$$\text{then } ht = -5(3 \sin 40^\circ)^2 + 90 \sin^2 40^\circ \\ = 45 \sin^2 40^\circ \\ \div 18.6 \text{ m.}$$

(ii) Speed at top pt $(\dot{y}=0)$: $\dot{x} = 30 \cos 40^\circ \div 23 \text{ m/s}$

$$(iii) x = 40 \Rightarrow t = \frac{4}{3 \cos 40^\circ} \Rightarrow y = -5 \left(\frac{4}{3 \cos 40^\circ} \right)^2 + \frac{4}{3 \cos 40^\circ} \times \frac{80 \sin 40^\circ}{23} \\ = -16.14743 \dots + 33.56 \dots \\ \div 18.4 \text{ m.}$$