

# NEWCASTLE GRAMMAR SCHOOL

Student Number: \_\_\_\_\_



## 2011 TRIAL HIGHER SCHOOL CERTIFICATE EXAMINATION

# Mathematics Extension 1

*Examination Date: Friday 19th August*

Examiner: Mr. M. Brain

### General Instructions

- Reading Time - 5 minutes
- Working Time - 2 hours
- Write using a blue or black pen
- Write your student number on every booklet
- Board-approved calculators may be used
- A table of standard integrals is provided in this paper
- All necessary working should be shown in every question
- Each question attempted is to be returned in a separate Writing Booklet clearly marked Questions 1 etc.
- If required, additional booklets may be requested

### Total marks - 84

- Attempt Questions 1- 7
- All questions are of equal value

**Question 1** (Start a new booklet)

**Marks**

- a) If  $y = (\tan^{-1} x)^2$  find  $\frac{dy}{dx}$  **2**
- b) Find the value of  $\sum_{n=2}^5 {}^n C_2$  **2**
- c) Solve  $\frac{2x-3}{x-2} \geq 1$  **4**
- d) Find  $\lim_{x \rightarrow 0} \frac{\sin 3x}{x}$  **1**
- e) Evaluate  $\int_0^3 \frac{x}{\sqrt{1+x}} dx$  using the substitution  $x = u^2 - 1$ , where  $u > 0$  **3**

**Question 2** (Start a new booklet)

**Marks**

- a) Find the size of the acute angle between the two lines with equations  $x + 2y + 1 = 0$  and  $2x - 3y + 6 = 0$ , correct to the nearest degree **4**
- b) A spherical balloon is expanding so that its volume,  $V \text{ cm}^3$ , increases at a constant rate of  $72 \text{ cm}^3$  per second. What is the rate of increase of the surface area when the radius is 12 cm? **4**
- c) Find **4**
- (i)  $\int \sin^2 x \, dx$
- (ii)  $\int \frac{dx}{\sqrt{9 + 4x^2}}$

**Question 3** (Start a new booklet)

**Marks**

- a) The rate of growth of a bacteria colony is proportional to the excess of the colony's population over 5000 and is given by

**5**

$$\frac{dN}{dt} = k(N - 5000)$$

- (i) Show that  $N = 5000 + Ae^{kt}$  is a solution to the differential equation above
- (ii) Given that the initial population was 15000 and had risen to 20000 after 2 days find the value of  $A$  and  $k$
- (iii) Hence calculate the expected population after 7 days

- b) The polynomial  $P(x) = x^5 + mx^3 + nx$  has a remainder of 5 when divided by  $(x - 2)$ , where  $m$  and  $n$  are constants.

**3**

- (i) Prove that  $P(x)$  is an odd function
- (ii) Hence find the remainder when  $P(x)$  is divided by  $(x + 2)$

- c) If  $\alpha, \beta$  and  $\gamma$  are the roots of the equation  $x^3 - 2x^2 + 3x + 7 = 0$  find the value of  $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}$


**4**

**Question 4** (Start a new booklet)

**Marks**

- a)  $ABC$  is a triangle inscribed in a circle.  $M$  is a point on the tangent to the circle at  $B$  and  $N$  is a point on  $AC$  produced so that  $MN$  is parallel to  $BA$

**4**



Copy the diagram into your answer booklet

- (i) State why  $\angle MBC = \angle BAC$
- (ii) Prove that  $MNCB$  is a cyclic quadrilateral
- b) Prove by mathematical induction that, for all integers  $n \geq 1$

**4**

$$1 \times 2^0 + 2 \times 2^1 + 3 \times 2^2 + \dots + n \times 2^{n-1} = 1 + (n-1)2^n$$

- c) (i) Prove that the equation  $\frac{\log_e x}{x} + 2 = 0$  has a solution between  $x = 0.4$  and  $x = 0.5$
- (ii) Use one application of Newton's method to find a closer approximation to the solution  $x = 0.4$ , correct to three decimal places

**4**

**Question 5** ( Start a new booklet)

**Marks**

- a) In Group A there are 5 men and 3 women. In Group B there are 4 men and 6 women. **4**
- (i) If one person is chosen at random from each group what is the probability that the two people chosen are of opposite sexes?
- (ii) If a group and then one person from that group is chosen at random what is the probability that the person chosen is a man?
- 
- b) Using  $\tan \frac{\theta}{2} = t$  show that  $\frac{1 - \cos \theta}{\sin \theta} = t$  **3**
- 
- c) A particle moves in a straight line such that its acceleration,  $a$ , is given by  $a = 3x^2$ , where  $x$  is displacement,  $v$  is velocity and  $t$  is time. Given that  $v = -\sqrt{2}$  and  $x = 1$  when  $t = 0$  find  $x$  as a function of time,  $t$  **5**

**Question 6** (Start a new booklet)

**Marks**

a) Consider the function  $f(x) = 3 \sin^{-1}\left(\frac{x}{2}\right)$  **5**

- (i) State the domain and range of this function
- (ii) Sketch the graph of  $y = f(x)$
- (iii) Find the slope of the graph at  $x = 0$

b) The displacement,  $x$  metres, of a particle, at  $t$  seconds is given by: **7**

$$x = 5 \cos(4\pi t)$$

- (i) Show that the acceleration of the particle can be expressed in the form:


$$\ddot{x} = -n^2 x$$

- (ii) State the period,  $P$ , of the motion
- (iii) Determine the maximum velocity of the particle
- (iv) Determine the maximum acceleration of the particle
- (v) Express  $v^2$  in terms of  $x$ , where  $v$  is the velocity of the particle

**QUESTION 7 IS ON THE NEXT PAGE**

- a) A vertical building stands with its base  $O$  on horizontal ground.  $A$  and  $B$  are two points on the building vertically above each other such that  $A$  is  $4h$  metres above  $O$  and  $B$  is  $h$  metres above  $O$ . A particle is projected horizontally with speed  $U$  metres per second from  $A$  and 10 seconds later a second particle is projected horizontally with speed  $V$  metres per second from  $B$ . The two particles hit the ground at the same point and at the same time.

**7**



- (i) Show that the horizontal and vertical displacements of particles  $A$  and  $B$ ,  $t$  seconds after the first particle is projected, are given by  $x_A = Ut$ ,  $y_A = 4h - \frac{1}{2}gt^2$  and  $x_B = V(t-10)$ ,  $y_B = h - \frac{1}{2}g(t-10)^2$ , respectively
- (ii) Find the time of flight of each particle
- (iii) Prove that  $V = 2U$
- b) The point  $P(2t, t^2)$  is on the parabola with equation  $x^2 = 4y$ , having its focus at  $F$ . The point  $M$  divides the interval  $FP$  externally in the ratio 3:1.
- (i) Show that the co-ordinates of  $M$  are  $x = 3t$  and  $y = \frac{1}{2}(3t^2 - 1)$
- (ii) Hence prove that the locus of  $M$  is also a parabola and determine the focal length of the locus of  $M$

**5**

## STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left( x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left( x + \sqrt{x^2 + a^2} \right)$$

NOTE :  $\ln x = \log_e x, \quad x > 0$

① a)  $y = (\tan^{-1} x)^2$

$$\therefore \frac{dy}{dx} = 2(\tan^{-1} x)^1 \times \frac{d}{dx}(\tan^{-1} x)$$

$$= 2 \tan^{-1} x \times \frac{1}{1+x^2}$$

$$= \frac{2 \tan^{-1} x}{1+x^2} \quad \text{①/2}$$

b)  $\sum_{n=2}^5 {}^n C_2 = {}^2 C_2 + {}^3 C_2 + {}^4 C_2 + {}^5 C_2$

$$= 1 + 3 + 6 + 10$$

(using calculator)

$$= \textcircled{20} \quad \text{①/2}$$

c)  $\frac{2x-3}{x-2} \geq 1$  x B.S. by (denom)<sup>2</sup>


$$\therefore \frac{2x-3}{x-2} \times (x-2)^2 \geq 1 \times (x-2)^2$$

$$\therefore (2x-3)(x-2) \geq x^2 - 4x + 4$$

$$\therefore 2x^2 - 4x - 3x + 6 \geq x^2 - 4x + 4$$

$$\therefore x^2 - 3x + 2 \geq 0$$

Consider graph of  $y = x^2 - 3x + 2$   
 $= (x-1)(x-2)$



$$\therefore x \leq 1 \text{ or } x \geq 2$$

but  $x \neq 2 \therefore x \leq 1 \text{ or } x > 2$  ①/4

d)  $\lim_{x \rightarrow 0} \frac{\sin 3x}{x} = \lim_{x \rightarrow 0} 3 \times \frac{\sin 3x}{3x}$

$$= 3 \times \lim_{x \rightarrow 0} \frac{\sin 3x}{3x} \quad \left. \begin{array}{l} \text{as } x \rightarrow 0 \\ 3x \rightarrow 0 \end{array} \right\}$$

$$= 3 \times 1$$

$$= \textcircled{3} \quad \text{①}$$

e)  $\int_0^3 \frac{x}{\sqrt{1+x}} dx$

$$x = u^2 - 1$$

$$\therefore dx = 2u du$$

$$= \int_1^2 \frac{u^2-1}{\sqrt{1+u^2-1}} \times 2u du$$

at  $x=3: u^2=4$   
 $u=2 (>0)$

$$= \int_1^2 \frac{u^2-1}{u} \times 2u du$$

$x=0: u^2=1$   
 $u=1 (>0)$

$$= 2 \int_1^2 (u^2-1) du$$

$$= 2 \left[ \frac{u^3}{3} - u \right]_1^2$$

$$= 2 \left( \left( \frac{8}{3} - 2 \right) - \left( \frac{1}{3} - 1 \right) \right)$$

$$= \textcircled{\frac{8}{3}} \quad \text{③}$$

② a)  $\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$

$m_1: 2y = -x - 1 \therefore y = -\frac{1}{2}x - \frac{1}{2} \therefore m_1 = -\frac{1}{2}$

$m_2: 3y = 2x + 6 \therefore y = \frac{2}{3}x + 2 \therefore m_2 = \frac{2}{3}$

$$\therefore \tan \theta = \left| \frac{-\frac{1}{2} - \frac{2}{3}}{1 - \frac{1}{2} \times \frac{2}{3}} \right|$$

$$= \left| \frac{-7}{4} \right|$$

$$= \frac{7}{4}$$

$$\therefore \theta = \tan^{-1} \left( \frac{7}{4} \right)$$

$$= 60.25 \dots$$

$$\theta = \textcircled{60^\circ} \text{ (nearest deg.)} \quad \text{④}$$

② b) S.A. = A, Volume = V

$$\therefore \frac{dA}{dt} = \frac{dA}{dV} \times \frac{dV}{dt} \dots \dots (1)$$

where  $\frac{dV}{dt} = 72 \text{ cm}^3/\text{s} \dots \dots (2)$

Now:  $\frac{dA}{dV} = \frac{dA}{dr} \times \frac{dr}{dV}$

where:  $A = 4\pi r^2 \therefore \frac{dA}{dr} = 8\pi r \dots \dots (3)$

$V = \frac{4}{3}\pi r^3 \therefore \frac{dV}{dr} = 4\pi r^2$

or  $\frac{dr}{dV} = \frac{1}{4\pi r^2} \dots \dots (4)$

$\therefore$  (2), (3), (4) into (1):

$$\begin{aligned} \frac{dA}{dt} &= 8\pi r \times \frac{1}{4\pi r^2} \times 72 \\ &= \frac{144}{r} \end{aligned}$$

$\therefore$  when  $r = 12$

$$\begin{aligned} \frac{dA}{dr} &= \frac{144}{12} \\ &= \boxed{12 \text{ cm}^2/\text{s}} \quad \textcircled{4} \end{aligned}$$

c) (i)  $\int \sin^2 x \, dx = \frac{1}{2} \int (1 - \cos 2x) \, dx$   
 $= \frac{1}{2} (x - \frac{1}{2} \sin 2x) + C$

or  $= \boxed{\frac{1}{2}x - \frac{1}{4} \sin 2x + C} \quad \textcircled{2}$

(ii)  $\int \frac{dx}{\sqrt{9+4x^2}} = \int \frac{1}{\sqrt{4(\frac{9}{4}+x^2)}} \, dx$

Using Standard Integrals sheet  $\rightarrow = \frac{1}{2} \int \frac{1}{\sqrt{(\frac{3}{2})^2+x^2}} \, dx$   
 $\rightarrow = \boxed{\frac{1}{2} \ln(x + \sqrt{9+4x^2})} \quad \textcircled{2}$

③ a) (i)

If  $N = 5000 + Ae^{kt} \dots \dots (1)$

then  $\frac{dN}{dt} = k \times Ae^{kt}$

but from (1):  $Ae^{kt} = N - 5000$

$\therefore \frac{dN}{dt} = k(N - 5000)$

$\therefore N = 5000 + Ae^{kt}$  is a solution to  $\frac{dN}{dt} = k(N - 5000)$  (QED)  $\textcircled{1}$

(ii) At  $t=0$ ,  $N = 15000$

$\therefore 15000 = 5000 + Ae^{k \times 0}$   
 $= 5000 + A$

$\therefore \boxed{A = 10000}$

At  $t=2$ ,  $N = 20000$

$\therefore 20000 = 5000 + 10000e^{2k}$

$\therefore 10000e^{2k} = 15000$

$e^{2k} = 1.5$

$\ln(e^{2k}) = \ln 1.5$

$\therefore 2k = \ln 1.5$

$k = \ln 1.5 \div 2$

$= 0.20273 \dots$

$\boxed{k = 0.2027} \quad \text{(4dp)} \quad \textcircled{3}$

(iii)  $\therefore$  At  $t=7$ :

$N = 5000 + 10000e^{0.2027 \times 7}$

$= 46325.7 \dots$

$\therefore \boxed{N = 46300} \quad \text{(nearest 100)} \quad \textcircled{1}$

b)  $P(x) = x^5 + mx^3 + nx$

(i)  $\therefore P(a) = a^5 + ma^3 + na$

$$P(-a) = (-a)^5 + m(-a)^3 + n(-a)$$

$$= -a^5 - ma^3 - na$$

$$= -(a^5 + ma^3 + na)$$

$$\therefore P(-a) = -\{-(a^5 + ma^3 + na)\}$$

$$= a^5 + ma^3 + na$$

i.e.  $P(a) = -P(-a) \therefore$  odd (QED) (1)

(ii) When  $P(x)$  divided by  $x+2$  remainder is  $P(-2)$

AND when divided by  $x-2$

remainder is  $P(2) = 5$  (given)

Now, from (i)  $P(a) = -P(-a)$

$$\therefore P(-2) = -P(-(-2))$$

$$= -P(2)$$

$$= -5 \quad (2)$$

c)  $\alpha, \beta, \gamma$  : roots of  $x^3 - 2x^2 + 3x + 7 = 0$

and:  $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma} = \frac{\alpha\beta + \alpha\gamma + \beta\gamma}{\alpha\beta\gamma}$

$$= \frac{\alpha\beta + \alpha\gamma + \beta\gamma}{\alpha\beta\gamma}$$

Now:  $\alpha\beta + \alpha\gamma + \beta\gamma = c/a$

$$= 3/1$$

$$= 3$$

$$\alpha\beta\gamma = -d/a$$

$$= -7/1$$

$$= -7$$

$$\therefore \frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma} = \frac{3}{-7} \text{ or } -\frac{3}{7} \quad (4)$$

(4) a) (i)  $\angle MBC = \angle BAC$

$\therefore$  angle between tangent and chord equals angle in alternate segment (1)

(ii)  $\angle BAC + \angle CNM = 180^\circ$

(co-int.  $\angle$ 's in  $\parallel$  lines)

and  $\angle BAC = \angle MBC$  (from (i))

$\therefore \angle MBC + \angle CNM = 180^\circ$

$\therefore$  MNCB is cyclic quadrilateral

(opp.  $\angle$ 's add to  $180^\circ$ ) (3)

b) Prove:  $1 \times 2^0 + 2 \times 2^1 + 3 \times 2^2 + \dots + n \times 2^{n-1} = 1 + (n-1)2^n$   
for  $n \geq 1$

• Prove true for  $n=1$

i.e. LHS =  $1 \times 2^0$       RHS =  $1 + (1-1) \times 2^1$

$$= 1 \times 1 \quad = 1 + 0$$

$$= 1 \quad = 1 \quad (1/2)$$

$\therefore$  true for  $n=1$

• Assume true for  $n=k$

i.e. assume:  $1 \times 2^0 + 2 \times 2^1 + \dots + k \times 2^{k-1} = 1 + (k-1)2^k$

$\therefore$  prove true for  $n=k+1$

i.e. prove:  $1 \times 2^0 + 2 \times 2^1 + \dots + k \times 2^{k-1} + (k+1)2^k = 1 + k \times 2^{k+1}$

Now: LHS =  $1 + (k-1)2^k + (k+1)2^k$  {from assumption.

$$= 1 + k \times 2^k - 2^k + k \times 2^k + 2^k$$

$$= 1 + 2k \times 2^k$$

$$= 1 + k \times 2^1 \times 2^k$$

$$= 1 + k \times 2^{k+1}$$

$$= \text{RHS}$$

$\therefore$  LHS = RHS (3)

•  $\therefore$  True for  $n=1$  and true for  $n=k+1$  when true for  $n=k$  (1/2)

$\therefore$  True for  $n=1, 2, 3, \dots$  (all  $n \geq 1$ ) (4)

(QED)

④ c) (i) Solution to  $\frac{\log_e x}{x} + 2 = 0$   
 corresponds to  $y = \frac{\log_e x}{x} + 2$  crossing  
 x-axis ( $y=0$ )

Now: when  $x=0.4$ :

$$\frac{\log_e x}{x} + 2 = \frac{\log_e 0.4}{0.4} + 2$$

$$= -0.29 \dots \text{ (below x-axis)}$$

when  $x=0.5$ :

$$\frac{\log_e x}{x} + 2 = +0.61 \dots \text{ (above x-axis)}$$

$\therefore$  **Sign change**  $\Rightarrow$  solution between  $x=0.4$  and  $x=0.5$  (QED)

(ii) For  $f(x) = \frac{\log_e x}{x} + 2$ :

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$$

$$\text{where } f'(x) = \frac{x \times \frac{1}{x} - \log_e x \times 1}{x^2}$$

$$= \frac{1 - \log_e x}{x^2}$$

$$\therefore x_2 = 0.4 - \frac{\left(\frac{\log_e 0.4}{0.4} + 2\right)}{\frac{1 - \log_e 0.4}{(0.4)^2}}$$

$$= 0.4242 \dots$$

$$\therefore x_2 = 0.424 \text{ (3dp)} \quad \text{③}$$

⑤ a)  $A = 5m, 3w$   
 $B = 4m, 6w$

(i)  $P(\text{opposite sexes}) = P(M, w \text{ or } wM)$

$$= P(M, w) + P(w, M)$$

$$= \frac{5}{8} \times \frac{6}{10} + \frac{3}{8} \times \frac{4}{10}$$

$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow$   
 A    B    A    B

$$= \frac{21}{40} \quad \text{②}$$

(ii)  $P(\text{man}) = P(\text{Group A, M or Group B, M})$

$$= P(\text{Group A, M}) + P(\text{Group B, M})$$

$$= \frac{1}{2} \times \frac{5}{8} + \frac{1}{2} \times \frac{4}{10}$$


$$= \frac{41}{80} \quad \text{②}$$

b)  $\tan \frac{\theta}{2} = t$

$$\therefore \tan(2 \times \frac{\theta}{2}) = \tan \theta$$

$$= \frac{2 \tan \frac{\theta}{2}}{1 - \tan^2 \frac{\theta}{2}}$$

$$= \frac{2t}{1-t^2}$$



$$\therefore \frac{1 - \cos \theta}{\sin \theta} = \frac{1 - \frac{1-t^2}{1+t^2}}{\frac{2t}{1+t^2}} \times \frac{1+t^2}{1+t^2}$$

$$= \frac{1+t^2 - 1+t^2}{2t}$$

$$= \frac{2t^2}{2t}$$

$$= t$$

$\therefore$  LHS = RHS (QED) ③

$$(5) c) \frac{d}{dx} \left( \frac{1}{2} v^2 \right) = a = 3x^2$$

$$\therefore \frac{1}{2} v^2 = \int 3x^2 dx$$

$$= x^3 + C$$

( $C_1 = 2x^3$ )

$$\therefore v^2 = 2x^3 + C_1$$

and at:  $v = -\sqrt{2}$ ,  $x = 1$

$$\therefore 2 = 2 + C_1$$

$$\therefore C_1 = 0$$

i.e.  $v^2 = 2x^3$

$$\therefore v = \pm \sqrt{2x^3}$$

But initially  $v = -\sqrt{2}$

$\therefore$  select:  $v = -\sqrt{2x^3}$

i.e.  $\frac{dx}{dt} = -\sqrt{2} x^{3/2}$

$$\therefore \frac{dt}{dx} = -\frac{1}{\sqrt{2}} x^{-3/2}$$

$$\therefore t = -\frac{1}{\sqrt{2}} \int x^{-3/2} dx$$

$$= -\frac{1}{\sqrt{2}} \times \frac{x^{-1/2}}{-1/2} + C_2$$

$$= \frac{2}{\sqrt{2}} x^{-1/2} + C_2$$

i.e.  $t = \sqrt{2} x^{-1/2} + C_2$

and at:  $t = 0$ ,  $x = 1$

$$\therefore 0 = \sqrt{2} + C_2$$

$$\therefore C_2 = -\sqrt{2}$$

$$\therefore t = \frac{\sqrt{2}}{\sqrt{x}} - \sqrt{2}$$

$$\therefore \sqrt{x} = \frac{\sqrt{2}}{t + \sqrt{2}}$$

$$\therefore x = \frac{2}{(t + \sqrt{2})^2} \quad (5)$$

$$(6) a) (i) \text{ Domain:}$$

$$y = \sin^{-1} x : -1 \leq x \leq 1$$

$$\therefore y = \sin^{-1} \left( \frac{x}{2} \right) : -1 \leq \frac{x}{2} \leq 1 \quad (x2)$$

$$\therefore -2 \leq x \leq 2$$

Range:


$$y = \sin^{-1} x : -\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$$

$$y = 3 \sin^{-1} x$$

i.e.  $\frac{y}{3} = \sin^{-1} x : -\frac{\pi}{2} \leq \frac{y}{3} \leq \frac{\pi}{2} \quad (x3)$

$$-\frac{3\pi}{2} \leq y \leq \frac{3\pi}{2} \quad (2)$$

(ii)



$$(iii) f(x) = 3 \sin^{-1} \left( \frac{x}{2} \right) \quad \left. \begin{array}{l} \text{using} \\ \text{Standard} \\ \text{Integrals!} \end{array} \right\}$$

$$\therefore f'(x) = 3 \times \frac{1}{\sqrt{2^2 - x^2}}$$

$$= \frac{3}{\sqrt{4 - x^2}}$$

$\therefore$  Slope at  $x = 0$  is  $f'(0)$

and  $f'(0) = \frac{3}{\sqrt{4 - 0^2}}$

$$= \frac{3}{2} \quad (2)$$

⑥ b)  $x = 5 \cos(4\pi t) \dots (1)$

(i) Express as  $\ddot{x} = -n^2 x$

from (1):  $\dot{x} = 5 \times -\sin(4\pi t) \times 4\pi$   
 $= -20\pi \sin(4\pi t)$

$\therefore \ddot{x} = -20\pi \times \cos(4\pi t) \times 4\pi$   
 $= -80\pi^2 \cos(4\pi t)$

$= -16\pi^2 (5 \cos(4\pi t))$  ②

i.e.  $\ddot{x} = -(4\pi)^2 x = -n^2 x$  (QED)

(ii) Period =  $\frac{2\pi}{n}$

$= \frac{2\pi}{4\pi}$

$= \frac{1}{2}$  (second) ①

(iii) Max Velocity at centre i.e. at  $x=0$

$v^2 = n^2(a^2 - x^2)$

where  $n = 4\pi$ ,  $a = 5$ ,  $x = 0$

$\therefore v^2 = (4\pi)^2 (5^2 - 0)$

$\therefore v = \pm 4\pi \times 5$

$\therefore v_{\max} = 20\pi$  m/s ②

(iv) Max. acceleration at amplitude (left)

i.e. at  $x=5$

$\therefore \ddot{x}_{\max} = -(4\pi)^2 \times 5$

$= 80\pi^2$  m/s<sup>2</sup> ①

(v) from (iii)  $v^2 = (4\pi)^2 (5^2 - x^2)$

$v^2 = 16\pi^2 (25 - x^2)$  ①

⑦ a)

(i) For 'any' projectile motion; fired from (0,0):

Horizontal

Vertical

$\ddot{x} = 0$

$\ddot{y} = -g$

$\therefore \dot{x} = C_1$

$\therefore \dot{y} = -gt + C_2$

but at  $t=0$ ,  $v_x = V \cos \theta$

but at  $t=0$ ,  $v_y = V \sin \theta$

$\therefore C_1 = V \cos \theta$

$\therefore C_2 = V \sin \theta$

$\therefore \dot{x} = V \cos \theta \dots (1)$

$\therefore \dot{y} = V \sin \theta - gt \dots (2)$

$\therefore x = Vt \cos \theta + C_3$

$\therefore y = Vt \sin \theta - \frac{1}{2}gt^2 + C_4$

but at  $t=0$ ,  $x=0$

but at  $t=0$ ,  $y=0$

$\therefore C_3 = 0$

$\therefore C_4 = 0$

$\therefore x = Vt \cos \theta \dots (3)$

$\therefore y = Vt \sin \theta - \frac{1}{2}gt^2 \dots (4)$

$\therefore$  in this question  $\theta = 0$  for both projectiles

$\therefore \sin \theta = 0$ ,  $\cos \theta = 1$

• for projectile A, time =  $t$

$\therefore$  for projectile B, time =  $t - 10$   
 (10 seconds later)

• AND for projectile A:

equation (2) gives:  $y = Vt \sin \theta - \frac{1}{2}gt^2 + C_4$

but at  $t=0$ ,  $y = 4h \therefore C_4 = 4h$

for projectile B

at  $t=0$ ,  $y = h \therefore C_4 = h$

$\therefore$  For A: (3) gives:  $x_A = Ut$

'(4)' gives:  $y_A = 4h - \frac{1}{2}gt^2$

For B: (3) gives:  $x_B = V(t-10)$

$y_B = h - \frac{1}{2}g(t-10)^2$

(QED) ④

⑦(a) (ii) Time of flight : when  $y=0$

ie.  $y(A) = 0$

$y(B) = 0$

ie.  $4h - \frac{1}{2}gt^2 = 0$

$\therefore h = \frac{1}{8}gt^2$  — (1)

and  $h = \frac{1}{2}g(t-10)^2$  — (2)

(1) into (2) :  $\frac{1}{8}gt^2 = \frac{1}{2}g(t-10)^2$

$\therefore t^2 = 4(t-10)^2$

$\therefore t = 2(t-10)$  or  $-t = 2(t-10)$

$\therefore t = 2t - 20 \quad \therefore -t = 2t - 20$

$\therefore t = 20$  on  $t = \frac{20}{3}$

( $t$  = time of flight for A which must be  $> 10$  seconds, for B to be projected 10 seconds after A)

$\therefore t = 20$  seconds

and  $t-10 = 10$  seconds

$\therefore$  Time of flight :  $A = 20$  sec  
 $B = 10$  sec

(iii) When both particles hit ground:

$r_A = r_B$

$\therefore 20u = v(20-10)$

$20u = 10v$

$\therefore v = 2u$  (QED) ①

b)(i)  $x^2 = 4ay \rightarrow x^2 = 4y \quad \therefore a=1$

$\therefore$  focus is  $(0,1)$ , P is  $(2t, t^2)$

Divide externally 3:1  $\left\{ \begin{array}{l} \therefore x = \frac{m x_2 + n x_1}{m+n} \\ \therefore y = \frac{m y_2 + n y_1}{m+n} \end{array} \right.$

$\therefore m:n = 3:-1 \quad \therefore x = \frac{3 \times 2t - 1 \times 0}{3-1}$

ie.  $x = 3t$  — (1)

and  $y = \frac{m y_2 + n y_1}{m+n}$

$\therefore y = \frac{3 \times t^2 - 1 \times 0}{3-1}$

$y = \frac{3t^2 - 1}{2}$  — (2) (QED) ③

(ii) from (1) :  $t = \frac{x}{3}$

sub into (2) :  $y = 3\left(\frac{x}{3}\right)^2 - 1$

$= \frac{x^2}{3} - 1$

$\therefore 2y = \frac{x^2}{3} - 1$

$\therefore 6y = x^2 - 3$  (QED)

Equation of locus :  $x^2 = 6y + 3$  ie parabola ①

and  $x^2 = 6\left(y + \frac{1}{2}\right)$

$= 4 \times \frac{1}{2} \left(y + \frac{1}{2}\right)$

$\therefore$  focal length of parabola is  $\left(\frac{1}{2}\right)$  ①