



Mrs Hickey
Mrs Quarles
Miss Lau
Mrs Stock
Mrs Choong

70

PYRMBLE LADIES' COLLEGE

YEAR 12

MATHEMATICS EXTENSION 1

HSC TRIAL EXAMINATION 2002

Time Allowed: 2 hours + 5 mins reading time

INSTRUCTIONS

- All questions should be attempted
- Write your name and your teacher's name on each page
- Start each question on a new page
- DO NOT staple the questions together
- Only approved calculators may be used
- A standard integral sheet is attached
- Marks might be deducted for careless or untidy work
- Hand this question paper in with your answers
- ALL rough working paper must be attached to the back of the last question
- Staple a coloured sheet of paper to the back of each question
- There are seven (7) questions in this paper
- All questions are of equal value

Question 1

Mark

(a) Evaluate $\lim_{x \rightarrow 0} \frac{\sin 3x}{x}$

1

(b) The point P (7, -1) divides the interval AB externally in the ratio 3 : 2.
If A is (-2, 5) find the coordinates of B.

2

(c) Solve for x

$$\frac{x+1}{x-2} < 2$$

2

(d) Find the gradient of the tangent to the curve $y = \tan^{-1}(2x)$ at the point where $x = \frac{1}{2}$.

2

(e) Evaluate $\int_0^1 \frac{1}{\sqrt{9-x^2}} dx$

2

(f) On the same number plane, sketch the graphs of

(i) $y = |2x - 1|$ and $y = |x + 1|$

2

(ii) Hence, or otherwise, solve $|2x - 1| \leq |x + 1|$

1

Question 2 (Start a new sheet of paper)

- (a) Prove that $\frac{\sin 2\theta}{\sin \theta} - \sec \theta = \frac{\cos 2\theta}{\cos \theta}$

2

- (b) Evaluate $\int_{\frac{1}{2}}^1 4t(2t-1)^3 dt$ by using the substitution $u = 2t-1$

4

- (c) The angle between the lines $y = 3x$ and $y = mx$ is 45° .
Find the value(s) of m .

3

- (d) Solve $\tan 2\theta - \cot \theta = 0$ where $0 \leq \theta \leq \pi$

3

Question 3 (Start a new sheet of paper)

Marks

- (a) Evaluate $\int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \cos^3\left(\frac{x}{2}\right) dx$

2

- (b) Use Mathematical Induction to prove that

(i) $4(1^3 + 2^3 + 3^3 + \dots + n^3) = n^2(n+1)^2$, for $n = 1, 2, 3, \dots$

3

(ii) Hence find the value of $\lim_{n \rightarrow \infty} \left(\frac{1^3 + 2^3 + 3^3 + \dots + n^3}{n^4} \right)$

1

- (c) (i) Express $\sin x + \sqrt{3} \cos x$ in the form $R \sin(x+\alpha)$
where $R > 0$ and $0 \leq \alpha \leq \frac{\pi}{2}$

2

- (ii) Hence sketch $y = \sin x + \sqrt{3} \cos x$ for $-2\pi \leq x \leq 2\pi$ showing
any x and y intercepts.

2

- (iii) Find the general solution to $\sin x + \sqrt{3} \cos x = \sqrt{2}$

2

Question 4 (Start a new sheet of paper)

Marks

- (a)
- α, β
- and
- γ
- are the roots of the equation
- $x^3 + 2x^2 - 3x + 5 = 0$

- (i) State the values of
- $\alpha + \beta + \gamma$
- ,
- $\alpha\beta + \alpha\gamma + \beta\gamma$

2

- (ii) Find the value of
- $\alpha^2 + \beta^2 + \gamma^2$

2

- (b) If a polynomial
- $P(x)$
- is divided by
- $(x+1)$
- the remainder is 5 and when
- $P(x)$
- is divided by
- $(2x+1)$
- the remainder is 3. Find the remainder when
- $P(x)$
- is divided by
- $(x+1)(2x+1)$
- .

3

- (c) From a point
- S
- the bearings of two points
- P
- and
- Q
- are found to be
- $331^\circ T$
- and
- $011^\circ T$
- respectively. From a point
- F
- , 7 km due north of
- S
- , the bearings of
- P
- and
- Q
- are
- $299^\circ T$
- and
- $020^\circ T$
- respectively.

- (i) Show that
- $PF = \sin 29^\circ \times \frac{7}{\sin 32^\circ}$

2

- (ii) By considering the triangle
- FPQ
- , show that if the distance between
- P
- and
- Q
- is
- d
- metres, then

$$d^2 = 49 \left(\frac{\sin^2 29^\circ}{\sin^2 32^\circ} + \frac{\sin^2 11^\circ}{\sin^2 9^\circ} - 2 \frac{\sin 29^\circ \sin 11^\circ \cos 81^\circ}{\sin 32^\circ \sin 9^\circ} \right)$$

3

Question 5 (Start a new sheet of paper)

Marks

- (a) Consider the function
- $f(x) = \frac{x-1}{x^2}$

- (i) Show that there is only one stationary point and determine its nature

3

- (ii) Determine the point of inflexion.

1

- (iii) What happens to
- $f(x)$
- as
- $x \rightarrow \pm\infty$
- ?

1

- (iv) What happens to
- $f(x)$
- as
- $x \rightarrow 0$
- ?

1

- (v) Sketch the curve showing all its essential features.
-
- (Use at least half a page.)

2

- (b) (i) Prove that
- $\frac{d}{dx} \left(\frac{1}{2} r^2 \right) = \dot{r}$

2

- (ii) An object moving in a straight line has an acceleration given by
- $\ddot{x} = x(8 - 3x)$
- where
- x
- metres is its position relative to a fixed point
- O
- .

At $x = 0$, it has a speed of 4 m/s. Find its speed when it is 1 m on the positive side of O .

2

Question 6 (Start a new sheet of paper)

Marks

- (a) A particle is oscillating in simple harmonic motion such that its displacement x metres from the origin is given by the equation $\frac{d^2x}{dt^2} = -16x$ where t is time in seconds.

- (i) Show that $x = a \cos(4t + \alpha)$ is a solution of motion for this particle. (a and α are constants).

1

- (ii) When $t = 0$, $v = 4$ m/s and $x = 5$ m. Show that the amplitude of the oscillation is $\sqrt{26}$ metres.

2

- (iii) What is the maximum speed of the particle?

1

- (b) $P(2ap, ap^2)$ and $Q(2aq, aq^2)$ are two points on the parabola $x^2 = 4ay$. The tangents at P and Q meet at T which is always on the parabola $x^2 = -4ay$.

- (i) Derive the equation of the tangent at P .

2

- (ii) Hence write down the equation of the tangent at Q .

1

- (iii) Show that T is the point $(a(q+p), apq)$.

1

- (iv) Show that $p^2 + q^2 = -6pq$

1

- (v) Find M , the midpoint of PQ .

1

- (vi) Hence, or otherwise, find the locus of M .

2

Question 7 (Start a new sheet of paper)

Marks

- (a) (i) On the same number plane, sketch the graphs of $y = \cos^{-1} x$ and $y = \sin^{-1}(\frac{x}{2})$. Label the important features.

2

- (ii) Show $y = \cos^{-1} x$ and $y = \sin^{-1}(\frac{x}{2})$ intersect at $x = \frac{2}{\sqrt{5}}$.

2

- (iii) Find the inverse function of $y = \sin^{-1}(\frac{x}{2})$

1

- (iv) Hence or otherwise find the area bounded by the x -axis and the graphs $y = \cos^{-1} x$ and $y = \sin^{-1}(\frac{x}{2})$
(answer correct to 2 decimal places.)

3

- (b) Wheat is the only crop grown on Sandy's property in outback NSW. Per hectare the amount of water, W , in kilolitres, used during irrigation times is given by

$$W = Cg^3 + \frac{D}{g}$$

where g is the amount of grain produced in tonnes per hectare and C and D are positive constants. There is a limited amount of water available for irrigation.

- (i) Show that, for maximum hectares under irrigation, production of grain per hectare, g , is given by

$$g = \left(\frac{D}{2C}\right)^{\frac{1}{4}}$$

2

- (ii) Show that for maximum grain produced on Sandy's property, grain production per hectare needs to be about 59% more than that given in part (i) above.

2

Pythagoras ladies college '07

Q1

a) $\lim_{x \rightarrow 0} \frac{\sin 3x}{x} = 3 \lim_{x \rightarrow 0} \frac{\sin 3x}{3x} = 3$

①

b) B is (1, 3)

②

or

$(-2, 5) \xrightarrow{-3:2} (x, y)$

$$\frac{-4-3x}{-3+2} = 7 \quad \frac{10-3y}{-3+2} = -1$$

$$-4-3x = -7$$

$$-3x = -3$$

$$x = 1$$

$$10-3y = 1$$


$$-3y = -9$$

$$y = 3$$

c) $\frac{x+1}{x-2} = 2 \quad x \neq 2$

$$x+1 = 2x-4$$

$$x = 5$$



$x > 5$ or $x < 2$

d) $y = \tan^{-1} 2x$

$$\frac{dy}{dx} = \frac{1}{1+(2x)^2}$$

$$= \frac{1}{1+4x^2}$$


when $x = \frac{1}{2}$

$$\begin{aligned} m_{\text{tang}} &= \frac{1}{1+4 \cdot \frac{1}{4}} \\ &= \frac{1}{2} \end{aligned}$$

Q1

$$\begin{aligned} \int_0^3 \frac{1}{\sqrt{9-x^2}} dx &= \left[\sin^{-1} \frac{x}{3} \right]_0^3 \\ &= \sin^{-1} \frac{3}{3} - \sin^{-1} 0 \\ &= \frac{\pi}{2}. \end{aligned}$$

f) $y = |2x-1|$ and $y = |x+1|$



$$2x-1 = x+1$$

$$x = 2$$

$$-2x+1 = x+1$$

$$-3x = 0$$

$$x = 0$$

$|2x-1| \leq |x+1|$ when $0 \leq x \leq 2$.

Q2.

$$\frac{\sin 2\theta}{\sin \theta} - \sec \theta = \frac{\cos 2\theta}{\cos \theta}$$

$$\begin{aligned} \text{a) LHS} &= \frac{2 \sin \theta \cos \theta}{\sin \theta} - \frac{1}{\cos \theta} \\ &= 2 \cos \theta - \frac{1}{\cos \theta} \\ &= \frac{2 \cos^2 \theta - 1}{\cos \theta} \\ &= \frac{\cos 2\theta}{\cos \theta}. \end{aligned}$$

$$\begin{aligned} \text{b) } u &= 2t-1 & \text{when } t=1 & u=1 \\ \frac{du}{dt} &= 2 & \text{when } t=\frac{1}{2} & u=0. \end{aligned}$$

$$\begin{aligned} \int_{\frac{1}{2}}^1 4t(2t-1) dt &= \int_0^1 2(u+1)u^{\frac{5}{2}} du \\ &= \int_0^1 u^6 + u^5 du \\ &= \left[\frac{u^7}{7} + \frac{u^6}{6} \right]_0^1 \\ &= \frac{1}{7} + \frac{1}{6} \\ &= \frac{13}{42} \end{aligned}$$

$$\text{c) } \tan \theta = \left| \frac{3-m}{1+3m} \right|$$

$$1 = \frac{3-m}{1+3m}$$

$$1+3m = 3-m$$

$$4m = 2$$

$$m = \frac{1}{2}$$

$$-1 = \frac{3-m}{1+3m}$$

$$-1-3m = 3-m$$

$$-2m = 4$$

$$m = -2$$

$$\text{d) } \tan 2\theta - \cot \theta = 0 \quad 0 \leq \theta \leq \pi \quad (3)$$

$$\frac{2 \tan \theta}{1 - \tan^2 \theta} - \frac{1}{\tan \theta} = 0$$

$$2 \tan^2 \theta = 1 - \tan^2 \theta$$


$$3 \tan^2 \theta = 1$$

$$\tan \theta = \pm \frac{1}{\sqrt{3}}$$

$$\theta = \frac{\pi}{6}, \frac{5\pi}{6}$$

Check $\theta = \frac{\pi}{2}$ $0-0=0$ True

\therefore Solns $\theta = \frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}$



(i) Consider $\triangle PFS$.

$$\frac{PF}{\sin 29} = \frac{7}{\sin 32}$$

$$PF = \sin 29 \times \frac{7}{\sin 32}$$

(ii) Similarly considering $\triangle QFS$

$$QF = \sin 11 \times \frac{7}{\sin 9}$$

Now considering $\triangle PQF$

$$\begin{aligned} PQ^2 &= PF^2 + QF^2 - 2 \cdot PF \cdot QF \cos 51^\circ \\ \text{i.e., } d^2 &= \sin^2 29 \times \frac{49}{\sin^2 32} + \sin^2 11 \times \frac{49}{\sin^2 9} - 2 \times \sin 29 \times \frac{1}{\sin 32} \times \\ &\quad \sin 11 \times \frac{7}{\sin 9} \times \cos 51^\circ \\ &= 49 \left[\frac{\sin^2 29}{\sin^2 32} + \frac{\sin^2 11}{\sin^2 9} - 2 \frac{\sin 29 \sin 11 \cos 51^\circ}{\sin 32 \sin 9} \right] \end{aligned}$$

Katie

Q4

$$x^3 + 2x^2 - 3x + 5 = 0$$

$$\alpha + \beta + \gamma = -\frac{b}{a} = -2$$

$$\alpha\beta + \alpha\gamma + \beta\gamma = \frac{c}{a} = -3$$

$$\begin{aligned} \alpha^2 + \beta^2 + \gamma^2 &= (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \alpha\gamma + \beta\gamma) \\ &= (-2)^2 - 2(-3) \\ &= 4 + 6 \\ &= 10 \end{aligned}$$

1/4 K.P.C.P

$$b) P(x) = (x+2)(x+1) + ax+b$$

$$P(1) = -a+b = 5 \quad \text{①}$$

$$P(-\frac{1}{2}) = -\frac{1}{2}a + b = 3 \quad \text{②}$$

$$\begin{aligned} \text{①-②} \quad -\frac{1}{2}a &= 2 \\ a &= -4 \\ b &= 1 \end{aligned}$$

$$\therefore \text{Remainder} = -4x + 1$$

Kate

Kate is the neatest writer

②

②

③

Q3.


$$\begin{aligned}
 \text{(i)} \lim_{n \rightarrow \infty} \left(\frac{1^3 + 2^3 + 3^3 + \dots + n^3}{n^4} \right) &= \lim_{n \rightarrow \infty} \frac{n^2(n+1)^2}{4n^4} \\
 &= \lim_{n \rightarrow \infty} \frac{n^4 + 2n^3 + n^2}{4n^4} \\
 &= \lim_{n \rightarrow \infty} \frac{1 + \frac{2}{n} + \frac{1}{n^2}}{4} \\
 &= \frac{1}{4}
 \end{aligned}$$

c. let $\sin x + \sqrt{3} \cos x = R \sin(x+\alpha)$

$$\begin{aligned}
 R \cos \alpha &= 1 & \textcircled{1} \\
 R \sin \alpha &= \sqrt{3} & \textcircled{2}
 \end{aligned}$$

$$\begin{aligned}
 \textcircled{2} \quad \tan \alpha &= \sqrt{3} \\
 \textcircled{1} \quad \alpha &= \frac{\pi}{3}
 \end{aligned}
 \qquad R^2 = 4$$

$$\therefore \sin x + \sqrt{3} \cos x = 2 \sin\left(x + \frac{\pi}{3}\right)$$



$$\begin{aligned}
 \text{(ii)} \quad 2 \sin\left(x + \frac{\pi}{3}\right) &= \sqrt{2} \\
 \sin\left(x + \frac{\pi}{3}\right) &= \frac{1}{\sqrt{2}} \\
 x + \frac{\pi}{3} &= n\pi + (-1)^n \frac{\pi}{4} \\
 \therefore x &= n\pi + (-1)^n \frac{\pi}{4} - \frac{\pi}{3}
 \end{aligned}$$

①

$$\text{Q3. } \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \cos^2\left(\frac{x}{2}\right) dx$$

$$\frac{1}{2} \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} (\cos x + 1) dx$$

$$\frac{1}{2} \left[\sin x + x \right]_{\frac{\pi}{6}}^{\frac{\pi}{2}}$$

$$= \frac{1}{2} \left[1 + \frac{\pi}{2} - \left(\frac{\pi}{6} + \frac{\pi}{6} \right) \right]$$

$$= \frac{1}{2} \left[\frac{\pi}{2} + \frac{\pi}{3} \right]$$

$$= \frac{1}{4} + \frac{\pi}{6}$$

②

$$\begin{aligned}
 \text{b) let } S_n &= 4(1^3 + 2^3 + \dots + n^3) \\
 \text{Required to prove } S_n &= n^2(n+1)^2 \\
 \text{For } n=1 \quad \text{LHS} &= 4(1^3) = 4 \\
 \text{RHS} &= 1^2(1+1)^2 = 4 \\
 \text{Statement is true for } n=1.
 \end{aligned}$$

Assume statement is true for $n=k$.

$$S_k = 4(1^3 + 2^3 + \dots + k^3) = k^2(k+1)^2$$

when $n=k+1$

$$\begin{aligned}
 S_{k+1} &= 4(1^3 + 2^3 + \dots + k^3 + (k+1)^3) \\
 &= 4(1^3 + 2^3 + \dots + k^3) + 4(k+1)^3 \\
 &= k^2(k+1)^2 + 4(k+1)^3 \\
 &= (k+1)^2(k^2 + 4(k+1)) \\
 &= (k+1)^2(k+2)^2
 \end{aligned}$$

Thus if it is true for $n=k$ it is true for $n=k+1$
it is true for $n=1$ hence it is true for $n=2$ and so on

②

③

Q6.

$$\frac{d^2x}{dt^2} = -16x$$

(i) $x = a \cos(4t + \alpha)$
 $\dot{x} = -4a \sin(4t + \alpha)$
 $\ddot{x} = -16a \cos(4t + \alpha)$

$$= -16x$$

$\therefore x = a \cos(4t + \alpha)$ is a soln

(ii) When $t=0$ $v=4$

$$4 = -4a \sin \alpha$$

$$\therefore -1 = a \sin \alpha \quad \textcircled{1}$$

when $t=0$ $x=5$

$$5 = a \cos \alpha \quad \textcircled{2}$$

$$\text{i.e. } \textcircled{1}^2 + \textcircled{2}^2 \quad (\sin^2 \alpha + \cos^2 \alpha = 1)$$

$$a^2 = (-1)^2 + (5)^2$$

$$a^2 = 26$$

$$a = \sqrt{26}$$

OR

using $v^2 = \dot{x}(a^2 - x^2)$

$$16 = 16(a^2 - 25)$$

$$1 = a^2 - 25$$

$$a^2 = 26$$

$$a = \sqrt{26}$$

(iii) Max speed occurs when $\sin(4t + \alpha) = 1$.

$$\dot{x} = -4a\sqrt{26} \sin(4t + \alpha)$$

$$= -4\sqrt{26}$$

$$\text{speed}_{\max} = |-4\sqrt{26}|$$

$$= 4\sqrt{26}$$

①

Q6

$$b \quad x^2 = 4ay$$

$$y = \frac{1}{4a} x^2$$

$$(i) \frac{dy}{dx} = \frac{1}{2a} x$$

$$m_{\text{tang}_P} = \frac{1}{2a} \times 2ap$$

 $= p$

Eq° of tangent at P

$$y - ap^2 = p(x - 2ap^2)$$

$$y = px - ap^2 \quad \textcircled{1}$$

②

$$(ii) y = qx - aq^2 \quad \textcircled{2} \quad \textcircled{1}$$

(iii) Solving eqns for tangents simultaneously.

$$0 = (p-q)x - a(p^2 - q^2)$$

$$x = a(p^2 - q^2) = a \frac{(p+q)(p-q)}{p-q}$$

$$x = a(p+q)$$

$$(i) y = ap(p+q) - ap^2$$

$$= apq$$

$\therefore T$ is $(a(p+q), apq)$

①

(iv) T lies on $x^2 = -4ay$

$$\text{i.e. } a^2(p+q)^2 = -4a^2pq$$

$$(p+q)^2 = -4pq$$

$$p^2 + q^2 + 2pq = -4pq$$

$$p^2 + q^2 = -6pq$$

$$(v) M \text{ is } \left(\frac{a(p+q)}{2}, \frac{ap^2 + q^2}{2} \right)$$

$$= \left(a(p+q), \frac{a(p^2 + q^2)}{2} \right)$$

②

①

①

①

①

Q 5

$$f(x) = \frac{x-1}{x^2}$$

$$\begin{aligned} \text{(i)} \quad f'(x) &= \frac{x^2 \cdot 1 - 2x(x-1)}{x^4} \\ &= \frac{x^2 - 2x^2 + 2x}{x^4} \\ &= \frac{x(2-x)}{x^4} \\ &= \frac{2-x}{x^3} \end{aligned}$$

$$f'(x) = 0 \text{ when } x=2.$$

$$\begin{aligned} f''(x) &= \frac{x^3 \cdot 1 - 3x^2(2-x)}{x^4} \\ &= \frac{-x^3 + 6x^2 + 3x^2}{x^4} \\ &= \frac{2x^2(x+3)}{x^4} \\ &= \frac{2(x+3)}{x^4} \end{aligned}$$

$$\text{when } x=2 \quad f''(x) < 0$$

\therefore Only stationary pt $(2, t)$ which is a max

$$\text{(ii)} \quad f''(x) = \frac{2(x+3)}{x^4} = 0 \text{ when } x=3$$

$$\begin{array}{c|c|c|c} x & 3^- & 3 & 3^+ \\ \hline f''(x) & - & 0 & + \end{array}$$

Change in concavity


\therefore There is a pt of inflection at $(3, \frac{2}{9})$

(3)

Q5

$$\begin{aligned} \text{(iii)} \quad \text{As } x \rightarrow \infty \quad f(x) &\rightarrow 0 \\ \text{As } x \rightarrow -\infty \quad f(x) &\rightarrow 0 \end{aligned}$$

$$\text{(iv)} \quad \text{As } x \rightarrow 0 \quad f(x) \rightarrow -\infty$$



(1)

$$\begin{aligned} b) \quad \frac{d}{dx} \left(\frac{1}{2} v^2 \right) &= \frac{d}{dv} \left(\frac{1}{2} v^2 \right) \frac{dv}{dx} \\ &= v \cdot \frac{dv}{dx} \\ &= \frac{dx}{dt} \cdot \frac{dv}{dx} \\ &= \frac{dv}{dt} \\ &= \ddot{x} \end{aligned}$$

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = 8x - 3x^2$$

$$\frac{1}{2} v^2 = \int 8x - 3x^2 dx$$

$$\frac{1}{2} v^2 = 4x^2 - x^3 + C$$

$$\text{when } x=0 \quad v=4 \quad \therefore C=8$$

$$v^2 = 8x^2 - 2x^3 + 16$$

$$\text{when } x=1 \quad v^2 = 8-2+16$$

(1)

(1)

(2)

(2)

(2)

Q6

For M

$$x = a(p+q)$$

$$x^2 = a^2(p^2 + q^2 + 2pq)$$

$$= a^2(-6pq + 2pq)$$

$$x^2 = -4apq \quad \textcircled{1}$$

(2)

$$y = \frac{a}{2}(p^2 + q^2)$$

$$= \frac{a}{2}x - 6pq$$

$$y = -3apq$$

$$\therefore \frac{y}{-3a} = pq$$

Sub in ①


$$x^2 = -4a^2 \times \frac{y}{-3a}$$

$$x^2 = \frac{4}{3}ay$$

$$3x^2 = 4ay$$

Q7

(i)



(ii)


Or

$$(ii) y = \cos^{-1} x$$

$$\text{when } x = \frac{2}{\sqrt{5}}$$


$$y = \cos^{-1} \frac{2}{\sqrt{5}}$$

$$\therefore \cos y = \frac{2}{\sqrt{5}}$$



$$\text{let } \cos^{-1} x = \sin^{-1} \frac{x}{\sqrt{5}} = \alpha.$$

$$\cos \alpha = x \quad \sin \alpha = \frac{x}{\sqrt{5}}$$



$$\text{Consider } y = \sin^{-1} x$$

$$y = \sin^{-1} \left(\frac{2}{\sqrt{5}} \right)$$

$$\therefore \sin y = \frac{1}{\sqrt{5}} \text{ True}$$

$$\therefore \text{Curves intersect at } x = \frac{2}{\sqrt{5}}.$$

$$(iii) y = \sin^{-1} \frac{x}{2}$$

$$\text{Inv. fn is } x^2 \quad \sin y = \frac{x}{2}$$

$$\therefore x = 2 \sin y$$

$$x^2 + \frac{x^2}{4} = 1$$

$$5x^2 = 4$$

$$x^2 = \frac{4}{5}$$

$$\sqrt{\frac{4}{5}}$$

$$x = \frac{2}{\sqrt{5}} \quad x > 0 \text{ from graph}$$

Q7.

$$(iv) y = \cos^{-1} x$$

$$\text{Inv. } x = \cos y.$$

Area required is same as



$$\text{Area} = \int_0^{0.46...} (\cos x - 2 \sin x) dx$$

$$[\sin x + 2 \cos x]_0^{0.46...}$$

$$[0.443... + 1.792... - 0 - 2]$$

$$= 0.24 \text{ units}^2.$$

Q7 b) To maximise hectares irrigated,
we need to minimise Water per hectare, (W)

$$\frac{dW}{dg} = 2Cg - \frac{D}{g^2} \quad \left. \begin{array}{l} \\ \end{array} \right\} \frac{1}{2}$$

$= 0$ for stationary pts.

$$\text{ie } g^3 = \frac{D}{2C} \quad \left. \begin{array}{l} \\ \end{array} \right\} \frac{1}{2}$$

$$g = \left(\frac{D}{2C}\right)^{\frac{1}{3}} \quad (1)$$

$$\frac{d^2W}{dg^2} = 2C + \frac{2D}{g^3} \quad \left. \begin{array}{l} \\ \end{array} \right\} \frac{1}{2}$$

> 0 since $C, D > 0$ \therefore minima T.P. $\left. \begin{array}{l} \\ \end{array} \right\} \frac{1}{2}$

(ii) Let G = tonnes of grain per kL water
We need to maximise G .

$$G = \frac{\text{tonnes of grain per hectare} \times \text{hectares}}{\text{per kL water}} \quad \left. \begin{array}{l} \\ \end{array} \right\} \frac{1}{2}$$

$$= g \times \frac{1}{Cg^2 + \frac{D}{g}}$$

$$\frac{dG}{dg} = \frac{Cg^2 + \frac{D}{g} - g(2Cg - \frac{D}{g^2})}{(Cg^2 + \frac{D}{g})^2} \quad \left. \begin{array}{l} \\ \end{array} \right\} \frac{1}{2}$$

$= 0$ for stationary pts

$$\text{ie } g^3 = \frac{2D}{C}$$

$$g = \left(\frac{2D}{C}\right)^{\frac{1}{3}} \quad (2)$$

$$\frac{dG}{dg} = -\frac{1}{3}\left(Cg^3 - 2D\right) \times \frac{1}{(Cg^2 + \frac{D}{g})^2}$$

$\frac{dG}{dg} < 0$	$\frac{dG}{dg} = 0$	$\frac{dG}{dg} > 0$
$\frac{dG}{dg} = \text{neg. pos.}$	0	$\frac{dG}{dg} = \text{pos. pos. pos. pos.} = \text{neg.}$

\therefore Max T.R.

Now from (1) & (2) above
comparing results

$$\sqrt[3]{\frac{2D}{C}} = \sqrt[3]{4}$$

$$\sqrt[3]{\frac{D}{2C}} = 1.587\dots$$

$\approx 59\%$ more than before