



**SYDNEY BOYS HIGH SCHOOL**  
**MOORE PARK, SURRY HILLS**

**2003**  
**TRIAL HIGHER SCHOOL**  
**CERTIFICATE**

# Mathematics      Extension 1

## General Instructions

- Reading Time – 5 Minutes
- Working time – 3 hours
- Write using black or blue pen. Pencil may be used for diagrams.
- Board approved calculators may be used.
- All necessary working should be shown in every question.

## Total Marks – 84

- Attempt all questions.
- All questions are of equal value.
- Each section is to be answered in a separate bundle, labeled Section A (Questions 1, 2, 3), Section B (Questions 4, 5, 6) and Section C (Questions 7 and 8).

Examiner: *A.M. Gainford*

*Note: This is an assessment task only and does not necessarily reflect the content or format of the Higher School Certificate.*

**Total marks - 84.**

**Attempt Questions 1-7.**

**All questions are of equal value.**

Answer each Section in a SEPARATE writing booklet. Extra writing booklets are available

**Section A** Use a SEPARATE writing booklet

<b>Question 1</b>	(12 marks)	<b>Marks</b>
(a)	Differentiate	
(i)	$x \sin 3x$	1
(ii)	$e^{1-x^2}$	1
(b)	Find the acute angle between the lines $3y = 2x + 8$ and $5x - y - 9 = 0$ .	2
(c)	Evaluate	
(i)	$\int_0^2 \frac{dx}{4+x^2}$	2
(ii)	$\int_0^1 \frac{x^2}{2+x^3} dx$	2
(d)	The letters of the word INTEGRAL are arranged in a row.  If one of these arrangements is selected at random, what is the probability that the vowels are in the same position?	2
(e)	Solve the inequality $\frac{-4}{x} > 0$ .	2

**Section A continued.**

**Question 2.** (12 marks)

**Marks**

(a) If  $\alpha$ ,  $\beta$ , and  $\gamma$  are the roots of the equation  $2x^3 - 5x^2 - 3x + 1 = 0$ , evaluate

(i)  $\alpha + \beta + \gamma$  and  $\alpha\beta\gamma$ .

**1**

(ii)  $\alpha^2 + \beta^2 + \gamma^2$ .

**2**


(b) Use the substitution  $u = x^2 + 4$  to find the exact value of  $\int_0^{2\sqrt{3}} \frac{x}{\sqrt{x^2 + 4}} dx$ .

**3**

(c) Determine the exact value of  $\cos \tan^{-1} \frac{8}{15}$ .

**2**

(d)



The diagram shows the graph of  $y = 2 \sin^{-1} 3x$ .

(i) Find the coordinates of A and C.

**2**

(ii) Find the gradient of the tangent at B.

**2**

**Section A continued.**


**Question 3.** (12 marks)

**Marks**

- (a) A function is defined as  $f(x) = 1 + e^{2x}$ . **2**  
 Find the inverse function  $f^{-1}(x)$  and state the domain and range.

- (b) Consider the quadratic expression  $Q(x) = (5k - 4)x^2 - 6x + (6k + 3)$ , **3**  
 where  $k$  is a constant.  
 Find the values of  $k$  for which  $Q(x) = 0$  has rational roots.

- (c)



$ABCD$  is a common tangent to the two circles.

- (i) Prove that  $ABG = DCH$ . **2**
- (ii) Prove that  $BCG \parallel BCH$ . **2**
- (d) Consider the series  $2^N + 2^{N-1} + 2^{N-2} + \dots + 2^{1-N} + 2^{-N}$ ,  
 where  $N$  is a positive integer.
- (i) Find an expression in terms of  $N$  for the number of terms in the series. **2**
- (ii) Find an expression in terms of  $N$  for the sum of the series. **1**

**Section B** Use a SEPARATE writing booklet.

**Question 4.** (12 marks)

**Marks**

(a) Consider the function  $f(\theta) = \frac{\sin \theta + \sin \frac{\theta}{2}}{1 + \cos \theta + \cos \frac{\theta}{2}}$

- (i) Show that  $f(\theta) = t$  where  $t = \tan \frac{\theta}{2}$ . **3**
- (ii) Write down the general solution of  $f(\theta) = 1$ . **1**

- (a) A certain particle moves along the straight line in accordance with the law:  $t = 2x^2 - 5x + 3$ , where  $x$  is measured in centimetres and  $t$  in seconds.

Initially, the particle is 1.5 centimetres to the right of the origin  $O$ , and moving away from  $O$ .

- (i) Show that the velocity,  $v \text{ cms}^{-1}$ , is given by **1**

$$v = \frac{1}{4x - 5}$$

- (ii) Find an expression for the acceleration,  $a \text{ cms}^{-2}$ , of the particle, in terms of  $x$ . **2**

- (iii) Find the velocity and acceleration of the particle when: **3**

( )  $x = 2 \text{ cm}$

( )  $t = 6 \text{ seconds}$

- (iv) Describe carefully in words the motion of the particle. **2**

**Section B continued.**

**Question 5.**

**Marks**


- a) (i) Prove the identity  $\frac{\cos y - \cos(y + 2)}{2\sin} = \sin(y + )$  **2**
- (ii) Hence prove by mathematical induction that for positive integers  $n$ , **4**  
 $\sin + \sin 3 + \sin 5 + \dots + \sin(2n - 1) = \frac{1 - \cos 2n}{2\sin}.$
- (b) (i) Show that the curve  $y = \frac{x^3 + 4}{x^2}$  has one stationary point and no **2**  
points of inflexion.
- (ii) Write down the equation(s) of any asymptotes. **1**
- (iii) Sketch the curve. **1**
- (iv) Hence, use the graph to find the values of  $k$  for which the **2**  
equation  $x^3 - kx^2 + 4 = 0$  has 3 real roots.

**Section C** Use a SEPARATE writing booklet.

**Question 6.** (12 marks)

**Marks**

The straight line  $y = mx + c$  meets the parabola  $x = 2t, y = t^2$  in real distinct points  $P$  and  $Q$  which correspond respectively to the values  $t = p$  and  $t = q$ .



- (i) Prove that  $pq = -c$ . 2
- (ii) Prove that  $p^2 + q^2 = 4m + 2c$ . 2
- (iii) Show that the equation of the normal to the parabola at  $P$  is  $x + py = 2p + p^3$ . 2
- (iv) The point  $N$  is the point of intersection of the normals to the parabola at  $P$  and  $Q$ . 2  
 Show that the coordinates at  $N$  are  $(-pq(p + q), (2 + p^2 + pq + q^2))$
- (v) If the chord  $PQ$  is free to move while maintaining a fixed gradient.
- ( ) Show that the locus of  $N$  is a straight line. 2
- ( ) Hence, or otherwise, show that this straight line is a normal to the parabola. 2

**Section C continued.**

**Question 7.** (12 marks)

**Marks**

- (a) When the polynomial  $P(x)$  is divided by  $(x + 4)$  the remainder is 5 and when  $P(x)$  is divided by  $(x - 1)$  the remainder is 9. Find the remainder when  $P(x)$  is divided by  $(x - 1)(x + 4)$ . **3**

- (b) A projectile is fired from a point on horizontal ground with initial speed  $V \text{ ms}^{-1}$  and angle of projection  $\theta$ . The cartesian equation of the path is given by


$$y = x \tan \theta - \frac{gx^2}{2V^2 \cos^2 \theta}$$

where  $x$  and  $y$  are the horizontal and vertical displacements of the particle from  $O$ , the point of projection.

The acceleration due to gravity is  $g$  and air resistance has been neglected.

- (i) Use the given equation to show that the maximum range  $R$  on the horizontal plane is given by  $R = \frac{V^2}{g}$ . **2**

- (ii) Show that to hit a target  $h$  metres above the ground at the same horizontal distance  $R$  using the same angle of projection  $\theta$ , the speed of projection must be increased to  $\frac{V^2}{\sqrt{V^2 - gh}}$ . **4**




Section C continued.

Question 7.

Marks

(c)



In the triangle  $ABC$ ,  $\angle BAC = 90^\circ$ .  $AD$  bisects  $\angle BAC$ .  
 $DH \perp AB$  and  $DK \perp AC$ .

Copy the diagram.

- (i) Show that  $\frac{AD}{DH} = \sqrt{2}$ . 1
- (ii) By considering the areas of the triangles or otherwise, 2  
show that  $\frac{\sqrt{2}}{AD} = \frac{1}{AB} + \frac{1}{AC}$

**THIS IS THE END OF THE PAPER**



**SYDNEY BOYS HIGH SCHOOL**  
MOORE PARK, SURRY HILLS

**2003**

**TRIAL HIGHER SCHOOL  
CERTIFICATE EXAMINATION**

**Mathematics    Extension 1**

**Sample Solutions**



QUESTION 3

(a)  $y = f(x) \quad y = 1 + e^{2x}$   
 $f^{-1}(x) \quad x = 1 + e^{2y}$   
 $e^{2y} = x - 1$   
 $2y = \log(x - 1)$   
 $y = \frac{1}{2} \log(x - 1)$

Domain  $x > 1$  Range All real  $y$

2

(b) Rational roots when  $\Delta = b^2 - 4ac = 0$  or has rational square root

$$36 - 4(5k - 4)(6k + 3) = 0$$

$$36 - 120k^2 + 36k + 48 = 0$$

$$-120k^2 + 36k + 84 = 0$$

$$10k^2 - 3k - 7 = 0$$

$$(10k + 7)(k - 1) = 0$$

rational roots when  $k = -\frac{7}{10}$  or 1

3

multiple solutions when  $-120k^2 + 36k + 84$  has rational roots

(c) (i)  $\angle ABC = \angle BEG$  (angle in alternate segment)

$\angle BEG = \angle CEH$  (vertically opposite)

$\angle CEH = \angle DCH$  (angle in alternate segment)

$\therefore \angle ABC = \angle DCH$  as required

2

(ii)  $\angle CBH = \angle BGC$  (alternate segment)

$\angle BCE = \angle CHF$  "

$\therefore \angle BGC = \angle HCB$  (angle sum of  $\Delta$ )

2

$\therefore \triangle BGC \cong \triangle BCH$  (equiangular)

(d) (i)  $a = 2^N \quad r = 2^{-1}$

$$S_n = \frac{a}{1-r}$$

$$\frac{a r^{n-1}}{h} = 2^{-N}$$

$$2^N (2^{-1})^{n-1} = 2^{-N}$$

$$2^{-n+1} = 2^{-2N}$$

$$-n+1 = -2N$$

$$n = 2N+1$$

2

$$= \frac{2^N}{1 - \frac{1}{2}}$$

$$= 2 \cdot 2^N = 2^{N+1}$$

1

$$(4) (a) \sin 2\theta = 2 \sin \theta \cos \theta$$

$$(i) \sin \theta = 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}$$

$$\cos 2\theta = \frac{\cos^2 \theta - \sin^2 \theta}{1 - 2 \sin^2 \theta}$$

$$2 \cos^2 \theta - 1$$

$$\text{so } \cos \theta = \frac{\cos^2 \frac{\theta}{2} - \sin^2 \frac{\theta}{2}}{1 - 2 \sin^2 \frac{\theta}{2}}$$

$$2 \cos^2 \frac{\theta}{2} - 1$$

$$f(\theta) = \frac{2 \sin \frac{\theta}{2} \cos \frac{\theta}{2} + \sin \frac{\theta}{2}}{1 + 2 \cos^2 \frac{\theta}{2} - 1 + \cos \frac{\theta}{2}}$$

$$= \frac{\sin \frac{\theta}{2} [2 \cos \frac{\theta}{2} + 1]}{\cos \frac{\theta}{2} [2 \cos \frac{\theta}{2} + 1]}$$

$$= \tan \frac{\theta}{2} = t \quad (3)$$

$$(ii) f(\theta) = \tan \frac{\theta}{2} = 1 \text{ general soln.}$$

$$\text{If } \tan \theta = a, \text{ then } \theta = n\pi + \tan^{-1}(a)$$

$$\tan \frac{\theta}{2} = 1, \text{ then } \frac{\theta}{2} = n\pi + \frac{\pi}{4}$$

$$\theta = 2n\pi + \frac{\pi}{2} \quad (1)$$

4 (b) (i)  $t = 2x^2 - 5x + 3$

$$\frac{dt}{dx} = 4x - 5$$

$$\frac{ds}{dt} = v = \frac{1}{4x-5} \quad (1)$$

(ii) using  $\ddot{x} = \frac{d}{dx} \left( \frac{1}{2} v^2 \right)$

$$= \frac{d}{dx} \left( \frac{1}{2(4x-5)^2} \right)$$

$$= \frac{d}{dx} \left[ \frac{1}{2} (4x-5)^{-2} \right]$$

$$= -(4x-5)^{-3} \times 4$$

$$= \frac{-4}{(4x-5)^3} \quad (2)$$

(iii) (a) when  $x=2$ ,  $v = \frac{1}{3} \text{ cm/s} \quad (2)$

$$a = -\frac{4}{27} \text{ cm/s}^2 \quad (2)$$

(b) when  $t=6$ ,  $6 = 2x^2 - 5x + 3$

$$(2x+1)(x-3) = 0$$

$$x = -\frac{1}{2}, x = 3$$

(1)

take  $x = 3$

at  $x=3$ ,  $v = \frac{1}{7} \text{ cm/s} \quad (1)$

$$a = \frac{4}{343} \quad (1)$$

(iv) particle is travelling to the right but is slowing down

(2)

$$(5) \quad (a) \quad (i) \quad \frac{\cos y - \cos(y+2\alpha)}{2 \sin \alpha} = \sin(y+\alpha)$$

$$\text{LHS} = \frac{\cos y - (\cos y \cos 2\alpha - \sin y \sin 2\alpha)}{2 \sin \alpha}$$

$$\frac{\cos y - (\cos y (1 - 2 \sin^2 \alpha) - \sin y \cdot 2 \sin \alpha \cos \alpha)}{2 \sin \alpha}$$

$$\frac{\cancel{\cos y} - \cancel{\cos y} + \cancel{2 \sin^2 \alpha} \cos y + \cancel{2 \sin \alpha} \cos \alpha \sin y}{\cancel{2 \sin \alpha}}$$

$$= \sin \alpha \cos y + \cos \alpha \sin y$$

$$= \sin(y+\alpha) \quad (2)$$

$$(ii) \quad \sin \alpha + \sin 3\alpha + \sin 5\alpha + \dots + \sin(2n-1)\alpha = \frac{1 - \cos 2n\alpha}{2 \sin \alpha}$$

step 1 Prove true for  $n=1$

$$\text{LHS} = \sin \alpha$$

$$\text{RHS} = \frac{1 - \cos 2\alpha}{2 \sin \alpha} = \frac{1 - (1 - 2 \sin^2 \alpha)}{2 \sin \alpha} = \sin \alpha = \text{LHS}$$

true for  $n=1$

step 2 Assume true for  $n=k$  (a positive integer) so

$$\sin \alpha + \sin 3\alpha + \sin 5\alpha + \dots + \sin(2k-1)\alpha = \frac{1 - \cos 2k\alpha}{2 \sin \alpha}$$

and we must prove it true for  $n=k+1$ , so

$$\sin \alpha + \sin 3\alpha + \sin 5\alpha + \dots + \sin(2k-1)\alpha + \sin(2k+1)\alpha = \frac{1 - \cos 2(k+1)\alpha}{2 \sin \alpha}$$

$$\text{LHS} = \frac{1 - \cos 2k\alpha}{2 \sin \alpha} + \sin(2k+1)\alpha$$

$$\frac{1 - \cos 2kd}{2 \sin d} + \sin(2kd + d).$$

now using (a)(i)  $\sin(y+d) = \frac{\cos y - \cos(y+2d)}{2 \sin d}$

$$\text{then } \sin(2kd+d) = \frac{\cos 2kd - \cos(2kd+2d)}{2 \sin d}$$

$$\text{now, } \frac{1 - \cos 2kd}{2 \sin d} + \frac{\cos 2kd - \cos 2(k+1)d}{2 \sin d}$$

$$= \frac{1 - \cos 2(k+1)d}{2 \sin d}$$

$\Rightarrow$  RHS =

True for  $n = k+1$ .

step 3 If the statement is true for  $n=k$ , then it is also true for  $n=k+1$ . Since the statement is true for  $n=1$ , it follows that it must also be true for  $n=2$  and so on.  $\therefore$  the statement is true for all positive integers  $n$ .

(4)

(5) (b) (i)  $y = \frac{x^3+4}{x^2} = \frac{x^3}{x^2} + \frac{4}{x^2} = x + 4x^{-2} = x + \frac{4}{x^2}$

$y' = 1 - 8x^{-3} = 1 - \frac{8}{x^3}$

$y'' = 24x^{-4} = \frac{24}{x^4}$

Stat points exist when  $y' = 0$ ,  $1 - \frac{8}{x^3} = 0$

$\frac{8}{x^3} = 1 \Rightarrow x^3 = 8$   
 $\Rightarrow x = 2$


At  $x=2$ ,  $y = 2 + \frac{4}{2^2} = 3$  (2, 3) (1) (min stat pt)  
 $y'' > 0$

Inflexions occur when  $y'' = 0$  and  $\exists$  a sign change

$\frac{24}{x^4} = 0 \Rightarrow 24 = 0x^4$   
 does not exist. (1)

(ii)  $y = \frac{x^3+4}{x^2} \Rightarrow x \neq 0$  (y axis) (1/2) vertical asymptote

$y = \frac{x^3(1 + \frac{4}{x^3})}{x^2} = x(1 + \frac{4}{x^3})$  and as  $x \rightarrow \infty$   $y \rightarrow x$ . (1/2) oblique asymptote



when  $y=0$ ,  
 $0 = \frac{x^3+4}{x^2}$

so  $x^3+4 = 0$   
 $x^3 = -4$   
 $x = \sqrt[3]{-4}$

$$5 \text{ (b) (iv)} \quad x^3 - kx^2 + 4 = 0$$

$$x^3 + 4 = kx^2$$

$$\text{So} \quad \frac{x^3 + 4}{x^2} = k.$$

$$\Rightarrow y = \frac{x^3 + 4}{x^2} = k$$

3 intersections will occur between  $y=k$  and  $y = \frac{x^3 + 4}{x^2}$  if


$$k > 3.$$

(2)

Solution — Section C

Question (6) [12]

$$x = 2t, y = t^2 \therefore y = \frac{x^2}{4}$$



Solve:

$$\begin{cases} y = \frac{x^2}{4} \\ y = mx + c \end{cases}$$

$$\therefore x^2 - 4mx - 4c = 0 \quad \text{--- (1)}$$

The roots to (1) are:  $2p, 2q$ .

$$\therefore \sum d_i : 2p + 2q = 4m$$

$$\text{i.e. } p + q = 2m \quad \text{--- (2)}$$

$$\text{Product of roots} : 4pq = -4c$$

$$(i) \therefore pq = -c \quad \text{--- (3) [2]}$$

$$(ii) \text{ Now, } p^2 + q^2 = (p+q)^2 - 2pq \\ = 4m^2 - 2(-c)$$

$$\therefore p^2 + q^2 = 4m^2 + 2c \quad \text{--- [2]}$$

$$(iii) \text{ gradient of } \text{tgt.} = p$$

$$\therefore \text{gradient of normal} = -\frac{1}{p}$$

$$\therefore \text{equation of normal: } y - p^2 = -\frac{1}{p}(x - 2p)$$

$$\therefore x + py = p^3 + 2q$$

(iv) The equation of normal at Q

$$x + qy = q^3 + 2q \quad \text{--- (5)}$$

$\therefore$  (4) - (5) we have:

$$(p-q)y = (p^3 - q^3) + 2(p-q)$$

$$\therefore y = 2 + p^2 + pq + q^2 \quad \text{--- (6)}$$

Substitute (6) into (4) we have:

$$x + 2p + p^3 + p^2q + pq^2 = p^3 + 2p$$

$$\therefore x = -p^2q - pq^2 = -pq(p+q)$$

$$\therefore N(-pq(p+q), (2 + p^2 + pq + q^2))$$

[2].

(8)

Question (6).

$$(V). \quad \begin{cases} pq = -c, & p+q = 2m \\ p^2+q^2 = 4m^2+2c. \end{cases}$$

The x-coord. of N becomes  $c(2m)$

The y-coord. of N becomes  $\sum 2+(4m^2+2c)-c^2$

$$\therefore N = (2mc, 4m^2+c+2)$$

(x) Chord PQ, whose equation is  $y = mx + c$ , is free to move whilst maintaining a fixed grad. i.e.  $m_{PQ} = m$  (a constant), but  $c$  is a variable.

$$\text{Now } x = 2mc, \Rightarrow c = \frac{x}{2m}$$

$$y = 2 + 4m^2 + \frac{x}{2m} \quad [2]$$

$$\therefore y = \frac{x}{2m} + 2(1+2m^2)$$

i.e. Equation of locus of N is a straight line with gradient  $\frac{1}{2m}$  and y-intercept  $2(1+m^2)$ .

The points of intersection of the locus of N and  $x^2 = 4y$  are found by solving

$$\begin{cases} y = \frac{x}{2m} + 2(1+2m^2) \\ x = 2t, \quad y = t^2 \end{cases}$$

$$\text{i.e. } t^2 = \frac{2t}{2m} + 2(1+2m^2) \quad (*)$$

$$mt^2 - t - 2m(1+2m^2) = 0$$

$$\therefore t = \frac{1 \pm \sqrt{1+8m^2(1+2m^2)}}{2m} = \frac{1 \pm (1+4m^2)}{2m}$$

$$\therefore t = \frac{1+2m^2}{m}, \quad \text{or } t = -2m$$

$\therefore$  locus of N cut parabola in 2 pt say U, V with parameters

$$t = \begin{cases} \frac{1+2m^2}{m} \\ -2m \end{cases} \quad [2]$$

From gradient of  $tg t$ , ( $=t$ )  $\Rightarrow$  the gradients of  $tg t$ s at U, V are

$$\frac{1+2m^2}{m}, \quad \text{and } -2m. \quad \text{In particular,}$$

the  $tg t$  at V has gradient  $-2m$  while the locus of N has gradient  $\frac{1}{2m}$ . Hence the locus of N is perp to  $tg t$  at V  $\Rightarrow$  normal at V

Question (7). [12]

(a)  $P(x) = (x+4)u(x) + 5$  [3]  
 $= (x-1)v(x) + 9.$

$\therefore P(-4) = 5, P(1) = 9. \quad \text{--- (1)}$

$P(x) = (x-1)(x+4)q(x) + (ax+b).$

From (1)  $\begin{cases} -4a+b = 5 \\ a+b = 9 \end{cases} \quad \text{--- (2)}$

$\therefore 5a = 4 \Rightarrow a = 4/5$

$\therefore b = 9 - 4/5 = 41/5$

i.e.  $\boxed{\frac{4x}{5} + \frac{41}{5}}$


(b) To find the range, set  $y=0.$

(i) i.e.  $x \left( \tan \theta - \frac{gx}{2v^2 \cos^2 \theta} \right) = 0.$

i.e.  $x = 0, \text{ or } x = \frac{2v^2 \cos^2 \theta \times \tan \theta}{g}$

i.e.  $\text{range} = \frac{v^2 (2 \sin \theta \cos \theta)}{g}$   
 $= \frac{v^2 \sin 2\theta}{g} \quad [2]$

$\therefore$  Maximum range occurs when  $\sin 2\theta = 1 \Rightarrow \boxed{R = \frac{v^2}{g}}$



Equation of higher trajectory ( $T_2$ ) is

(1)  $\boxed{h = R \tan \theta - \frac{gR^2}{2U^2 \cos^2 \theta}} \quad (\text{velocity})$

When the speed of projectile was  $V$ , the range was:

$\boxed{R = \frac{V^2 \sin 2\theta}{g}} \quad \text{--- (2)}$

Substitute (2) into (1) we have.

$h = \frac{V^2 \sin 2\theta \tan \theta}{g} - \frac{V^4 \sin^2 2\theta}{g U^2 \cos^2 \theta}$

Note:  $\sin 2\theta = 2 \sin \theta \cos \theta, \tan \theta = \frac{\sin \theta}{\cos \theta}$

$\therefore h = \frac{2V^2 \sin^2 \theta}{g} - \frac{2V^4 \sin^2 \theta}{g U^2}$


When  $v = V$ , Range is  $R_{\max} \therefore \theta = 45^\circ$

$\therefore h = \frac{V^2}{g} - \frac{V^4}{g U^2} \quad [4]$

$\therefore U^2 g h = V^2 U^2 - V^4, U^2 (v^2 - gh) =$

$\therefore U^2 = \frac{V^4}{v^2 - gh} \therefore \boxed{U = \frac{V^2}{\sqrt{v^2 - gh}}}$

∴ Question  
7(c)



∴ AD bisects  $\angle BAC (=90^\circ)$

∴  $\angle BAD = \angle DAC = 45^\circ$

⇒  $\angle HDA = \angle KDA = 45^\circ$

(Angle sum of a  $\Delta$ ).

i.e.  $\Delta AHD$  is isos. ⇒  $AH = DH$ .

In  $\Delta AHD$ ,  $AD^2 = AH^2 + DH^2$

(Pythagoras)  $= 2DH^2$

$$\therefore \left(\frac{AD}{DH}\right)^2 = 2$$

(i) ⇒  $\frac{AD}{DH} = \sqrt{2}$  [1]

$$\therefore \boxed{\frac{1}{DH} = \frac{\sqrt{2}}{AD}} \quad \text{--- (1)}$$

(ii)

$\Delta AHD \cong \Delta AKD$ . (AAS).

∴  $DH = DK$ . --- (2)

Area of  $\Delta ABC$

$$= \frac{1}{2} AB \cdot AC.$$

but area of  $\Delta ABC$

$$= \text{area of } \Delta ABD + \text{area of } \Delta ACD.$$

$$\text{Area of } \Delta ABD = \frac{1}{2} AB \cdot DH$$

$$\text{Area of } \Delta ACD = \frac{1}{2} AC \cdot DK.$$

from (2) ∴  $DK = DH$

$$\therefore \text{area of } \Delta ACD = \frac{1}{2} AC \cdot DH.$$

$$\therefore \frac{1}{2} AB \cdot AC = \frac{1}{2} (AB \cdot DH + AC \cdot DH)$$

$$\therefore DH (AB + AC) = AB \cdot AC. \quad [2]$$

$$\therefore \frac{AB + AC}{AB \cdot AC} = \frac{1}{DH}$$

$$\therefore \boxed{\frac{1}{AC} + \frac{1}{AB} = \frac{\sqrt{2}}{AD}} \quad \text{from (1)}$$