

# **PENRITH HIGH SCHOOL**



## **MATHEMATICS EXTENSION 2 2012**

### **HSC Trial**

**Assessor: Mr Ferguson**

**General Instructions:**

- Reading time – 5 minutes
- Working time – **3 hours**
- Write using black or blue pen. Black pen is preferred
- Board-approved calculators may be used.
- A table of standard integrals is provided at the back of this paper.
- A multiple choice answer sheet is provided at the back of this paper.
- Show all necessary working in Questions 11 – 16.
- Work on this question paper will not be marked.

#### **Section 1**

**Total marks – 100**

**SECTION 1 – Pages 2 – 5**

**10 marks**

- Attempt Questions 1 – 10
- Allow about 15 minutes for this section.

**SECTION 2 – Pages 6 – 12**

**90 marks**

- Attempt Questions 11 – 16
- Allow about 2 hours 45 minutes for this section.

#### **Section 2**

<b>Question</b>	<b>Mark</b>
1	
2	
3	
4	
5	

<b>Question</b>	<b>Mark</b>
6	
7	
8	
9	
10	
<b>Total</b>	<b>/10</b>

<b>Question</b>	<b>Mark</b>
11	/15
12	/15
13	/15
14	/15
15	/15
16	/15


<b>Total</b>	<b>/100</b>
%	

**This paper MUST NOT be removed from the examination room**

***Student Number: .....***


**SECTION 1:** Circle the correct answer on the multiple choice answer sheet

- 1 The diagram shows the graph of the function  $y = f(x)$ .




Which of the following is the graph of  $y = |f(x)|$ ?


(A)




(B)



(C)




(D)



**2** Let  $z = 4 + i$ . What is the value of  $i\bar{z}$ ?

- (A)  $-1 - 4i$
- (B)  $-1 + 4i$
- (C)  $1 - 4i$
- (D)  $1 + 4i$

**3** Consider the Argand diagram below.



Which inequality could define the shaded area?

- |  |  |
|--|--|
| <p>(A) <math> z - i  \leq 2</math> and <math>0 \leq \arg(z - 1) \leq \frac{3\pi}{4}</math></p> | <p>(B) <math> z + i  \leq 2</math> and <math>0 \leq \arg(z - 1) \leq \frac{3\pi}{4}</math></p> |
| <p>(C) <math> z - i  \leq 2</math> and <math>0 \leq \arg(z - 1) \leq \frac{\pi}{4}</math></p>  | <p>(D) <math> z + i  \leq 2</math> and <math>0 \leq \arg(z - 1) \leq \frac{\pi}{4}</math></p>  |

**4** Consider the hyperbola with the equation  $\frac{x^2}{9} - \frac{y^2}{5} = 1$ .

What are the coordinates of the vertex of the hyperbola?

- |                                    |                                    |
|------------------------------------|------------------------------------|
| <p>(A) <math>(\pm 3, 0)</math></p> | <p>(B) <math>(0, \pm 3)</math></p> |
| <p>(C) <math>(0, \pm 9)</math></p> | <p>(D) <math>(\pm 9, 0)</math></p> |

**5** The points  $P(cp, \frac{c}{p})$  and  $Q(cq, \frac{c}{q})$  lie on the same branch of the hyperbola  $xy = c^2$  ( $p \neq q$ ). The tangents at  $P$  and  $Q$  meet at the point  $T$ . What is the equation of the normal to the hyperbola at  $P$ ?

- (A)  $p^2x - py + c - cp^4 = 0$
- (B)  $p^3x - py + c - cp^4 = 0$
- (C)  $x + p^2y - 2c = 0$
- (D)  $x + p^2y - 2cp = 0$

**6** What is the value of  $\int \sec x dx$ ? Use the substitution  $t = \tan \frac{x}{2}$ .

- (A)  $\ln |(t+1)(t-1)| + c$       (B)  $\ln \left| \frac{1+t}{1-t} \right| + c$   
(C)  $\ln |(1+t)(1-t)| + c$       (D)  $\ln \left| \frac{t+1}{t-1} \right| + c$

**7** Let  $I_n = \int_0^x \sin^n t dt$ , where  $0 \leq x \leq \frac{\pi}{2}$ .

Which of the following is the correct expression for  $I_n$ ?

- (A)  $I_n = \left( \frac{n-1}{n} \right) I_{n-2}$  with  $n \geq 2$ .  
(B)  $I_n = \left( \frac{n+1}{n} \right) I_{n-2}$  with  $n \geq 2$ .  
(C)  $I_n = n(n-1)I_{n-2}$  with  $n \geq 2$ .  
(D)  $I_n = n(n+1)I_{n-2}$  with  $n \geq 2$ .

**8** The region enclosed by  $y = x^3$ ,  $y = 0$  and  $x = 2$  is rotated around the y-axis to produce a solid. What is the volume of this solid?

- (A)  $\frac{8\pi}{5}$  units<sup>3</sup>  
(B)  $\frac{32\pi}{5}$  units<sup>3</sup>  
(C)  $\frac{64\pi}{5}$  units<sup>3</sup>  
(D)  $\frac{16\pi}{5}$  units<sup>3</sup>

**9** What is the angle at which a road must be banked so that a car may round a curve with a radius of 100 metres at 90 km/h without sliding? Assume that the road is smooth and gravity to be  $9.8 \text{ ms}^{-2}$ .

- (A)  $83^\circ 10'$       (B)  $32^\circ 32'$   
(C)  $83^\circ 6'$       (D)  $32^\circ 53'$

**10** The polynomial equation  $x^3 + 4x^2 - 2x - 5 = 0$  has roots  $\alpha$ ,  $\beta$  and  $\gamma$ . Which of the following polynomial equations have roots  $\alpha^2$ ,  $\beta^2$  and  $\gamma^2$ ?

- (A)  $x^3 - 20x^2 - 44x - 25 = 0$
- (B)  $x^3 - 20x^2 + 44x - 25 = 0$
- (C)  $x^3 - 4x^2 + 5x - 1 = 0$
- (D)  $x^3 + 4x^2 + 5x - 1 = 0$

## SECTION 2

**Question 11** (15 marks) (Use a new page to write your answers)

(a) Find (i)  $\int \frac{t^2 - 1}{t^3} dt$ . 4

(ii)  $\int \frac{dx}{\sqrt{6-x-x^2}}$

(b) Evaluate (i)  $\int_0^1 \frac{x}{(x+1)(2x+1)} dx$  3

(ii)  $\int_0^{\frac{\pi}{4}} x \tan^2 x dx$  3

(c) (i) If  $I_n = \int_0^{\frac{\pi}{2}} x^n \cos x dx$ , show that for  $n > 1$  , 3

$$I_n = \left(\frac{\pi}{2}\right)^n - n(n-1)I_{n-2}$$

(ii) Hence find the area of the finite region bounded by the curve 2

$$y = x^4 \cos x \text{ and the } x \text{ axis for } 0 \leq x \leq \frac{\pi}{2} .$$

**Question 12** (15 marks) (Use a new page to write your answers)

(a) Given that  $z = \sqrt{2} - \sqrt{2}i$  and  $w = -\sqrt{2}$ , find, in the form  $x + iy$ :

(i)  $wz^2$  1

(ii)  $\arg z$  1

(iii)  $\frac{z}{z+w}$  2

(iv)  $|z|$  1

(v)  $z^{10}$  2

(b) Find the values of real numbers  $a$  and  $b$  such that  $(a+ib)^2 = 5-12i$  2

(c) Draw Argand diagrams to represent the following regions 2

(i)  $1 \leq |z+4-3i| \leq 3$


(ii)  $\frac{\pi}{6} \leq \arg z \leq \frac{\pi}{3}$

(d) (i) Show that  $\frac{1+\cos\theta+i\sin\theta}{1-\cos\theta-i\sin\theta} = i \cot \frac{\theta}{2}$  2

(ii) Hence solve  $\left(\frac{z-1}{z+1}\right)^8 = -1$  2

**Question 13** (15 marks) (Use a new page to write your answers)

- (a) The diagram shows the graph of the function  $f(x) = x^2 + x - 2$ . On separate diagrams sketch the following graphs, showing clearly any intercepts on the coordinate axes and the equations of any asymptotes.




(i)  $y = |f(x)|$  1

(ii)  $y = [f(x)]^2$  1

(iii)  $y = \frac{1}{f(x)}$  2

(iv)  $y = \log_e f(x)$  2


- (b) The horizontal base of a solid is the area enclosed by the curve  $|x|^{\frac{1}{2}} + |y|^{\frac{1}{2}} = 1$ .  
 Vertical cross sections taken perpendicular to the  $x$ -axis are equilateral triangles with one side in the base.



- (i) Show that the volume of the solid is given by  $V = 2\sqrt{3} \int_0^1 (1-\sqrt{x})^4 dx$  2
- (ii) Use the substitution of  $u = 1 - \sqrt{x}$  to evaluate this integral. 3

- (c) The tangent  $AE$  is parallel to the chord  $DC$ .

- (i) Prove that  $(AB)^2 = BC \cdot BE$  3
- (ii) Hence or otherwise prove that  $\frac{AC}{AE} = \sqrt{\frac{BC}{BE}}$  1



**Question 14** (15 marks) (Use a new page to write your answers)

- (a) The equation of an ellipse is given by  $4x^2 + 9y^2 = 36$ .
- (i) Find  $S$  and  $S'$  the foci of the ellipse 2
- (ii) Find the equations of the directrices  $M$  and  $M'$  1
- (iii) Sketch the ellipse showing foci, directrices and axial intercepts. 2
- (iv) Let  $P$  be any point on the ellipse.  
Show  $SP + S'P = 6$  2
- (v) Find the equation of the chord of contact from an external point  $(3, 2)$  1
- 
- (b) (i) Sketch the rectangular hyperbola  $xy = c^2$ , labelling the point  $P\left(ct, \frac{c}{t}\right)$  on it. 1
- (ii) Show that the equations of the tangent and normal to the hyperbola at  $P$  are  $x + t^2 y = 2ct$  and  $ty + ct^4 = t^3 x + c$  respectively. 3
- (iii) If the tangent at  $P$  meets the coordinate axes at  $X$  and  $Y$  respectively and the normal at  $P$  meets the lines  $y = x$  and  $y = -x$  at  $R$  and  $S$  respectively, prove that the quadrilateral  $RYSX$  is a rhombus. 3

**Question 15** (15 marks) (Use a new page to write your answers)

- (a) When a certain polynomial is divided by  $x+1$ ,  $x-3$  the respective remainders are 6 and -2. Find the remainder when this polynomial is divided by  $x^2 - 2x - 3$ . 3

- (b) The cubic equation  $x^3 + px + q = 0$  has 3 non-zero roots  $\alpha, \beta, \gamma$ . 3

Find, in terms of the constants  $p, q$  the values of

(i)  $\alpha^2 + \beta^2 + \gamma^2$

(ii)  $\alpha^3 + \beta^3 + \gamma^3$ .

- (c) If  $\alpha, \beta, \gamma$  are the roots of the equation  $3x^3 - 5x^2 - 4x + 3 = 0$ , 3  
find the cubic equation with roots  $\alpha - 1, \beta - 1, \gamma - 1$ .

- (d) A polynomial of degree  $n$  is given by  $P(x) = x^n + ax - b$ . It is given that the polynomial has a double root at  $x = \alpha$ .

- (i) Find the derived polynomial  $P'(x)$  and show that  $\alpha^{n-1} = -\frac{a}{n}$ . 3

- (ii) Show that  $\left(\frac{a}{n}\right)^n + \left(\frac{b}{n-1}\right)^{n-1} = 0$ . 2

- (iii) Hence deduce that the double root is  $\frac{bn}{a(n-1)}$ . 1

**Question 16** (15 marks) (Use a new page to write your answers)

- (a) For  $a > 0, b > 0, c > 0$  and  $d > 0$  and given that  $\frac{a+b}{2} \geq \sqrt{ab}$ , show that 2


$$\frac{a+b+c+d}{4} \geq \sqrt[4]{abcd}$$

- (b) (i) Use De Moivre's theorem to express  $\tan 5\theta$  in terms of powers of  $\tan \theta$ . 3

- (ii) Hence show that  $x^4 - 10x^2 + 5 = 0$  has roots  $\pm \tan \frac{\pi}{5}$  and  $\pm \tan \frac{2\pi}{5}$ . 2

- (iii) Deduce that  $\tan \frac{\pi}{5} \cdot \tan \frac{2\pi}{5} \cdot \tan \frac{3\pi}{5} \cdot \tan \frac{4\pi}{5} = 5$  1

- (c) A mass 10 kg, centre  $B$  is connected by light rods to sleeves  $A$  and  $C$  which revolve freely about the vertical axis  $AC$  but do not move vertically.



- (i) Given  $AC = 2$  metres, show that the radius of the circular path of rotation of  $B$  is  $\frac{\sqrt{3}}{2}$  metres. 1

- (ii) Find the tensions in the rods  $AB, BC$  when the mass makes 90 revolutions per minute about the vertical axis. 3

- (d) Given that  $a_n = \sqrt{2 + a_{n-1}}$  for integers  $n \geq 1$  and  $a_0 = 1$ , by mathematical induction prove that for  $n \geq 1$  :

$$\sqrt{2} < a_n < 2$$

## Section 1

1) B.

$$\begin{aligned} 2) \quad iz &= i(4+i) \\ &= 4i + i^2 \\ &= 4i - 1 \\ &= -1 + 4i \\ \bar{iz} &= -1 - 4i \\ &= A. \end{aligned}$$

3) A.

$$\begin{aligned} 4) \quad \text{Let } y &= 0 \\ \frac{x^2}{9} &= 1 \\ x^2 &= 9 \\ x &= \pm 3. \end{aligned}$$

$(\pm 3, 0)$

A

$$\begin{aligned} 5) \quad \text{Normal and } P(cp, \frac{c}{p}) \\ \therefore y - \frac{c}{p} &= p^2(x - cp) \\ py - c &= p^3(x - cp) \\ py - c &= p^3x - p^4c \\ p^3x - py + c - cp^4 &= 0 \end{aligned}$$

$$\begin{aligned} x = ct &\quad \frac{dx}{dt} = c \\ y = \frac{c}{t} &\quad \frac{dy}{dt} = -\frac{c}{t^2} \\ \therefore \frac{dy}{dx} &= -\frac{1}{t^2} \end{aligned}$$

gradient of normal is  $t^2$   
in this case is  $p^2$

B

6)  $t = \tan \frac{x}{2}$

$$\begin{aligned} \frac{dt}{dx} &= \frac{1}{2} \sec^2 \frac{x}{2} \\ &= \frac{1}{2}(1+t^2) \\ dx &= \frac{2dt}{1+t^2} \end{aligned}$$

$$\int \sec x \, dx = \int \frac{1+t^2}{1-t^2} \cdot \frac{2}{1+t^2} dt$$

$$\begin{aligned} &= \int \frac{2}{1-t^2} dt \\ &= \int \frac{A}{1-t} + \frac{B}{1+t} dt \\ A(1+t) + B(1-t) &= 2 \end{aligned}$$

$$\text{Let } t = -1 \quad \therefore B = 1$$

$$\begin{aligned} t &= 1 \quad \therefore A = 1 \\ \therefore \int \frac{1}{1-t} + \int \frac{1}{1+t} &= -\ln(1-t) + \ln(1+t) = \ln \frac{1+t}{1-t} + C. \quad B \end{aligned}$$

$$7) \quad I_n = \int_0^{\pi} \sin^n x \, dx \quad 0 \leq x \leq \frac{\pi}{2}$$

$$\int_0^{\pi} \sin^{n-1} x \sin x \, dx$$

$$u = \sin x \quad u' = (n-1)\sin^{n-2} x \cos x \\ v' = \sin x \quad v = -\cos x$$

$$I_n = [-\cos x \sin^{n-1} x] + (n-1) \int \cos^2 x \sin^{n-2} x \, dx$$

$$\begin{aligned} I_n &= (n-1) \int (1 - \sin^2 x) \sin^{n-2} x \, dx \\ &= (n-1) \int \sin^{n-2} x - (n-1) \int \sin^n x \, dx \end{aligned}$$

$$I_n (1+n-1) = (n-1) \int \sin^{n-2} x \, dx \\ n I_n = n-1 I_{n-2}$$

$$I_n = \frac{n-1}{n} (I_{n-2})$$

A

$$\begin{aligned} 8) \quad & \int_0^2 \pi x y \, dx \\ & \text{Diagram: A quarter circle from } (0,0) \text{ to } (2,2) \\ & = \int_0^2 \pi x \cdot x^3 \, dx \\ & = \int_0^2 \pi x^4 \, dx \\ & = \pi \left[ \frac{x^5}{5} \right]_0^2 \\ & = \frac{32\pi}{5} \quad B \end{aligned}$$

$$9) \quad \tan \theta = \frac{v^2}{rg}$$

$$= \frac{25^2}{100 \times 9.8}$$

$$= 0.6377$$

$$\theta = 32^\circ 32'$$

B

10)

$$\alpha \text{ & } \beta \text{ satisfy } x^3 + 4x^2 - 2x - 5 = 0$$

$$\alpha^2 \beta^2 \gamma^2 \text{ satisfy } (\alpha^{\frac{1}{2}})^3 + 4(\alpha^{\frac{1}{2}})^2 - 2(\alpha^{\frac{1}{2}}) - 5 = 0$$

$$x^{\frac{3}{2}} + 4x - 2x^{\frac{1}{2}} - 5 = 0$$

$$2x^{\frac{3}{2}} - 2x^{\frac{1}{2}} = -4x + 5$$

$$x^{\frac{1}{2}}(2x-2) = -4x + 5$$

$$x(x-2)^2 = (-4x+5)^2$$

$$x(x^2 - 4x + 4) = 16x^2 - 40x + 25$$

$$x^3 - 4x^2 + 4x = 16x^2 - 40x + 25$$

$$x^3 - 20x^2 + 44x - 25 = 0$$

B

### Section 2. Question 11

$$a) i) \int \frac{t^2-1}{t^3} dt$$

$$\int \frac{t^2}{t^3} - \int \frac{1}{t^3}$$

$$\int \frac{1}{t} - \int t^{-3}$$

$$\ln t - \frac{t^{-2}}{-2}$$

$$\ln t + \frac{1}{2t^2} + C$$

$$ii) \int \frac{dx}{\sqrt{6-x-x^2}}$$

$$\int \frac{dx}{\sqrt{-(x^2+x-6)}}$$

$$\int \frac{dx}{\sqrt{-(x+\frac{1}{2})^2 - \frac{25}{4}}}$$

$$= \int \frac{dx}{\sqrt{\frac{25}{4} - (x+\frac{1}{2})^2}}$$

$$= \sin^{-1} \frac{x+\frac{1}{2}}{\frac{5}{2}}$$

$$= \sin^{-1} \left( \frac{2x+1}{5} \right) + C.$$

$$b) i) \int_0^1 \frac{x}{(x+1)(2x+1)} dx \quad \left( \frac{A}{x+1} + \frac{B}{2x+1} \right) = x \\ A(2x+1) + B(x+1) = x$$

$$\text{let } x=-1 \quad -A = -1$$

$$A = 1$$

$$x = -\frac{1}{2} \quad \frac{1}{2}B = -\frac{1}{2}$$

$$B = -1$$

$$\int_0^1 \frac{1}{x+1} + \int_0^1 \frac{1}{2x+1} dx \\ = \int_0^1 \frac{1}{x+1} - \frac{1}{2} \int_0^1 \frac{2}{2x+1} dx \quad \left[ \ln(x+1) - \frac{1}{2} \ln(2x+1) \right]_0^1 \\ = \ln\left(\frac{2}{3}\right) - \ln\left(\frac{2\sqrt{3}}{5}\right)$$

$$\begin{aligned}
 & \text{(ii)} \int_0^{\frac{\pi}{4}} x \tan x dx \quad u = x \quad \frac{du}{dx} = 1 \\
 & \quad \frac{du}{dx} = \sec^2 x - 1 \quad v = \tan x - x \\
 & \therefore I = \left[ x(\tan x - x) \right]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} (\tan x - x) dx \\
 & = \left[ x(\tan x - x) \right]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} \frac{\sin x}{\cos x} - x dx \\
 & = \left[ x(\tan x - x) + \ln(\cos x) + \frac{x^2}{2} \right]_0^{\frac{\pi}{4}} \\
 & = \left[ x \tan x - \frac{x^2}{2} + \ln(\cos x) \right]_0^{\frac{\pi}{4}} \\
 & = \frac{\pi}{4} - \frac{\pi^2}{32} + \ln \sqrt{2}.
 \end{aligned}$$

$$\begin{aligned}
 & \text{(iii)} I_n = \int_0^{\frac{\pi}{2}} x^n \cos x dx \quad u = x^n \quad \frac{du}{dx} = n x^{n-1} \\
 & \quad \frac{du}{dx} = \cos x \quad v = \sin x \\
 & \left[ x^n \sin x \right]_0^{\frac{\pi}{2}} - \int \sin x n x^{n-1} dx \\
 & \left( \frac{\pi}{2} \right)^n + n \int_0^{\frac{\pi}{2}} x^{n-1} \sin x dx \quad u = x^{n-1} \quad \frac{du}{dx} = (n-1)x^{n-2} \\
 & \quad \frac{du}{dx} = \sin x \quad v = -\cos x \\
 & \left( \frac{\pi}{2} \right)^n + n \left[ x^{n-1} \cdot \cos x \right]_0^{\frac{\pi}{2}} - n \int \cos x (n-1)x^{n-2} dx \\
 & = \left( \frac{\pi}{2} \right)^n + 0 - n(n-1) I_{n-2}.
 \end{aligned}$$

$$I_n = \left( \frac{\pi}{2} \right)^n - n(n-1) I_{n-2}.$$

$$\text{(iv)} \quad \int_0^{\frac{\pi}{2}} x^4 dx$$

$$\begin{aligned}
 A &= I_4 \\
 &= \left( \frac{\pi}{2} \right)^4 - 4 \times 3 \left[ \left( \frac{\pi}{2} \right)^2 - 2 \int_0^{\frac{\pi}{2}} \cos x dx \right] \\
 &= \left( \frac{\pi}{2} \right)^4 - 3\pi^2 + 24 \left[ \sin x \right]_0^{\frac{\pi}{2}} \\
 &= \left( \frac{\pi}{2} \right)^4 - 3\pi^2 + 24
 \end{aligned}$$

Question 12

$$\begin{aligned}
 & \text{(i)} \quad -\sqrt{2} (\sqrt{2} - \sqrt{2}i)^2 \\
 & = -\sqrt{2} (2 - 4i + 2i^2) \\
 & = -\sqrt{2} (2 - 4i) \\
 & = 4\sqrt{2}i
 \end{aligned}$$

$$\begin{aligned}
 & \text{(ii)} \quad \begin{array}{c} \sqrt{2} \\ \diagdown \\ 2 \\ \diagup \\ \sqrt{2} \end{array} \\
 & = -\frac{\pi}{4}
 \end{aligned}$$

$$\begin{aligned}
 & \text{(iii)} \quad \frac{z}{z+w} \\
 & \quad \frac{\sqrt{2}-\sqrt{2}i}{\sqrt{2}-\sqrt{2}i+\sqrt{2}} \\
 & = \frac{\sqrt{2}-\sqrt{2}i}{-\sqrt{2}i} \times \frac{\sqrt{2}+i}{\sqrt{2}+i} \\
 & = \frac{2i-2i^2}{-2i^2} \\
 & = \frac{2i+2}{2} \\
 & = 1+i
 \end{aligned}$$

$$\begin{aligned}
 & \text{(iv)} \quad |z| \\
 & = \sqrt{\sqrt{2}^2 + \sqrt{2}^2} \\
 & = \sqrt{4} \\
 & = 2
 \end{aligned}$$

$$\begin{aligned}
 & \text{(v)} \quad z^{10} = 2 \left( \text{cis } -\frac{\pi}{4} \right)^{10} \\
 & = 2^{10} \text{cis } -\frac{10\pi}{4} = 1024 \text{cis } -\frac{2\pi}{4} = \text{cis } -\frac{\pi}{2} = -i 1024
 \end{aligned}$$

$$b) (a+bi)^2 = 5-12i$$

$$a^2 + 2abi - b^2 = 5-12i$$

$$a^2 - b^2 = 5 \quad \text{--- (1)}$$

$$2ab = -12 \quad \text{--- (2)}$$

$$ab = -6$$

$$b = -\frac{6}{a}$$

$$a^2 - \frac{36}{a^2} = 5$$

$$a^4 - 5a^2 - 36 = 0$$

$$(a^2-9)(a^2+4)=0$$


$$a^2=9 \quad \therefore a=\pm 3$$

$$2(\pm 3)b = -12$$

$$\therefore b=\pm 2$$

$$a=3, b=2$$

$$\text{or } a=-3, b=+2$$



$$d(i) \frac{1+\cos\theta+i\sin\theta}{1-\cos\theta-i\sin\theta} = \frac{2\cos^2\frac{\theta}{2} + i2\sin\frac{\theta}{2}\cos\frac{\theta}{2}}{2\sin^2\frac{\theta}{2} - i2\sin\frac{\theta}{2}\cos\frac{\theta}{2}}$$

$$= \frac{2\cos^2\frac{\theta}{2}(\cos\frac{\theta}{2} + i\sin\frac{\theta}{2})}{2\sin^2\frac{\theta}{2}(\sin\frac{\theta}{2} - i\cos\frac{\theta}{2})}$$

$$= \frac{\cos\frac{\theta}{2}(\cos\frac{\theta}{2} + i\sin\frac{\theta}{2})}{-\sin\frac{\theta}{2}(\cos\frac{\theta}{2} + i\sin\frac{\theta}{2})}$$

$$\text{since } \sin\theta - i\cos\theta = -i(\cos\theta + i\sin\theta)$$

$$= i\cot\frac{\theta}{2} \quad \text{since } \frac{1}{-i} = i$$

c.

$$\frac{(z-1)^8}{(z+1)^4} = -1 \Rightarrow \frac{z-1}{z+1} = \sqrt[8]{-1}$$

$$\frac{z-1}{z+1} = \sqrt[8]{\cos(7\pi + 2k\pi)} = \cos\left(\frac{(2k+1)\pi}{8}\right) \quad k=0, \pm 1, \pm 2$$

$$z-1 = \left(\cos\left(\frac{(2k+1)\pi}{8}\right)\right)(z+1)$$

$$z-1 = \cos\left(\frac{(2k+1)\pi}{8}\right)z + \cos\left(\frac{(2k+1)\pi}{8}\right)$$

$$z\left(1 - \cos\left(\frac{(2k+1)\pi}{8}\right)\right) = \cos\left(\frac{(2k+1)\pi}{8}\right) + 1$$

$$\therefore z = \frac{1 + \cos\left(\frac{(2k+1)\pi}{8}\right)}{1 - \cos\left(\frac{(2k+1)\pi}{8}\right)}$$

$$= i\cot\left(\frac{(2k+1)\pi}{16}\right) \quad \text{from (1)}$$

$$= i\cot\frac{\pi}{16}, i\cot\frac{3\pi}{16}$$

$$= \pm i\cot\frac{\pi}{16}, \pm i\cot\frac{3\pi}{16} \quad \text{since } \cot x \text{ is an odd function}$$

d(i)

Alternatively Let  $t = \tan\frac{\theta}{2}$ .

$$\text{L.H.S} \quad \frac{1 - t^2}{1 + t^2} + \frac{i2t}{1 + t^2}$$

$$= \frac{1 - t^2 - i2t}{1 + t^2}$$

$$= \frac{2 + i2t}{2t^2 - i2t}$$




$$= \frac{1 + it}{t^2 - it}$$

$$= \frac{i(t-i)}{t(t-i)}$$

$$= \frac{i}{t} = i\cot\frac{\theta}{2} = \text{R.H.S}$$


Question 3

i)  $y = |f(x)|$



iv)  $y = \log f(x)$


As  $x \rightarrow -2$  or  $1$   $f(x) \rightarrow 0$   $\log f(x) \rightarrow +\infty$



crosses  $x$  axis when  $\ln f(x) = 0$   
ie when  $f(x) = 1$

$$\begin{aligned}x^2 + x - 2 &= 1 \\x^2 + x - 3 &= 0 \\x &= \frac{-1 \pm \sqrt{13}}{2}\end{aligned}$$

(b) i)



$$A = 2y \cdot \sqrt{3}y \cdot \frac{1}{2}$$

$$A = \sqrt{3}y^2$$

$$\int V = \sqrt{3}y^2 dx$$

$$V = \sum_{x \geq 0} \sqrt{3}y^2 dx$$

$$= \int_0^1 \sqrt{3}y^2 dx$$

$$= \sqrt{3} \int_0^1 (1-x)^4 dx$$

$$= 2\sqrt{3} \int_0^1 (1-u)^4 du$$

(ii)  $u = 1-x$   $x = 1-u$   $dx = -du$

$$u=0 \quad u=1$$

$$u=1 \quad u=0$$

$$= 2\sqrt{3} \int_1^0 u^4 \times -2(1-u) du$$

$$= 4\sqrt{3} \int_1^0 u^4 - u^5 du$$

$$= 4\sqrt{3} \left[ \frac{1}{5}u^5 - \frac{1}{6}u^6 \right]_1^0$$

$$= 4\sqrt{3} \left[ \frac{1}{5} - \frac{1}{6} \right]$$

$$= \frac{4\sqrt{3}}{30} = \frac{2\sqrt{3}}{15}$$

C) i) Aim prove  $(AB)^2 = BC \cdot BE$

Proof: In  $\triangle ABC$  and  $\triangle EBA$ ,

$\angle AEB = \angle CDE$  (alternate  $\angle$ 's on parallel lines)

$\angle CDE = \angle CAB$  (angles in the same segment)

$\therefore \angle AEB = \angle CAB - A$

$\angle BAE = \angle BCA$  (angle in the alternate segment)

$\therefore \triangle ABC \sim \triangle EBA$  equiangular.

$$\therefore \frac{AB}{BC} = \frac{BE}{AB} \text{ or } AB^2 = BC \cdot BE$$

ii)  $\frac{AC}{AE} = \frac{BC}{BA}$  since similar triangles have sides in proportion.

$$AB^2 = BC \cdot BE$$

$$\therefore AB = \sqrt{BC \cdot BE}$$

$$\frac{AC}{AE} = \frac{BC}{\sqrt{BC \cdot BE}}$$

$$= \frac{BC}{\sqrt{BC} \sqrt{BE}}$$

$$\frac{AC}{AE} = \frac{\sqrt{BC}}{\sqrt{BE}}$$

### Question 14

$$(a) 4x^2 + 9y^2 = 36$$

$$(i) \frac{4x^2}{36} + \frac{9y^2}{36} = 1$$

$$a^2 = 9 \quad b^2 = 4 \quad b^2 = a^2(1-e^2)$$

$$4 = 9(1-e^2)$$

$$\frac{4}{9} = 1 - e^2$$

$$e^2 = \frac{5}{9}$$

$$e = \frac{\sqrt{5}}{3}$$

$$S(ae, 0) \quad S'(ae, 0)$$

$$S(\sqrt{5}, 0) \quad S'(-\sqrt{5}, 0)$$

$$(i) x = \pm \frac{a}{e}$$


$$= \pm \frac{3}{\frac{\sqrt{5}}{3}}$$

$$= \pm \frac{9}{\sqrt{5}}$$

$$M: x = \frac{9}{\sqrt{5}}$$

$$M': x = -\frac{9}{\sqrt{5}}$$

(iii)



$$(IV) SP + SP' = 6$$

$$PS = ePM$$

$P'S' = eP'M'$  where  $M$  and  $M'$  are the feet of the perpendiculars from  $P$  to  $m$  and  $m'$ .

$$PS + P'S' = e(PM + PM')$$

$$= e(MM')$$

$$= e\left(\frac{a}{e} + \frac{a}{e}\right)$$

$$= \frac{2ae}{e}$$

$$PS + P'S' = 2a$$

$$a = 3.$$


$$\therefore SP + SP' = 6$$

$$(V) \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$\frac{3x}{9} + \frac{2y}{4} = 1$$

$$\frac{x}{3} + \frac{y}{2} = 1.$$

b(i)



$$(VI) xy = c^2$$

$$y = \frac{c^2}{x}$$

$$\frac{dy}{dx} = -\frac{c^2}{x^2}$$

$$\text{at } P, \text{ grad tangent} = -\frac{1}{t^2}$$

$$\therefore \text{grad normal} = t^2.$$

$$\tan \text{ of tang } y - \frac{c}{t} = -\frac{1}{t^2}(x - ct)$$

$$t^3y - ct = -x + ct$$

$$\text{Equation of } \frac{x + t^2y}{\text{normal}} = 2ct$$

$$y - \frac{c}{t} = t^2(x - ct)$$

$$ty - c = t^3x - ct^4$$

$$ty + ct^4 = t^3x + c.$$

$$(VI) \quad \begin{array}{l} \cancel{(2ct, 0)} \\ Y(0, \frac{2c}{t}) \end{array}$$

$$R\left(\frac{c(t^{2+1})}{t}; \frac{c(t^{2+1})}{t}\right)$$

$$S\left(\frac{c(t^{2+1})}{t}, -\frac{c(t^{2+1})}{t}\right)$$

$$\text{Midpoint } XY = (ct, \frac{c}{t})$$

$$\text{Midpoint } RS = \left(\frac{2ct^2}{t}, \frac{2c}{t}\right)$$

$$= ct, \frac{c}{t}$$

$$\text{Grad } XY = \frac{\frac{c}{t}}{-2ct}$$

$$= -\frac{1}{t^2}$$

$$\text{Grad } RS = \frac{\frac{2ct^2}{t}}{\frac{2c}{t}}$$

$$= t^2$$

$$\therefore RS \perp XY$$

$\therefore RDSX$  is a rhombus

Question 15

(a)  $P(x) = (x+1)(x-3)$ .  $\therefore P(x) + ax+b$ .

$$P(-1) = -a+b = b \quad \text{--- (1)}$$

$$P(3) = 3a+b = -2 \quad \text{--- (2)}$$

$$(1) - (2) \quad -4a = 8 \\ a = -2$$

$$\therefore b = 4$$

$$x^2 - 2x - 3 = (x+1)(x-3)$$

when divided by  $x^2 - 2x - 3$

$$\therefore P(x) = -2x + 4.$$

(b) (i)  $\alpha, \beta, \gamma \quad \therefore \alpha^2 + \beta^2 + \gamma^2 = (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \gamma\alpha) = 0 - 2p.$

$$\alpha^3 + \beta^3 + \gamma^3 = -2p.$$

(ii)

$$\begin{aligned} \alpha^3 + p\alpha + q &= 0 \\ \beta^3 + p\beta + q &= 0 \\ \gamma^3 + p\gamma + q &= 0 \end{aligned}$$

$$\alpha^3 + \beta^3 + \gamma^3 + p(\alpha + \beta + \gamma) + 3q = 0$$

$$\alpha^3 + \beta^3 + \gamma^3 + p \cdot 0 + 3q = 0$$

$$\therefore \alpha^3 + \beta^3 + \gamma^3 = -3q$$

(c) Let  $y = x-1$ .

$$\therefore x = y+1.$$

$$3(y+1)^3 - 5(y+1)^2 - 4(y+1) + 3 = 0$$

$$3(y^3 + 3y^2 + 3y + 1) - 5(y^2 + 2y + 1) - 4(y+1) + 3 = 0$$

$$3y^3 + 9y^2 + 9y + 3 - 5y^2 - 10y - 5 - 4y - 4 + 3 = 0$$

$$3y^3 + 4y^2 - 5y - 3 = 0$$

in terms of  $x$

$$3x^3 + 4x^2 - 5x - 3 = 0$$

(d) (i)  $P(x) = x^n + ax - b$

double root if  $x = \alpha$

$$P(x) = nx^{n-1} + a$$

$$P''(x) = n(n-1)x^{n-2}$$

$$\text{note } P(x) = 0 \Rightarrow \alpha^n + a\alpha - b = 0$$

$$P'(\alpha) = 0 \Rightarrow n\alpha^{n-1} + a = 0$$

$$\therefore \alpha^{n-1} = -\frac{a}{n}$$

(ii)  $P(x) = \alpha^n + a\alpha - b = 0 \quad \text{--- (1)}$

$$P'(\alpha) = n\alpha^{n-1} + a = 0 \quad \text{--- (2)}$$

$$n\alpha^{n-1} + a\alpha = 0 \quad \text{--- (2)}$$

$$(1) - \alpha(2) \quad (1-n)\alpha^n - b = 0$$

$$\alpha^n = \frac{b}{1-n} \quad \text{--- (3)}$$

$$\text{also } \alpha^{n-1} = -\frac{a}{n} \quad \text{--- (4)}$$

$$\text{from (3)} \quad (\alpha^n)^{n-1} = \left(\frac{b}{1-n}\right)^{n-1}$$

$$\text{from (4)} \quad (\alpha^{n-1})^n = \left(-\frac{a}{n}\right)^n$$

$$\left(\frac{b}{1-n}\right)^{n-1} = \left(-\frac{a}{n}\right)^n$$

$$\left(\frac{b}{n-1}\right)^{n-1} = (-1)^n \left(\frac{a}{n}\right)^n$$

$$(-1)^{n-1} \left(\frac{b}{n-1}\right)^{n-1} = (-1)^{n-1} \left(\frac{a}{n}\right)^n$$

$$\left(\frac{a}{n}\right)^n + \left(\frac{b}{n-1}\right)^{n-1} = 0$$

(iii) double root is  $\omega$

$$\omega = \frac{\omega^n}{\omega^{n-1}} \quad \text{use (3)}$$

$$\omega = \frac{\omega^n}{\omega^{n-1}} \quad \text{use (4)}$$

$$= \left(\frac{b}{1-n}\right) / -\frac{a}{n}$$

$$\omega = \frac{bn}{-a(1-n)} = \frac{bn}{a(n-1)}$$

Question 16.

(a) Let  $x = \frac{a+b}{2}$

$$y = \frac{ct+d}{2}$$

$$\therefore \frac{xy}{2} = \frac{ab+ctd}{4}$$

$$\text{Now } \frac{xy}{2} \geq \sqrt{xy}$$

$$\therefore \frac{ab+ctd}{4} \geq \sqrt{\sqrt{ac}\sqrt{cd}}$$

$$\frac{ab+ctd}{4} \geq \sqrt[4]{abcd}$$

b) i)  $\text{cis } 5\theta = (\text{cis } \theta)^5$

$$\cos 5\theta + i \sin 5\theta = \cos^5 \theta + i \cdot 5 \cos^4 \theta \sin \theta - 10 \cos^3 \theta \sin^3 \theta - i(10 \cos^3 \theta \sin^3 \theta + 5 \cos \theta \sin^4 \theta + i \sin^5 \theta)$$

$$\therefore \cos 5\theta = \cos^5 \theta - 10 \cos^3 \theta \sin^3 \theta + 5 \cos \theta \sin^4 \theta$$

$$\sin 5\theta = 5 \cos^4 \theta \sin \theta - 10 \cos^3 \theta \sin^3 \theta + 5 \cos \theta \sin^4 \theta$$

$$\tan 5\theta = \frac{5 \cos^4 \theta \sin \theta - 10 \cos^3 \theta \sin^3 \theta + 5 \cos \theta \sin^4 \theta}{\cos^5 \theta - 10 \cos^3 \theta \sin^2 \theta + 5 \cos \theta \sin^4 \theta}$$

$$= \frac{5 \tan \theta - 10 \tan^3 \theta + \tan^5 \theta}{1 - 10 \tan^2 \theta + 5 \tan^4 \theta}$$

$$= \tan \theta \cdot \frac{5 - 10 \tan^2 \theta + \tan^4 \theta}{1 - 10 \tan^2 \theta + 5 \tan^4 \theta}$$

(ii) Let  $x = \tan \theta$  then  $\tan 5\theta = 0 \quad \therefore x^4 - 10x^2 + 5 = 0$

$$5\theta = 0 \text{ or } \pi \text{ or } 2\pi, \dots$$

$$\theta = \frac{\pi}{5}, \frac{2\pi}{5}, \frac{3\pi}{5}, \frac{4\pi}{5}$$

$$\tan \frac{3\pi}{5} = -\tan \left(\pi - \frac{3\pi}{5}\right)$$


$$= -\tan \frac{2\pi}{5} \quad \text{and} \quad \tan \frac{4\pi}{5} = -\tan \frac{\pi}{5}$$

root  $x = \pm \tan \frac{\pi}{5}$  and  $\pm \tan \frac{2\pi}{5}$ ,  
product of roots  $(\tan \frac{\pi}{5})(\tan \frac{2\pi}{5})(\tan \frac{3\pi}{5})(\tan \frac{4\pi}{5}) = 5 = 5$ .

$$\alpha \beta \gamma \delta = \frac{c}{a}$$

Question 16(c)

(i)



$$\cos 60 = \frac{AB}{2}$$


$$AB = 1.$$

$$\sin 60 = \frac{r}{AB}$$

$$r = \frac{\sqrt{3}}{2}.$$

(ii) Let tension in rods AB and BC

be  $T_1$  and  $T_2$  respectively



$$\sum F_V = 0$$

$$T_1 \cos 60 = T_2 \cos 30 + mg$$

$$T_1 \left(\frac{1}{2}\right) = T_2 \left(\frac{\sqrt{3}}{2}\right) + 10g$$

$$T_1 = T_2 \sqrt{3} + 20g$$

$$\sum F_H = mrw^2$$

$$T_1 \sin 60 + T_2 \sin 30 = mrw^2$$

$$T_1 \left(\frac{\sqrt{3}}{2}\right) + T_2 \left(\frac{1}{2}\right) = 10 \left(\frac{\sqrt{3}}{2}\right) (3\pi)^2$$

$$T_1 = 90\pi^2 - \frac{T_2}{\sqrt{3}}$$

$$T_2 \sqrt{3} + 20g = 90\pi^2 - \frac{T_2}{\sqrt{3}}$$

$$T_2 = \frac{5\sqrt{3}}{2} (9\pi^2 - 2g)$$

$$T_1 = \frac{5}{2} (27\pi^2 + 2g)$$

so tensions in AB and BC are  
 $\sum (27\pi^2 + 2g) N$   
 $\text{and } \sum \frac{5\sqrt{3}}{2} (9\pi^2 - 2g) N$

16(d).  $n=1$

$$a_1 < \sqrt{2+q_0} = \sqrt{3}.$$

Since  $\sqrt{2} < \sqrt{3} < 2$  is true for  $n=1$ .

Assume true for  $n=k$

$$\sqrt{2} < a_k < 2 \quad (A)$$

Now prove true for  $n=k+1$

$$\sqrt{2} < a_{k+1} < 2 \quad (B)$$

From (A)  $\sqrt{2} < a_k < 2$

$$2+\sqrt{2} < 2+a_k < 4$$

$$\sqrt{2+\sqrt{2}} < \sqrt{2+a_k} < 2$$

$$\sqrt{2+\sqrt{2}} < a_{k+1} < 2$$

$$\text{Now } 2 < 2+\sqrt{2} \Rightarrow \sqrt{2} < \sqrt{2+\sqrt{2}}$$

$$\therefore \sqrt{2} < a_{k+1} < 2$$

$\therefore$  proved by mathematical induction.