

Name:

Maths Teacher:

SYDNEY TECHNICAL HIGH SCHOOL



Year 12

Mathematics Extension 2

TRIAL HSC

2016

Time allowed: 3 hours plus 5 minutes reading time

General Instructions:

- Reading time – 5 minutes
- Working time – 3 hours
- Marks for each question are indicated on the question.
- Board - Approved calculators may be used
- In Questions 11- 16, show relevant mathematical reasoning and/or calculations.
- Full marks may not be awarded for careless work or illegible writing
- ***Begin each question on a new page***
- Write using black pen
- All answers are to be in the writing booklet provided
- A BOSTES reference sheet is provided at the rear of this Question Booklet, and may be removed at any time.

Total marks – 100

Section I Multiple Choice

10 Marks

- Attempt Questions 1-10
- Allow 15 minutes for this section

Section II

90 Marks

- Attempt Questions 11-16
- Allow 2 hours 45 minutes for this section

Section I

10 marks

Attempt Questions 1- 10

Allow about 15 minutes for this section

Use the multiple- choice answer sheet located in your answer booklet for Questions 1 -10

1. Which conic has eccentricity $\frac{\sqrt{3}}{3}$?

(A) $\frac{x^2}{3} + \frac{y^2}{2} = 1$

(B) $\frac{x^2}{3^2} + \frac{y^2}{2^2} = 1$

(C) $\frac{x^2}{3} - \frac{y^2}{2} = 1$

(D) $\frac{x^2}{3^2} - \frac{y^2}{2^2} = 1$

2. What value of z satisfies; $z^2 = 20i - 21$?





(A) $-2+5i$

(B) $2-5i$

(C) $2+5i$

(D) $5-2i$

3. Which graph represents the curve, $y=(x+3)^2(x-1)^3$?



4. The polynomial $2x^4 - 17x^3 + 45x^2 - 27x - 27$ has a triple root at $x = \alpha$.

What is the value of α ?

(A) $-\frac{1}{2}$

(B) $\frac{1}{2}$

(C) -3

(D) 3

5. If $z_1 = 1+2i$ and $z_2 = 3-i$ then $z_1 \div \overline{z_2}$ is,

(A) $\frac{1}{2} - \frac{1}{2}i$

(B) $\frac{1}{2} + \frac{1}{2}i$

(C) $4+3i$

(D) $\frac{5}{8} + \frac{5}{8}i$

6. Which expression is equal to, $\int \frac{x^2}{\cos^2 x} dx$?

(A) $2x \tan x - 2 \int \tan x dx$

(B) $\frac{1}{3}(x^3 \sec^2 x - \int x^3 \tan x dx)$

(C) $x^2 \tan x - 2 \int x \tan x dx$

(D) $x^2 \tan x - 2 \int x \sec^2 x dx$

7. What is the natural domain of the function $f(x) = \frac{1}{2}(x\sqrt{x^2-1} - \ln(x+\sqrt{x^2-1}))$?

(A) $x \leq -1$ or $x \geq 1$

(B) $-1 \leq x \leq 1$

(C) $x \geq 1$

(D) $x \leq -1$

8. If α, β, δ are the roots of $x^3 + x - 1 = 0$, then an equation with roots

$$\frac{(\alpha+1)}{2}, \frac{(\beta+1)}{2}, \frac{(\delta+1)}{2}$$
 is?

- (A) $x^3 - 3x^2 + 4x - 3 = 0$
(B) $x^3 + 3x^2 + 4x + 1 = 0$
(C) $x^3 - 6x^2 + 16x - 24 = 0$
(D) $8x^3 - 12x^2 + 8x - 3 = 0$


9. The complex number Z satisfies $|Z + 2| = 1$

What is the smallest positive value of the $\arg(z)$ on the Argand diagram?

- (A) $\frac{\pi}{3}$
(B) $\frac{5\pi}{6}$
(C) $\frac{2\pi}{3}$
(D) $\frac{\pi}{6}$

10. The base of a solid is the region bounded by the parabola $x = 4y - y^2$ and the y -axis.

Vertical cross sections are right angled isosceles triangles perpendicular to the x -axis as shown.



Which integral represents the volume of this solid?

- (A) $\int_0^4 2\sqrt{4-x} dx$
(B) $\int_0^4 \pi(4-x) dx$
(C) $\int_0^4 (8-2x) dx$
(D) $\int_0^4 (16-4x) dx$

Question 11 (15 marks)

(a) Express $\frac{18+4i}{3-i}$ in the form, $x+iy$, where x and y are real.

2

(b) Consider the complex numbers $z = -1 + \sqrt{3}i$ and $w = \sqrt{2} \left(\cos\left(\frac{-\pi}{4}\right) + i \sin\left(\frac{-\pi}{4}\right) \right)$

(I) Evaluate $|z|$

1

(II) Evaluate $\arg(z)$

1

(III) Find the argument of $\frac{w}{z}$

2

(c) (i) Find A, B and C such that

3


$$\frac{1}{x(x^2+4)} = \frac{A}{x} + \frac{Bx+C}{x^2+4}$$

(ii) Hence, or otherwise, find;

$$\int \frac{dx}{x(x^2+4)}$$

2

(d)



A particle moves along the x -axis. At time, $t=0$, the particle is at $x=0$.

Its velocity v at time t is shown on the graph above.

Copy or trace this graph onto your answer page.

(i) At what time is the acceleration the greatest? Explain your answer.

1

(ii) At what time does the particle first return to $x=0$? Explain your answer.

1

(iii) Sketch the displacement time graph for the particle in the interval, $0 \leq t \leq 9$.

2

Question 12 (15 marks) START THIS QUESTION ON A NEW PAGE.

(a) Find $\int x\sqrt{x+1}dx$

2

(b) Evaluate

(i) $\int_0^{\frac{\pi}{4}} \sin x \cos 2x dx$

2

(ii) $\int_1^e \frac{\ln x}{x^2} dx$

2

(c) Find the equation of the normal to the curve, $3x^2y^3 + 4xy^2 = 6 + y$ at the point (1,1).

4

(d)

(i) Prove that,

$$\cos(A-B)x - \cos(A+B)x = 2 \sin Ax \sin Bx$$

1

(ii) Using the above result, express the equation $\sin 3x \sin x = 2 \cos 2x + 1$,
as a quadratic equation in terms of $\cos 2x$

2

(iii) Hence, solve, $\sin 3x \sin x = 2 \cos 2x + 1$ for $0 \leq x \leq 2\pi$

2

Question 13 (15 marks) START THIS QUESTION ON A NEW PAGE.

- (a) The function $y = f(x)$ is defined by the equation;

$$f(x) = \frac{x(x-4)}{4}$$

Without the use of calculus, draw sketches of each of the following, clearly labelling any
Intercepts, asymptotes and turning points.

- | | | |
|-------|------------------------|---|
| (i) | $y = f(x)$ | 1 |
| (ii) | $y^2 = f(x)$ | 2 |
| (iii) | $y = \frac{x x-4 }{4}$ | 2 |
| (iv) | $y = \tan^{-1} f(x)$ | 2 |
| (v) | $y = e^{f(x)}$ | 2 |

- (b) Sketch the locus of z satisfying

- | | | |
|------|-------------------------------------|---|
| (i) | $Re(z) = z $ | 2 |
| (ii) | $Im(z) \geq 2$ and $ z - 1 \leq 2$ | 2 |

- (c) Write down the domain and range of $y = 2 \sin^{-1} \sqrt{1-x^2}$ 2

Question 14 (15 marks) START THIS QUESTION ON A NEW PAGE.

- (a) Use the substitution $t = \tan \frac{x}{2}$ to find

4

$$\int_0^{\frac{\pi}{2}} \frac{1}{5 + 4\cos x + 3\sin x} dx$$

- (b) The area enclosed by the curves $y = \sqrt{x}$ and $y = x^2$ is rotated about the y -axis.


Use the method of **cylindrical shells** to find the volume of the solid formed.

4

Question 14 continues on the next page....

Question 14 continued....

(c)




$P(a \sec \theta, b \tan \theta)$ lies on the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$.

The tangent at P meets the line $x = -a$ and $x = a$ at R and Q respectively.

- | | | |
|-------|---|---|
| (i) | Show that the equation of the tangent is given by $\frac{x \sec \theta}{a} - \frac{y \tan \theta}{b} = 1$. | 2 |
| (ii) | Find the coordinates of Q and R . | 1 |
| (iii) | Show that QR subtends a right angle at the focus $S(ae, 0)$. | 2 |
| (iv) | Deduce that Q, S, R, S' are concyclic. | 2 |

Question 15 (15 marks) START THIS QUESTION ON A NEW PAGE.

- (a) In the diagram, AB and AC are tangents from A to the circle with centre O , meeting the circle at B and C respectively. ADE is a secant of the circle. G is the midpoint of DE . CG produced meets the circle at F .



- | | | |
|-------|--|---|
| (i) | Copy the diagram, using about one third of the page, into your answer booklet and prove that $ABOC$ and $AOGC$ are cyclic quadrilaterals | 3 |
| (ii) | Explain why $\angle OGF = \angle OAC$. | 1 |
| (iii) | Prove that $BF \parallel AE$ | 3 |

(b)

(i) Let $I_n = \int_0^1 x^n \sqrt{1-x^3} dx$ for $n \geq 2$.

Show that: $I_n = \frac{2n-4}{2n+5} \cdot I_{n-3}$ for $n \geq 5$

(ii) Hence find I_8 2

- (c) A sequence of numbers is given by $T_1 = 6$ $T_2 = 27$ and $T_n = 6T_{n-1} - 9T_{n-2}$ for $n \geq 3$.

Prove by Mathematical Induction that:

$$T_n = (n+1) \times 3^n \text{ for } n \geq 1 \quad \text{3}$$

Question 16 (15 marks) START THIS QUESTION ON A NEW PAGE.

(a) Show that the minimum value of $ae^{mx} + be^{-mx}$ is $2\sqrt{ab}$

if a, b and m are all positive constants.

4

(b) A particle of mass 1 kilogram is projected upwards under gravity (g) with a speed of $2k$

in a medium in which resistance to motion is $\frac{g}{k^2}$ times the square of the speed, where k is a positive constant.

(i) Show that the maximum height (H) reached by the particle is

$$H = \frac{k^2}{2g} \ln 5$$

3


(ii) Show that the speed with which the particle returns to its starting point

$$\text{is given by } V = \frac{2k}{\sqrt{5}}$$

4

(c) The shaded region in the diagram is bounded by the curves $y = \sin x$, $y = \cos x$ and the line $y = 1$.

This region is rotated around the y -axis.



Calculate the volume of the solid formed, using the process of **Volume by Slicing**.

4

STHS - Ext 2 Trial - Suggestion solution

Section 1

1. A 2. C 3. C 4. D 5. B
 6. C * 7. C 8. D 9. B 10. C
 (all given)

Section 2


Question 11

$$\begin{aligned} \text{a) } & \frac{18+4i}{3-i} \times \frac{3+i}{3+i} \\ & = \frac{54+18i+12i-4}{10} \\ & = \frac{50+30i}{10} \\ & = 5+3i \end{aligned}$$

$$\begin{aligned} \text{equating: } & 0 = A + B \\ & B = -\frac{1}{4} \\ & 0 = C \\ \text{ie } & A = \frac{1}{4}, B = -\frac{1}{4}, C = 0 \\ \text{ii. } & \text{Now } \int \frac{dx}{x(x^2+4)} = \int \frac{1}{4x} - \frac{1}{x^2+4} dx \end{aligned}$$

$$\begin{aligned} \text{b) } & \omega = \sqrt{2} \operatorname{cis}(-\pi/4) \quad z = -1 + \sqrt{3}i \\ \text{i) } & |z| = \sqrt{1+3} \\ & = 2 \\ \text{ii) } & \arg(z) = 2\pi/3 \\ \text{iii) } & \arg\left(\frac{\omega}{z}\right) = \arg(\omega) - \arg(z) \\ & = -\pi/4 - 2\pi/3 \\ & = -\frac{11\pi}{12}. \end{aligned}$$

$$\begin{aligned} \text{c) } & \frac{1}{x(x^2+4)} = \frac{A}{x} + \frac{Bx+C}{x^2+4} \\ \therefore & 1 = A(x^2+4) + x(Bx+C) \\ \text{let } x=0 & \\ 1 = A(4) & \rightarrow A = \frac{1}{4} \end{aligned}$$



Question 12

$$\text{a) } \int x \sqrt{x+1} dx$$

one method:

$$\text{let } u = x+1$$

$$\frac{du}{dx} = 1 \quad \therefore du = dx$$

$$= \int (u-1)\sqrt{u} du$$

$$= \int u\sqrt{u} - \sqrt{u} du$$

$$= \int u^{3/2} - u^{1/2} du$$

$$= \frac{2}{5}u^{5/2} - \frac{2}{3}u^{3/2}$$

$$= \frac{2}{5}(x+1)^{5/2} - \frac{2}{3}(x+1)^{3/2} + C$$

$$\text{c) } 3x^2y^3 + 4xy^2 = 6+y \text{ @}(1,1)$$

$$\text{b) i. } \int_0^{\pi/4} \sin x \cos 2x dx$$

$$= \int_0^{\pi/4} \sin x (2\cos^2 x - 1) dx$$

$$= \int_0^{\pi/4} 2\sin x (\cos x)^2 - \sin x dx$$

$$= \left[-\frac{2}{3} \cos^3 x + \cos x \right]_0^{\pi/4}$$

$$= -\frac{2}{3} \left(\frac{1}{\sqrt{2}} \right)^3 + \frac{1}{\sqrt{2}} - \left[-\frac{2}{3}(1)^3 + 1 \right]$$

$$\text{ii. } \int_1^e \frac{\ln x}{x^2} dx$$

$$= \int_1^e x^{-2} \ln x dx$$

$$= \ln x \cdot \frac{x^{-1}}{-1} - \int_1^e \frac{1}{x} \cdot \frac{-1}{x^2} dx$$

$$= -\frac{\ln x}{x} \Big|_1^e + \int_1^e x^{-2} dx$$

$$= -\left[\frac{1}{e} - 0 \right] + \left[-\frac{1}{x} \right]_1^e$$

$$= -\frac{1}{e} + \left[-\frac{1}{e} - (-1) \right]$$

$$= 1 - \frac{2}{e}$$

$$3x^2y^3 + 4xy^2 = \frac{dy}{dx}(1-9x^2y^2-8xy)$$

$$3x^2 \left[3y^2 \frac{dy}{dx} \right] + y^3 \cdot 6x + 4x^2 y \frac{dy}{dx} + 4y^2 = \frac{dy}{dx}$$

$$6xy^3 + 4x^2y = \frac{dy}{dx}(1-9x^2y^2-8xy)$$

$$\text{at } (1,1)$$

$$\frac{dy}{dx} = \frac{10}{-16}$$

$$M_T = -\frac{5}{8} \quad \therefore M_N = \frac{8}{5}$$

$$y-1 = \frac{8}{5}(x-1)$$

$$5y-5 = 8x-8$$

$$8x-5y-3=0$$

Question 12 - con't.

d)

i. Prove that

$$\cos(A-B)x - \cos(A+B)x = 2\sin Ax \sin Bx$$

$$\text{LHS} = \cos Ax \cos Bx + \sin Ax \sin Bx - [\cos Ax \cos Bx - \sin Ax \sin Bx]$$

$$= 2\sin Ax \sin Bx$$

= RHS.

$$\text{ii. } \sin 3x \sin x = 2\cos 2x + 1$$

$$A=3$$

$$B=1$$

$$\therefore [\cos(3-1)x - \cos(3+1)x] \div 2 = \cos 2x + 1$$

$$\cos 2x - \cos 4x = 2\cos 2x + 2$$

$$\cos 2x - [2\cos^2 2x - 1] = 2\cos 2x + 2$$

$$\cos 2x - 2\cos^2 2x + 1 = 2\cos 2x + 2$$

$$2\cos^2 2x + 3\cos 2x + 1 = 0$$

iii. hence,

$$(2\cos 2x + 1)(\cos 2x + 1) = 0$$

$$\cos 2x = -\frac{1}{2} \quad \text{or} \quad \cos 2x = -1$$


$$2x = \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{8\pi}{3}, \frac{10\pi}{3} \quad \text{or} \quad 2x = \pi, 3\pi$$

$$x = \frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{5\pi}{3} \quad x = \frac{\pi}{2}, \frac{3\pi}{2}$$


Question 13

a) $f(x) = \frac{x(x-4)}{4}$

1.



II.



b) $R(z) = |z|$


let $z = x + iy$

$$x = \sqrt{x^2 + y^2} \quad x \geq 0$$

$$x^2 = x^2 + y^2$$


$$y^2 = 0$$

$y = 0$ but $x \geq 0$




II. $\operatorname{Im}(z) \geq 2 \quad |z-1| \leq 2$

$\therefore y \geq 2$ circle centre $(1, 0)$



c) $y = 2\sin^{-1}\sqrt{1-x^2}$

IV.



$\therefore -1 \leq x \leq 1$

R; $0 \leq y \leq \pi$

Question 14.

a)

$$x = \frac{\pi}{2}, t = \tan \frac{\pi}{4} = 1$$

$$x=0, t=\tan 0=0$$

$$\therefore \int_0^1 \frac{1}{5+4\left(\frac{1-t^2}{1+t^2}\right) + 3\left(\frac{2t}{1+t^2}\right)} \cdot \frac{2}{t^2+1} dt$$

$$= \int_0^1 \frac{2}{5(1+t^2) + 4(1-t^2) + 6t} dt$$

$$= \int_0^1 \frac{2}{t^2+6t+9} dt$$

$$= 2 \int_0^1 \frac{1}{(t+3)^2} dt$$

$$= 2 \int_0^1 (t+3)^{-2} dt$$

$$= -2 \left[\frac{1}{t+3} \right]_0^1$$

$$= \left(-\frac{2}{4} \right) - \left(-\frac{2}{3} \right) = \frac{1}{6}$$

c)

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

$$2x/a^2 - 2y/b^2 \cdot \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = \frac{x}{y} \frac{b^2}{a^2}$$

$$P(a \sec \theta, b \tan \theta)$$

$$M_T = \frac{b}{a \sin \theta}$$

Now

$$y - \frac{b \sin \theta}{\cos \theta} = \frac{b}{a \sin \theta} \left(x - \frac{a}{\cos \theta} \right)$$

$$a \sin \theta y - a b \sin^2 \theta = b x - a b$$

$$\frac{\sin \theta y}{\cos \theta} - \frac{a b \sin^2 \theta}{\cos \theta} = \frac{b x}{\cos \theta} - \frac{a b}{\cos \theta}$$

$$(\div ab)$$

$$\frac{\sin \theta y}{b} - \frac{x}{a} = \frac{\sin^2 \theta - 1}{\cos \theta}$$

$$\frac{-\tan \theta y}{b} + \frac{x}{a \cos \theta} = 1$$

or

$$\frac{x \sec \theta}{a} - \frac{\tan \theta y}{b} = 1$$

b)

at Q $x=a$

$$\frac{a \sec \theta}{a} - \frac{\tan \theta y}{b} = 1$$

$$\frac{1}{\cos \theta} - \frac{y}{b} \frac{\sin \theta}{\cos \theta} = \frac{\cos \theta}{\cos \theta}$$

$$1 - \frac{y}{b} \sin \theta = \cos \theta$$

$$y = b(1 - \cos \theta)$$

$$\sin \theta$$

$$\Delta V = 2\pi x (\sqrt{x} - x^2) \Delta x$$

$$\lim_{\Delta x \rightarrow 0} \sum 2\pi x (\sqrt{x} - x^2) \Delta x$$

$$= 2\pi \int_0^1 x^{3/2} - x^3 dx$$

$$= \left[\frac{2}{5} x^{5/2} - \frac{1}{4} x^4 \right]_0^1 \times 2\pi$$

$$= 2\pi \left[\frac{2}{5} - \frac{1}{4} - 0 \right] = \frac{3\pi}{10} u^3$$

Q

$$\left[a, \frac{b(1-\cos \theta)}{\sin \theta} \right]$$

at R = -a

iv) $M_{RS1} = \frac{b(1-\cos \theta)}{a(e-i)\sin \theta}$

$$- \frac{a \sec \theta}{a} - \frac{y \tan \theta}{b} = 1$$

$$-1 - \frac{y}{b} \sin \theta = \cos \theta$$

$$\therefore M_{RS1} \times M_{RS1}$$

$$y = -\frac{b - b \cos \theta}{\sin \theta} = \frac{b^2(1-\cos^2 \theta)}{-a^2(e^2-1)\sin^2 \theta}$$

$$\text{or } y = \frac{-b(1+\sec \theta)}{\tan \theta} = \frac{b^2}{-b^2} = -1$$

iii) $S(ae, 0)$

$$M_{SQ} = 0 - \frac{b(1-\cos \theta)}{\sin \theta}$$

$$ae - a$$

$$= -\frac{b(1-\cos \theta)}{\sin \theta}$$

$$a(e-i)$$

making $\triangle QSR$ a cyclic quad. as opposite angles are supplementary.

$M_{RS} = \frac{b(1+\cos \theta)}{\sin \theta}$

Now $M_{SQ} \times M_{RS}$

$$= -\frac{b(1-\cos \theta)}{\sin \theta} \times \frac{b(1+\cos \theta)}{a(e-i)}$$

$$= -\frac{b^2(1-\cos^2 \theta)}{a^2(e^2-1)\sin^2 \theta}$$

$$= -\frac{b^2}{a^2(e^2-1)}$$

but

$$\frac{a^2(e^2-1)}{a^2(e^2-1)} = b^2$$

$$= -1 \therefore \angle QSR = 90^\circ$$

(5)

(6)

Question 15.

Join AO, BF BC

as BO = OC radii

$$\angle OCB = \angle CBO \text{ (equal angles)}$$

$$I. \angle ABO = \angle COA = 90^\circ$$

α opposite equal sides

(radii to tangent at point of contact is 90°)

Now in $\triangle BOC$

$$\angle BOC = 180 - 2\alpha \text{ (angle sum)}$$

\therefore opposite angles in

$$\therefore \angle BFC = 90 - \alpha$$

ABOC are supplementary and

(angle at the circumference is

ABOC is a cyclic quadrilateral.

half the angle at the centre on

$$\text{Now, } \angle ABO = 90^\circ$$

arc BC)

(AO is a diameter or Line from

$$\therefore \angle BFC = \angle FGE (90 - \alpha)$$

midpt to centre is perpendicular)

and the alternate angles

$$\angle OGA = \angle OCA \quad (\text{angles at} \\ \text{circumference} \\ \text{of circle } OGCA) \quad = 90^\circ$$

are equal

$$\therefore BF \parallel AE$$

$\therefore AOGC$ is a cyclic quad

as opposite angles are
supplementary.

$$II. \angle OGF = \angle OAC$$

exterior angle of a cyclic
quadrilateral equals opposite
interior angle ($AOGC$).

$$III. \text{ let } \angle OGF = \angle OAC = \alpha$$

$$\therefore \angle FGE = 90^\circ - \alpha \quad (\text{straight line})$$

and

$$\angle OBC = \angle OAC \quad (\text{angles in the} \\ \text{same segment} \\ = \alpha \quad \text{of } ABOC)$$

Question 15 con't

$$I_n = \int_0^1 x^n \sqrt{1-x^3} dx \quad n \geq 2$$

$$= \int_0^1 x^{n-2} \cdot x^2 \sqrt{1-x^3} dx$$

$$= \left[x^{n-2} (1-x^3)^{3/2} \right]_0^1 - \int_0^1 (n-2)x^{n-3} \cdot -2(1-x^3) \sqrt{1-x^3} dx$$

$$I_n = 0 + \frac{2(n-2)}{9} \int_0^1 x^{n-3} \sqrt{1-x^3} dx - \frac{2(n-2)}{9} \int_0^1 x^n \sqrt{1-x^3} dx$$

$$I_n \left[1 + \frac{2(n-2)}{9} \right] = \frac{2(n-2)}{9} \int_0^1 x^{n-3} \sqrt{1-x^3} dx$$

$$I_n \left[\frac{9+2n-4}{9} \right] = \frac{2n-4}{9} I_{n-3} \frac{dx}{x}$$

$$I_n = \frac{2n-4}{2n+5} I_{n-3}$$

ii)

$$J_8 = \left(\frac{16-4}{16+5} \right) I_5$$

$$= \frac{12}{21} \left[\frac{(10-4) I_2}{10+5} \right] \quad \text{Now } I_2 = \int_0^1 x^2 \sqrt{1-x^3} dx \\ = \frac{12}{21} \cdot \frac{6}{15} \cdot \frac{2}{9}$$

$$= \frac{(1-x^3)^{3/2}}{3/2 \cdot -3}$$

$$= \frac{16}{315} \quad = \left[\frac{-2(1-x^3)^{3/2}}{9} \right]_0^1$$

$$= -2 \left[0 - 1^{3/2} \right] \quad = -2 \left[\frac{2}{9} \right]$$

Question 15 con't

c)

$$T_1 = 6 \quad T_2 = 27$$

$$T_n = 6T_{n-1} - 9T_{n-2} \quad n \geq 3$$

$$T_n = (n+1)3^n \quad \text{for } n \geq 1$$

Test $n=1$

$$T_1 = (1+1) \times 3^1$$

$$= 2 \times 3$$

$$= 6 \quad \text{which is given}$$

\therefore True for $n=1$

Assume true for $n=k$

$$\text{ie } T_k = (k+1)3^k$$

where

$$T_k = 6T_{k-1} - 9T_{k-2}$$

Prove true for $n=k+1$

aim to prove

$$T_{k+1} = 6T_k - 9T_{k-1} = (k+1+1) \cdot 3^{k+1}$$

$$T_{k+1} = 6T_k - 9T_{k-1}$$

$$= 6(k+1) \cdot 3^k - 9[k(3^{k-1})] \quad \text{By assumption}$$

$$= 2(k+1) \times 3 \times 3^k - 3^2 \cdot k \cdot 3^{k-1}$$

$$= 2(k+1) \cdot 3^{k+1} - k \cdot 3^{k+1}$$

$$= (2k+2 - k) \cdot 3^{k+1}$$

$$= (k+2) \cdot 3^{k+1}$$

$$= (k+1+1) \cdot 3^{k+1}$$

as required

(Statement Required).

(9)

Question 16

$$\text{let } P = ae^{mx} + be^{-mx}$$

$$\frac{dP}{dx} = mae^{mx} - mbe^{-mx} = 0$$

$$\frac{dP}{dx} = mae^{mx} = be^{-mx}$$

$$\therefore m^2 = -mg - g \frac{v^2}{K^2} \quad m=1$$

$$ae^{mx} = b e^{-mx}$$

$$e^{2mx}$$

$$e^{2mx} = b/a$$

$$2mx = \ln(b/a)$$

$$\sqrt{\frac{dv}{dx}} = -g \left(\frac{K^2 + v^2}{K^2} \right)$$

$$x = \frac{1}{2m} \ln \left(\frac{b}{a} \right)$$

$$\frac{dv}{dx} = -g \left(\frac{K^2 + v^2}{v K^2} \right)$$

$$\frac{dx}{dv} = -\frac{1}{g} \frac{v K^2}{K^2 + v^2}$$

$$\frac{d^2P}{dx^2} = m^2 ae^{mx} + m^2 be^{-mx}$$

$$\text{at } x = \frac{1}{2m} \ln(b/a)$$

$$x = -\frac{K^2}{g} \int \frac{v}{K^2 + v^2} dv$$

$$\frac{d^2P}{dx^2} > 0 \quad \text{as } e^{-mx} > 0$$

$$x = -\frac{K^2}{g} \cdot \frac{1}{2} \ln(K^2 + v^2) + C_1$$

$$\text{and } a, b, m > 0$$

$$x = 0$$

$$v = 2k$$

\therefore min value is when

$$x = \frac{1}{2m} \ln(b/a)$$

$$\therefore C_1 = \frac{K^2}{2g} \ln(5k^2)$$

$$P = m \left(\frac{1}{2m} \ln(b/a) \right) - m \left(\frac{1}{2m} \ln(b/a) \right) \therefore$$

$$ae^{m \left(\frac{1}{2m} \ln(b/a) \right)} + be^{-m \left(\frac{1}{2m} \ln(b/a) \right)} \quad x = -\frac{K^2}{2g} \ln(K^2 + v^2) + \frac{K^2 \ln(5k^2)}{2g}$$

$$= ae^{\frac{1}{2} \ln(b/a)} + b e^{-\frac{1}{2} \ln(b/a)}$$

$$= a e^{\ln \sqrt{b/a}} + b e^{\ln \sqrt{b/a}}$$

$$\max \text{ height } v=0$$

$$= a \sqrt{\frac{b}{a}} + b \sqrt{\frac{a}{b}}$$

$$x = -\frac{K^2}{2g} \ln K^2 + \frac{K^2}{2g} \ln(5k^2)$$

$$= K^2/2g \ln \left[\frac{5k^2}{K^2} \right]$$

$$= \frac{k^2}{2g} \ln 5.$$

$$= \sqrt{ab} + \sqrt{ba} = 2\sqrt{ab}.$$

(10)

Question 16 cont'

b) $x=0 t=0 v=0$ $\frac{k^2}{2g} \ln 5 = -\frac{k^2}{2g} \ln(k^2 - v^2) + \frac{k^2}{2g} \ln(k^2)$

$\downarrow + \downarrow mg$

$(m=1)$

$\uparrow R$

$\ln 5 = -\ln(k^2 - v^2) + \ln k^2$

$\ln 5 = \ln \left(\frac{k^2}{k^2 - v^2} \right)$

$\ddot{x} = g - R$

$\ddot{x} = g - \frac{g}{k^2} v^2$

$-5v^2 = -4k^2$

$v^2 = \frac{4k^2}{5}$

$v dv = g - \frac{gv^2}{k^2}$

$\therefore v = \sqrt{\frac{4k^2}{5}}$

$\frac{dv}{dx} = \frac{g}{v} - \frac{gv}{k^2}$

$v > 0$

$= \frac{gk^2 - gv^2}{vk^2}$

$V = \frac{2k}{\sqrt{5}}$

$\frac{dx}{dv} = \frac{vk^2}{gk^2 - gv^2}$

$x = \int \frac{vk^2}{gk^2 - gv^2} dv$

$x = \frac{k^2}{g} \times -1 \ln(k^2 - v^2) + C_2$

$x=0$

$v=0$

$C_2 = \frac{k^2}{2g} \ln k^2$

$x = -\frac{k^2}{2g} \ln(k^2 - v^2) + \frac{k^2}{2g} \ln k^2$

Now $x = \frac{k^2}{2g} \ln 5$

Question 16 cont'

$\Delta V = \frac{\pi^2}{2} \left[\frac{\pi}{2} - 2(0) + 0 - \left(\frac{\pi}{2} \cdot \frac{1}{\sqrt{2}} - 2 \frac{\pi}{2\sqrt{2}} \frac{-2\pi}{\sqrt{2}} + \frac{2}{\sqrt{2}} \right) \right]$

$x \text{ lies of } y = \cos x$

$\therefore x = \cos^{-1} y$

(I used symmetry can do $\sin^{-1} y$)

$r = x = \cos^{-1} y$

$R = \frac{\pi}{2} - x = \frac{\pi}{2} - \cos^{-1} y$

$\Delta V = \pi (R^2 - r^2) \Delta y$

$= \pi \left[\frac{\pi}{2} - \cos^{-1} y - \cos y \right] \left[\frac{\pi}{2} - \cos^{-1} y + \cos y \right] \Delta y$

$= \pi \left[\frac{\pi}{2} - 2\cos^{-1} y \right] \left[\frac{\pi}{2} \right] \Delta y$

$= \frac{\pi^2}{2} \left[\frac{\pi}{2} - 2\cos^{-1} y \right] \Delta y$

other solutions such as $\frac{\pi^2}{2} \int_{-\frac{\pi}{2}}^1 2\sin^{-1} y - \frac{\pi}{2} dy$ can be used.

Total volume $\ddot{\Sigma}$

$= \lim_{\Delta y \rightarrow 0} \sum_{y \in \Sigma}^1 \frac{\pi^2}{2} \left[\frac{\pi}{2} - 2\cos^{-1} y \right] \Delta y$

$= \frac{\pi^2}{2} \int_{-\frac{\pi}{2}}^1 \frac{\pi}{2} - 2\cos^{-1} y dy$

$= \frac{\pi^2}{2} \left[\frac{\pi}{2} y - \left[2y\cos^{-1} y - \int 2y \cdot \frac{-1}{\sqrt{1-y^2}} dy \right] \right]$

$= \frac{\pi^2}{2} \left[\frac{\pi}{2} y - 2y\cos^{-1} y + 2\sqrt{1-y^2} \right]_0^1$